**A PRIORI ERROR ESTIMATES FOR NUMERICAL METHODS
FOR SCALAR CONSERVATION LAWS
PART III: MULTIDIMENSIONAL FLUX-SPLITTING
MONOTONE SCHEMES ON NON-CARTESIAN GRIDS***

BERNARDO COCKBURN[†], PIERRE-ALAIN GREMAUD[‡], AND JIMMY XIANGRONG
YANG[†]

Abstract. This paper is the third of a series in which a general theory of a priori error estimates for scalar conservation laws is constructed. In this paper, we consider multidimensional flux-splitting monotone schemes defined on non-Cartesian grids. We identify those schemes which are *consistent* and prove that the $L^\infty(0, T; L^1(\mathbb{R}^d))$ -norm of the error goes to zero as $(\Delta x)^{1/2}$ when the discretization parameter Δx goes to zero. Moreover, we show that *nonconsistent* schemes can converge at optimal rates of $(\Delta x)^{1/2}$ because (i) the *conservation form* of the schemes and (ii) the so-called *consistency of the numerical fluxes* allow the regularity properties of the approximate solution to compensate for their lack of *consistency*.

Key words. a priori error estimates, irregular grids, monotone schemes, conservation laws, supraconvergence

AMS subject classifications. 65M60, 65N30, 35L65

PII. S0036142997316165

1. Introduction. This is the third of a series of papers in which we develop a theory of a priori error estimates, that is, estimates given solely in terms of the exact solution, for numerical methods for the scalar conservation law [16]

$$(1.1a) \quad v_t + \nabla \cdot f(v) = 0, \quad \text{in } (0, T) \times \mathbb{R}^d,$$

$$(1.1b) \quad v(0) = v_0, \quad \text{on } \mathbb{R}^d.$$

In the first paper of this series [6], a new general approach for obtaining a priori error estimates for numerical methods for scalar conservation laws was introduced. The approach was then used to obtain optimal a priori error estimates for the Engquist–Osher scheme [9] on one-dimensional uniform grids; in contrast with previous work, [4], [5], [18], [19], [20], [21], [23], [24], [25], [27], [29], [31], (see also [28]), no regularity properties of the approximate solution were *explicitly* used. In the second paper of this series [7], an extension of the above result to the case of flux-splitting monotone schemes defined on irregular Cartesian grids was undertaken. As it is well known, see, for example, [13], [26], and [30], if the numerical fluxes do not properly take into account the irregularity of the grids, a *loss of consistency* is generated and, as a consequence, a new term in the truncation error appears. In [7] it was shown that this new term could be controlled without using regularity properties of the approximate solution *only* for a subset of the schemes which were called *consistent* schemes. For

*Received by the editors February 7, 1997; accepted for publication October 30, 1997; published electronically July 9, 1998.

<http://www.siam.org/journals/sinum/35-5/31616.html>

[†]School of Mathematics, University of Minnesota, 127 Vincent Hall, Minneapolis, MN 55455 (cockburn@math.umn.edu). The first author was partially supported by the National Science Foundation grant DMS-9407952 and by the University of Minnesota Supercomputer Institute.

[‡]Center for Research in Scientific Computation and Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205. This author was partially supported by the Army Research Office through grant DAAH04-95-1-0419.

FIG. 1. Initial condition v_0 and triangulation with $\Delta x = 1/8$.

this *new* class of schemes, optimal a priori error estimates were proven. In this paper, we consider the case of flux-splitting monotone schemes of the form

$$(1.2) \quad u_K^{n+1} = u_K^n - \frac{\Delta t}{|K|} \sum_{e \in \partial K} |e| f_{e,K}(u_K^n, u_{K_e}^n),$$

defined on non-Cartesian grids (see section 2.a), we identify those schemes which are *consistent* and we prove an optimal a priori error estimate for them; more precisely, we show that the $L^\infty(0, T; L^1(\mathbb{R}^d))$ -norm of the error goes to zero as $(\Delta x)^{1/2}$ where Δx is the maximum diameter of the finite volumes K . Moreover, we uncover the mechanism that allows optimal a priori error estimates to hold even for *nonconsistent* schemes. Extending a similar result obtained in the one-dimensional case [7], we show that the regularity properties of the approximate solution can compensate for the loss of *consistency* of the scheme thanks to (i) the *conservation form* of the schemes and to (ii) the so-called *consistency of the numerical fluxes*; the nonlinear nature of the equations does not play any role in this mechanism.

The error estimate presented in this paper is the first a priori error estimate and the first optimal error estimate for numerical schemes for scalar conservation laws defined on non-Cartesian grids. As pointed out above, see also the discussion in [5], all previous work in error estimation for nonlinear conservation laws has been devoted to a posteriori error estimates. As a consequence, upper bounds of the total variation of the approximate solution had to be obtained. Since it is not a trivial matter to obtain such bounds even for simple schemes, this requirement has significantly slowed down the progress in this area. Indeed, there are only a few upper bounds of the total variation available in the literature. In [17] and [8], for monotone schemes in uniform Cartesian grids, and in [27], for monotone schemes in nonuniform Cartesian grids, it was proven that the total variation of the approximate solution does not increase in time. Unfortunately, the techniques used in the above mentioned papers are of no use in the non-Cartesian grids case since in this case the total variation of the approximate solutions *does* increase with time, even when the grids are uniform. For example, consider the simple case of the linear conservation law in two dimensions with $f(v) = (0, b)v$ with initial condition v_0 equal to one in the shaded area shown in Figure 1 and equal to zero elsewhere.

Then for the simple Lax–Friedrichs-like scheme (1.2) with

$$f_{e,K}(v, w) = \frac{1}{2}(f(v) + f(w)) \cdot n_{e,K} - \frac{1}{2\lambda}(w - v),$$

FIG. 2. The ratio $R(N) = |u_h(\Delta t)|_{TV(R^2)} / |u_h(0)|_{TV(R^2)}$.

and grids of equilateral triangles as shown in Figure 1, a computation shows that the ratio R of the total variation of the approximate solution $u_h(\Delta t)$ after a single time-step to the total variation of the initial data $u_h(0) = v_0$ as a function of $\Delta x = 1/N$ is bigger than one, as is shown in Figure 2; therein, we have taken $\lambda b = .9$ and $cfl = 2\sqrt{3} \frac{\Delta t}{\lambda \Delta x} = .9$.

The other technique to obtain upper bounds for the total variation of approximate solutions of numerical schemes for nonlinear conservation laws is an extension, obtained in [4] (see also [15]) of an argument introduced in [2] to obtain the convergence of finite difference methods [3] in the framework of measure-valued solutions. This technique, which is essentially an a priori L^2 -estimate, was used in [4], [31], [25], and [5] to establish that the total variation of the approximate solution blows up like $(\Delta x)^{-1/2}$ as Δx goes to zero. This upper bound results in a posteriori error estimates with a rate of convergence of $(\Delta x)^{1/4}$ for finite volume schemes [4], [31], [25], of $(\Delta x)^{1/4}$ for the discontinuous Galerkin method [5], and of $(\Delta x)^{1/8}$ for the streamline diffusion method [5]. Unfortunately, this technique does not lead to optimal error estimates since the total variation is bounded only in terms of the L^2 -norm of the initial data instead of in terms of its total variation. The advantage of our approach is that no estimate of the total variation is required to obtain optimal error estimates for the so-called *consistent* schemes.

We have to point out, however, that *consistent* schemes of the form (1.2), (i) can only be defined on fairly uniform grids and (ii) have to have fairly restrictive numerical fluxes. For example, upwinding-like schemes are not *consistent*; see a related result in [12]. This unfortunate situation is, of course, related to the fact that the schemes of the form (1.2) update the value at the volume K by *only* using the information stored in those volumes K_e sharing a common face e with K . We believe that the consistency requirements for schemes of a wider stencil, as in, for example, the Lax–Friedrichs-like scheme considered in [1], should be weaker. However, we do not explore this issue in this paper since this would only further complicate an already very technical analysis.

For schemes that are not *consistent*, we show that optimal error estimates can still be obtained in spite of the loss of consistency. This phenomenon, sometimes called *supraconvergence*, has been studied by several authors in different contexts; see, for example, [22], [14], [11], and [10]. *Supraconvergence* for schemes for the linear conservation laws has been studied in [33], [34] for the one-dimensional case, and in [32] for nonuniform Cartesian grids. Therein, the degeneracy from second-order to first-order accuracy, due to the irregularities of the grids, “predicted” by a naive consideration

of the L^∞ -like truncation error is studied. By a careful consideration of the *structure* of that truncation error, it is shown that this degeneracy does not actually take place provided that the scheme is stable and the exact solution is smooth enough. Our approach is similar in that we show that the *structure* of the truncation error allows the stability properties of the scheme to compensate for its lack of consistency. However, we study the general nonlinear conservation laws, work with an L^1 -like truncation error and half-order accurate schemes, require minimal regularity of the exact solution, and consider general non-Cartesian grids.

Finally, let us emphasize that the results presented in this paper are not a straightforward extension of those obtained in our second paper of this series [7] for irregular Cartesian grids. Although the technique to obtain a priori error estimates introduced in [6] is independent of the space dimension, the irregularity of the grids plays a greater role in the multidimensional case and renders the manipulation of the truncation error a much more delicate undertaking.

The paper is organized as follows. In section 2, the numerical schemes under consideration and related technical assumptions are presented, and the main results are stated and discussed. In section 3, we give a proof of our main a priori error estimate, Theorem 2.1. In section 4, we give an explanation of the *supraconvergence* phenomenon and the proof of the corresponding a priori error estimate, Theorem 2.3. We end with some concluding remarks in section 5.

2. The numerical schemes and main results.

2a. The numerical schemes. We consider, for the sake of simplicity, uniform partition of $\mathbb{R}^+$ of the form $\{t^n = n\Delta t\}_{n \in \mathbb{N}}$ and triangulations of $\mathbb{R}^d$, $\mathcal{T}_{\Delta x} = \{K\}$, made of nonoverlapping polyhedra. We require that for any two elements K and K' , $\overline{K} \cap \overline{K}'$ is either a face e of both K and K' with nonzero $(d-1)$ -Lebesgue measure $|e|$ or has Hausdorff dimension less than $d-1$. The maximum diameter of the sets K of the triangulation $\mathcal{T}_{\Delta x}$ is denoted by Δx .

Given these partitions, we define an approximation u to the entropy solution v of (1.1) as the piecewise-constant function

$$(2.1) \quad u(t, x) = u_K^n, \quad \text{for } (t, x) \in [t^n, t^{n+1}) \times K$$

constructed as follows. At $t = 0$, the degrees of freedom of u are given by

$$(2.2a) \quad u_K^0 = \frac{1}{|K|} \int_K u_0(s) ds,$$

where $|K|$ is the measure of the finite volume K . The remaining degrees of freedom are defined by the following flux-splitting scheme:

$$(2.2b) \quad u_K^{n+1} = u_K^n - \frac{\Delta t}{|K|} \sum_{e \in \partial K} |e| f_{e,K}(u_K^n, u_{K_e}^n),$$

where K_e denotes the finite volume sharing the face e with the finite volume K and the numerical flux $f_{e,K}(v, w)$ has the form

$$(2.3a) \quad f_{e,K}(v, w) = f_{cent;e,K}(v, w) - f_{visc;e,K}(v, w),$$

with

$$(2.3b) \quad f_{cent;e,K}(v, w) = a_{e,K} f(v) \cdot n_{e,K} + b_{e,K} f(w) \cdot n_{e,K},$$

TABLE 1
Values of the mesh related parameters on simple grids.

mesh type	aspect ratio	d	$\mathbb{D}_{\Delta x}$	$\iota_{\Delta x}$
equilateral	N/A	2	4	3
Cartesian triangular	λ	2	$2\sqrt{4+\lambda^2}$	3
Cartesian rectangular	λ	2	$\sqrt{4+\lambda^2}$	4
Cartesian rectangular	λ, μ	3	$\sqrt{4+\lambda^2+\mu^2}$	6

where $n_{e,K}$ is the outward unit normal at the face e of the finite volume K and

$$(2.3c) \quad f_{visc;e,K}(v, w) = \alpha_e (N_e(w) - N_e(v)).$$

We assume that the flux $f_{e,K}$ is *consistent* with the nonlinearity $f \cdot n_{e,K}$ and *conservative*, i.e., that $f_{e,K}(u, u) = f(u) \cdot n_{e,K}$ and $f_{e,K}(u_K, u_{K_e}) + f_{e,K_e}(u_{K_e}, u_K) = 0$, respectively. For the schemes under consideration, these properties are equivalent to the following conditions:

$$(2.3d) \quad a_{e,K} + b_{e,K} = 1, \quad (\text{consistency}),$$

$$(2.3e) \quad a_{e,K_e} = b_{e,K}, \quad b_{e,K_e} = a_{e,K}, \quad (\text{conservativity}).$$

We ask that

$$(2.3f) \quad 0 \leq a_{e,K}, b_{e,K} \leq 1.$$

Finally, we require that for every finite volume K and each of its faces e we have

$$(2.3g) \quad \alpha_e N'_e(\cdot) \geq b_{e,K} |f'(\cdot) \cdot n_{e,K}|,$$

and normalize α_e and N'_e by asking that for all bounded intervals I :

$$(2.3h) \quad \alpha_e \leq 1, \quad \sup_{c \in I} N'_e(c) \leq \sup_{c \in I} \|f'(c)\|_{l^\infty}.$$

As it is well known, this condition ensures the *monotonicity* of the scheme under the following condition on the size of the time-step Δt :

$$(2.4) \quad \frac{\Delta t}{|K|} \sum_{e \in \partial K} |e| (\alpha_e N'_e(u_K^n) - b_{e,K} f'(u_K^n) \cdot n_{e,K}) \leq 1, \quad \forall K \in \mathcal{T}_{\Delta x}, n \geq 0.$$

Our results also involve two grid-related quantities which we introduce next. Given a face $e \in \mathcal{E}_{\Delta x} \equiv \bigcup \{e : e \in \partial K \text{ for all } K \in \mathcal{T}_{\Delta x}\}$, we define C_e to be the convex hull of $K_e^+ \cup K_e^-$, d_e to be the diameter of C_e , and we set

$$(2.5) \quad \mathbb{D}_{\Delta x} = \sup_{e \in \mathcal{E}_{\Delta x}} \frac{|e| d_e}{\max\{|K_e^+|, |K_e^-|\}}.$$

Also, given the finite volume K , we denote by ι_K the number of sets C_e for which $|K \cap C_e| \neq 0$ for some $e \in \mathcal{E}_{\Delta x}$ and we set

$$(2.6) \quad \iota_{\Delta x} = \sup_{K \in \mathcal{T}_{\Delta x}} \iota_K.$$

Table 1 gives the values of those parameters in some elementary cases.

2b. Consistency and a priori error estimates.

The generalization of the notion of *consistency* introduced in [7] for schemes on irregular Cartesian grids will be based on the following two quantities:

$$(2.7a) \quad \Delta_K = \max_{1 \leq j \leq d} \sum_{i=1}^d \left| \sum_{e \in \partial K} |e| (\delta_e)_j (n_{e,K})_i \right|,$$

$$(2.7b) \quad \mathbb{A}_K = \sup_{I \in \mathbb{R}} \frac{1}{\sup_{c \in I} \|f'(c)\|} \sup_{c \in I} \sum_{i=1}^d \left| \sum_{e \in \partial K} |e| \alpha_e N'_e(c) (x_{K_e} - x_K)_i \right|,$$

where

$$(2.7c) \quad \delta_e = (x_e - x_K) - b_{e,K_e} (x_{K_e} - x_K),$$

where x_K denotes the barycenter of the finite volume K and x_e the barycenter of the face e . We say that the scheme (2.2b) with fluxes satisfying the conditions (2.3) is *consistent* with respect to the family of triangulations $\{\mathcal{T}_{\Delta x}\}_{\Delta x > 0}$ if $\Delta_K = \mathbb{A}_K \equiv 0$. If we consider, for instance, a grid of equilateral triangles or any affine transformation of it and choose

$$a_{e,K} = b_{e,K} = 1/2, \quad \alpha_e = 1/2, \quad N'_e(c) = \|f'(c)\| \quad \text{or} \quad N'_e(c) = 1/\lambda,$$

then the corresponding scheme is *consistent*. Note that the first choice of N'_e corresponds to an Engquist–Osher-like scheme with isotropic artificial viscosity whereas the second choice corresponds to the Lax–Friedrichs like scheme described in the introduction.

The conditions for *consistency* for the schemes under consideration impose heavy restrictions on the grids and on the numerical fluxes. Indeed, $\Delta_K \equiv 0$ if $\delta_e \equiv 0$, that is, if

$$(x_c - x_K) = b_{e,K_e} (x_{K_e} - x_K),$$

or, equivalently, if

$$x_e = b_{e,K_e} x_{K_e} + b_{e,K} x_K.$$

This means not only that the value of the scalar b_{e,K_e} is determined, but that the barycenters x_{K_e} and x_K and the barycenter of the face e , x_e , must lie on the same line! This condition is trivially satisfied for Cartesian or affine transformations of Cartesian grids, but that is not the case for general non-Cartesian grids. Moreover, $\mathbb{A}_K \equiv 0$ if and only if

$$x_K = \sum_{e \in \partial K} \zeta_{e,K}(c) x_{K_e}, \quad \text{where } \zeta_{e,K}(c) = \frac{|c| \alpha_e N'_e(c)}{\sum_{e \in \partial K} |e| \alpha_e N'_e(c)}.$$

This means, not only that the quantities $\zeta_{e,K}(c)$ should be independent of c , but that the barycenter x_K has to be equal to a very precise convex combination of the barycenters x_{K_e} . This condition is *impossible* to satisfy for upwinding-like fluxes unless the grids are Cartesian or affine transformations of Cartesian grids.

For example, consider the linear case $f(v) = (0, b) v$ in two-space dimensions and triangulations made of acute triangles. For the upwinding scheme $N'_e(c) = |(0, b) \cdot$

$n_{e,K} \mid$. If for some face e of the finite volume K the velocity $(0, b)$ is perpendicular to $n_{e,K}$, we have $N'_e(c) = 0$ and the previous condition is impossible to satisfy. This lack of consistency of upwinding-like schemes has been pointed out in [12] in the framework of a formal L^∞ -like truncation error analysis.

Fortunately, the consistency condition can be relaxed. We say that a scheme is *p-consistent* if there are two nonnegative constants C_δ and C_α such that

$$(2.8) \quad |\delta|_{var,1/2} \leq C_\delta (\Delta x)^p, \quad |\alpha|_{var,1/2} \leq C_\alpha (\Delta x)^p,$$

where

$$(2.9) \quad |\delta|_{var,1/2} = \sup_{x \in \mathbb{R}^d} \frac{\sum_{x: |x-x_K| \leq (\Delta x)^{1/2}} \Delta_K}{(\Delta x)^{d/2}},$$

$$(2.10) \quad |\alpha|_{var,1/2} = \sup_{x \in \mathbb{R}^d} \frac{\sum_{x: |x-x_K| \leq (\Delta x)^{1/2}} \mathbb{A}_K}{(\Delta x)^{d/2}}.$$

For instance, if we consider the two-dimensional case and define

$$b_{e,K} = \frac{(x_e - x_{K_e}) \cdot (x_K - x_{K_e})}{\|x_K - x_{K_e}\|^2}, \quad a_{e,K} = 1 - b_{e,K},$$

$$\alpha_e = \max\{a_{e,K}, a_{e,K_e}\}, \quad N'_e(c) = \|f'(c)\| \quad \text{or} \quad N'_e(c) = 1/\lambda,$$

then the conservativity condition (2.3e) is satisfied. Moreover, (2.3f) is verified provided that

$$\|x_e - x_{K_e}\| \leq \|x_K - x_{K_e}\| \quad \text{and} \quad |\theta_e| \leq \frac{\pi}{2},$$

where $\theta_e = \angle(x_e - x_{K_e}, x_K - x_{K_e})$. In particular, those conditions are satisfied if the triangulation is acute. We observe that

$$\Delta_K \leq 2\iota_{\Delta x} \mathbb{D}_{\Delta x} \sup_{e \in \mathcal{E}_{\Delta x}} \left(\frac{\|\delta_e\|_{l^\infty}}{d_e} \max\{K_e^+, K_e^-\} \right),$$

$$\mathbb{A}_K \leq d_{\iota_{\Delta x}} \Delta x \sup_{e \in \mathcal{E}_{\Delta x}} \left\| x_K - \sum_{e \subset \partial K} \zeta_{e,K}(c) x_{K_e} \right\|_{\ell^\infty}.$$

Assuming now the triangulation to be uniformly regular, i.e., $\Delta x \leq \kappa h_K$ for any $K \subset \mathcal{T}_{\Delta x}$ (h_K = diameter of K), gives

$$|\delta|_{var,1/2} \leq 2\kappa^2 \iota_{\Delta x} \mathbb{D}_{\Delta x} \sup_{e \in \mathcal{E}_{\Delta x}} \frac{\|\delta_e\|_{l^\infty}}{d_e},$$

$$|\alpha|_{var,1/2} \leq 2d_{\iota_{\Delta x}} \frac{\kappa^2}{\Delta x} \sup_{e \in \mathcal{E}_{\Delta x}} \left\| x_K - \sum_{e \subset \partial K} \zeta_{e,K}(c) x_{K_e} \right\|_{\ell^\infty}.$$

Without further assumptions on the grid, we have $|\delta|_{var,1/2} = \mathcal{O}(1)$ and $|\alpha|_{var,1/2} = \mathcal{O}(1)$ and the above scheme is 0-consistent, i.e., inconsistent (case treated in Theorem 2.3). However, if we assume, for instance,

$$\|\delta_e\|_{\ell^\infty} = \mathcal{O}(\Delta x^{1+p}),$$

$$\left\| \sum_{e \in \partial K} (x_{K_e} - x_K) \right\|_{\ell^\infty} = \mathcal{O}(\Delta x^{1+p}),$$

then the scheme is p -consistent (case treated in Corollary 2.2). The first of the above assumptions amounts to having, for $e \in \mathcal{E}_{\Delta x}$, the three nodes x_K , x_e , x_{K_e} lining up as Δx goes to zero. The second assumption means, for instance, that the grid becomes asymptotically equilateral, if simplicial elements are used.

We are now ready to state our error estimate which, following [6], is expressed in terms of the numerical viscosity associated to the scheme under consideration and in terms of the measure of consistency introduced above.

THEOREM 2.1. *Let the Courant–Friedrichs–Levy condition (2.4) be satisfied. Let u be the piecewise-constant solution given by the scheme (2.2) with coefficients satisfying (2.3), let v be the entropy solution. Then,*

$$\begin{aligned} \|u(t^N) - v(t^N)\|_{L^1(\mathbb{R}^d)} &\leq 2 \|u_0 - v_0\|_{L^1(\mathbb{R}^d)} + 8 \|v_0\|_{TV(\mathbb{R}^d)} \sqrt{2t^N \|\nu_v\|} (\Delta x)^{1/2} \\ &\quad + C \|v_0\|_{TV(\mathbb{R}^d)} (\|\delta\|_{var,1/2} + \|\alpha\|_{var,1/2}) \\ &\quad + \|v_0\|_{TV(\mathbb{R}^d)} (b_1(\Delta x)^{3/4} + b_2 \Delta x), \end{aligned}$$

where $\|\nu_v\| = \sup_K \sup_{t \in (0, t^N)} \max_{1 \leq i, j \leq d} |\nu_K^{ij}(v(t, x)) / \Delta x|$ and the local viscosity coefficient ν_K^{ij} is given by

$$\begin{aligned} \nu_K^{ij}(c) &= \frac{\Delta t}{2} f'_i(c) f'_j(c) - \frac{1}{|K|} \sum_{e \in \partial K} |e| [b_{e, K_e} f'(c) \cdot n_{e, K} + \alpha_e N'_e(c)] \xi_{e, K}^{ij}, \\ \xi_{e, K}^{ij} &= \frac{1}{2} \{ (x_{K_e} - x_K)_i (x_{K_e} - x_K)_j + \mathbb{I}_{K_e}^{ij} - \mathbb{I}_K^{ij} \}, \\ \mathbb{I}_\Omega^{ij} &= \frac{1}{|\Omega|} \int_\Omega (x' - x_\Omega)_i (x' - x_\Omega)_j dx'. \end{aligned}$$

The constant C is given by

$$C = 4 \|f'(v)\| \left(t^N + \sqrt{2t^N \|\nu_v\|} \right) (1 + b_0 (\Delta x)^{1/4}),$$

and $\|f'(v)\|$ is given by

$$\|f'(v)\| = \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \|f'(v(t, x))\|_{\ell^\infty}.$$

The constants b_0 , b_1 , and b_2 are locally bounded functions that depend solely on the quantities $\|f'(v)\| \Delta t / \Delta x$, $\|f'(v)\| / \|\nu_v\|$, $\{t^N \|\nu_v\|\}^{1/2}$, $\mathbb{D}_{\Delta x}$, and ι_Δ .

An immediate consequence of this result follows.

COROLLARY 2.2 (p -consistent schemes). *With the notation and under the assumptions of Theorem 2.1, if the scheme is p -consistent, we have*

$$\begin{aligned} \|u(t^N) - v(t^N)\|_{L^1(\mathbb{R}^d)} &\leq 2 \|u_0 - v_0\|_{L^1(\mathbb{R}^d)} + 8 \|v_0\|_{TV(\mathbb{R}^d)} \sqrt{2t^N \|\nu_v\|} (\Delta x)^{1/2} \\ &\quad + \mathcal{O}((\Delta x)^{\min\{p, 3/4\}}). \end{aligned}$$

2c. Supraconvergence and a posteriori error estimates. For *nonconsistent* schemes in arbitrary triangulations, and this includes notably the upwinding-like

schemes mentioned in the previous section, it is possible to obtain an error estimate in terms of the following quantities:

$$(2.11a) \quad \|\delta\|_{\ell^\infty(\mathcal{E}_{\Delta x})/\mathbb{R}^d} = \inf_{\hat{\delta} \in \mathbb{R}^d} \sup_{e \in \mathcal{E}_{\Delta x}} \max_{1 \leq j \leq d} |(\delta_e - \hat{\delta})_j|,$$

$$(2.11b) \quad \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})} = \sup_{e \in \mathcal{E}_{\Delta x}} |\alpha_e| \max_{1 \leq j \leq d} |(x_{K_e} - x_K)_j|,$$

which are always of order Δx , by (2.7c), (2.3f), and (2.3h), instead of in terms of the quantities $|\delta|_{var,1/2}$ and $|\alpha|_{var,1/2}$. The price to pay is that the error estimate now must depend on the smoothness properties of the approximate solution; see Lemma 4.1.

THEOREM 2.3. *Let the Courant–Friedrichs–Levy condition (2.4) be satisfied. Let u be the piecewise-constant solution given by the scheme (2.2) with coefficients satisfying (2.3); let v be the entropy solution. Then,*

$$\begin{aligned} \|u(t^N) - v(t^N)\|_{L^1(\mathbb{R}^d)} &\leq 2 \|u_0 - v_0\|_{L^1(\mathbb{R}^d)} + 8 \|v_0\|_{TV(\mathbb{R}^d)} \sqrt{2t^N} \|\nu_v\| (\Delta x)^{1/2} \\ &+ \frac{2\sqrt{2}d}{\sqrt{t^N} \|\nu_v\| \Delta x} \left\{ \|\delta\|_{\ell^\infty(\mathcal{E}_{\Delta x})/\mathbb{R}^d} + \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})} \right\} \\ &\quad \left\{ \mathbb{D}_{\Delta x} \iota_{\Delta x} \|f'(v)\| t^N \|v_0\|_{TV(\mathbb{R}^d)} + \frac{1}{\iota_{inf}} \|f'(u)\| \|u\|_{L^1(0,t^N;TV(\mathbb{R}^d))} \right\} \\ &+ \|v_0\|_{TV(\mathbb{R}^d)} (b_1(\Delta x)^{3/4} + b_2 \Delta x), \end{aligned}$$

where $\iota_{inf} = \inf_{K \in \mathcal{T}_{\Delta x}} \iota_K$. The constants b_1 and b_2 are locally bounded functions that depend solely on the quantities $\|f'(v)\| \Delta t / \Delta x$, $\|f'(v)\| / \|\nu_v\|$, $\{t^N \|\nu_v\|\}^{1/2}$, $\mathbb{D}_{\Delta x}$, and $\iota_{\Delta x}$, where $\|\nu_v\|$ is defined in Theorem 2.1.

Beside the estimate itself, which is new, an explanation of why it is at all possible is worth considering. As will be shown in section 4, it is a combination of both *consistency* and *conservativity* of the numerical fluxes, which makes it possible by forcing the quantity $D_{e,K} = (x_e - x_K) - b_{e,K_e}(x_{K_e} - x_K)$ to be such that $D_{e,K} = D_{e,K_e} \equiv \delta_e$. Indeed, we have

$$\begin{aligned} D_{e,K} &= (x_e - x_K) - b_{e,K_e}(x_{K_e} - x_K) \\ &= (x_e - x_K) - a_{e,K}(x_{K_e} - x_K), && \text{by conservativity (2.3e),} \\ &= (x_e - x_K) - (1 - b_{e,K})(x_{K_e} - x_K), && \text{by consistency (2.3d),} \\ &= (x_e - x_{K_e}) - b_{e,K}(x_K - x_{K_e}) \\ &= D_{e,K_e}. \end{aligned}$$

3. Proof of Theorem 2.1. This section is entirely devoted to the proof of Theorem 2.1. It has exactly the same structure as section 3 of [7]. In all the proofs, we assume that the entropy solution v is smooth since the general case can be obtained by a standard density argument; see the proof of Proposition 5.5 in [6]. The case in which v is piecewise smooth is treated in [35].

3a. The approximation inequality. Along the lines of [6], we consider the form

$$E(u, v; t^N) = \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \Psi_K^n(v(t, x)) \phi(t, x, t^{n+1}, x_K) |K| \Delta t \, dx \, dt,$$

where $U(\cdot)$ is the absolute value function $|\cdot|$, and Ψ_K^n is given by

$$\Psi_K^n(c) = U'(u_K^n - c) \left\{ \frac{u_K^{n+1} - u_K^n}{\Delta t} + \frac{1}{|K|} \sum_{e \in \partial K} |e| f_{e,K}(u_K^n, u_{K_e}^n) \right\}.$$

The term $\phi(t, x, t^{n+1}, x_K)$ is an averaged test function defined by

$$(3.1) \quad \phi(t, x, t^{n+1}, x_K) = \frac{1}{|K|} \int_K \varphi(t, x, t^{n+1}, x') dx',$$

where x_K denotes the barycenter of the finite volume K , and where the function $\varphi(t, x, t', x')$ is defined as follows:

$$(3.2a) \quad \varphi(t, x, t', x') = w_{\epsilon_t}(t - t') \prod_{i=1}^d \eta_{\epsilon_x}(x_i - x'_i),$$

$x = (x_1, \dots, x_d)$, $x' = (x'_1, \dots, x'_d)$, and t, t' being, respectively, points in $\mathbb{R}^d$ and $\mathbb{R}^+$. Finally, the functions w_{ϵ_t} and η_{ϵ_x} are constructed as follows:

$$(3.2b) \quad w_{\epsilon_t}(s) = \frac{1}{\epsilon_t} w\left(\frac{s}{\epsilon_t}\right), \quad \eta_{\epsilon_x}(s) = \frac{1}{\epsilon_x} \eta\left(\frac{s}{\epsilon_x}\right), \quad \forall s \in \mathbb{R},$$

where ϵ_t and ϵ_x are two arbitrary positive numbers. Both w and η satisfy the following conditions: (i) $\eta(t) \geq 0$, for $t > 0$, (ii) $\eta(t) = \eta(-t)$, for $t > 0$, (iii) the support of η is $[-1, 1]$, and (iv) $\int_0^1 \eta(r) dr = 1/2$. For future reference, we set

$$(3.2c) \quad W(t) = \int_0^t w_{\epsilon_t}(s) ds.$$

Furthermore, the functions w and η are taken to be smooth approximations to $\chi \equiv \chi_0$ and χ_ϵ , where

$$\chi_\epsilon(x) = \begin{cases} (1 + \epsilon)/2 & \text{for } |x| \leq (1 - \epsilon)/(1 + \epsilon), \\ (1 + \epsilon)^2 (1 - |x|)/4\epsilon & \text{for } |x| \in [(1 - \epsilon)/(1 + \epsilon), 1], \\ 0 & \text{elsewhere.} \end{cases}$$

It is easy to verify that we can find a sequence of functions η such that

$$(3.3a) \quad \lim_{\eta \rightarrow \chi_\epsilon} |\eta|_{TV(\mathbb{R})} = |\chi_\epsilon|_{TV(\mathbb{R})} = 1 + \epsilon,$$

$$(3.3b) \quad \lim_{\eta \rightarrow \chi_\epsilon} |\eta'|_{TV(\mathbb{R})} = |\chi'_\epsilon|_{TV(\mathbb{R})} = 2 + \epsilon + 1/\epsilon,$$

$$(3.3c) \quad \lim_{\eta \rightarrow \chi_\epsilon} \|\eta\|_{L^\infty(\mathbb{R})} = \|\chi_\epsilon\|_{L^\infty(\mathbb{R})} = (1 + \epsilon)/2.$$

We now describe the main ideas on which the derivation of our estimates is based. From the identity

$$0 = E(u, v; t^N) = -E_{div}^*(u, v; t^N) + E_{diss}(u, v; t^N) + T_{err}(u, v; t^N),$$

and the structure of the term $T_{err}(u, v; t^N)$, the following approximation inequality can be derived [6, Proposition 7.6]:

$$\begin{aligned} e(t^N) &\leq 2e(0) + 8(\epsilon_x + \epsilon_t \|f'(v)\|) |v_0|_{TV(\mathbb{R}^d)} + 2\|f'(v)\| |v_0|_{TV(\mathbb{R}^d)} \Delta t \\ &\quad + 2 \lim_{w \rightarrow \chi} \sup_{1 \leq n \leq N} \{E_{div}^*(u, v; t^n)/W(t^n) - E_{diss}(u, v; t^n)/W(t^n)\}, \end{aligned}$$

where $e(t)$ denotes the error $\|u(t) - v(t)\|_{L^1(\mathbb{R}^d)}$. Now, to obtain the error estimate, it is enough to estimate the forms $E_{div}^*(u, v; t^n)$ and $E_{diss}(u, v; t^n)$. The remainder of this section is devoted to identifying these forms and estimating them.

We start by rewriting $\Psi_K^n(c)$,

$$\begin{aligned}\Psi_K^n(c) &= U'(u_K^n - c) \frac{1}{\Delta t} \left\{ u_K^{n+1} - u_K^n + \frac{\Delta t}{|K|} \sum_{e \in \partial K} |e| f_{e,K}(u_K^n, u_{K_e}^n) \right\} \\ &= U'(u_K^n - c) \frac{1}{\Delta t} \left\{ u_K^{n+1} - u_K^n + \frac{\Delta t}{|K|} \sum_{e \in \partial K} (p_{e,K}(u_K^n) - p_{e,K}(u_{K_e}^n)) \right\},\end{aligned}$$

where

$$p_{e,K}(s) = -|e| [b_{e,K} f(s) \cdot n_{e,K} - \alpha_e N_e(s)],$$

and proceed as in [6]. Here, the form $E_{diss}(u, v; t^N)$ is

$$E_{diss}(u, v; t^N) = \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} LRED_K^n(v(t, x)) \phi(t, x, t^{n+1}, x_K) |K| \Delta t \, dx \, dt,$$

where the local rate of entropy dissipation $LRED_K^n(c)$ is given by

$$\begin{aligned}LRED_K^n(c) &= \frac{1}{\Delta t} \int_{u_K^{n+1}}^{u_K^n} (p_K(u_K^n) - p_K(s)) U''(s - c) \, ds \\ &\quad + \frac{1}{|K|} \sum_{e \in \partial K} \int_{u_K^{n+1}}^{u_{K_e}^n} (p_{e,K}(u_{K_e}^n) - p_{e,K}(s)) U''(s - c) \, ds,\end{aligned}$$

and $p_K(s) = s - \frac{\Delta t}{|K|} \sum_{e \in \partial K} p_{e,K}(s)$.

The dual form $E_{div}^*(u, v; t^N)$ reads

$$\begin{aligned}E_{div}^*(u, v; t^N) &= - \int_0^{t^N} \int_{\mathbb{R}^d} \int_0^{t^N} \int_{\mathbb{R}^d} U(u(t', x') - v(t, x)) \varphi_t(t, x, t', x') \, dx \, dt \, dx' \, dt' \\ &\quad + \int_0^{t^N} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} U(u(t', x') - v(t^N, x)) \varphi(t^N, x, t', x') \, dx \, dx' \, dt' \\ &\quad - \int_0^{t^N} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} U(u(t', x') - v_0(x)) \varphi(0, x, t', x') \, dx \, dx' \, dt' \\ &\quad - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{e \in \partial K} |e| b_{e,K} [\phi(t, x, t^{n+1}, x_K) \\ &\quad \quad \quad - \phi(t, x, t^{n+1}, x_{K_e})] F(u_K^n, v) \cdot n_{e,K} \Delta t \, dx \, dt \\ &\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{e \in \partial K} |e| \alpha_e [\phi(t, x, t^{n+1}, x_{K_e}) \\ &\quad \quad \quad - \phi(t, x, t^{n+1}, x_K)] \mathcal{N}_e(u_K^n, v) \Delta t \, dx \, dt,\end{aligned}$$

where the functions $F(u, c)$ and $\mathcal{N}_e(u, c)$ are defined as follows:

$$(3.4) \quad F(u, c) = \int_c^u f'(s) U'(s - c) \, ds, \quad \mathcal{N}_e(u, c) = \int_c^u N_e'(s) U'(s - c) \, ds.$$

3b. Estimate of $E_{diss}(u, v; t^n)$.

PROPOSITION 3.1. *Under condition (2.3g) on the viscosity term N , and if the Courant–Friedrichs–Levy (CFL) condition (2.4) is satisfied, the local rate of entropy dissipation $LRED_j^n(c)$ is nonnegative. Hence,*

$$-E_{diss}(u, v; t^n) \leq 0.$$

Sketch of the proof. The above conditions ensure that the functions $p_{e,K}(s)$ and $p_K(s)$ are nondecreasing in s . The result follows from this fact and the definition of $E_{diss}(u_h, v; t^n)$. $\square$

3c. Estimate of $E_{div}^*(u, v; t^N)$.

PROPOSITION 3.2. *We have*

$$\lim_{w \rightarrow \chi} \sup_{1 \leq n \leq N} \{E_{div}^*(u, v; t^n)/W(t^n)\} \leq TEW_{visc} + TEW_{cons} + TEW_{h.o.t.},$$

where

$$\begin{aligned} TEW_{visc}(u, v; t^n) &\leq C_0 t^N d \left\{ 2 \frac{|\eta|_{TV(\mathbb{R})}}{\epsilon_x} \left(1 + \frac{\Delta t}{\epsilon_t} \right) \right\} \|\nu_v\| \Delta x, \\ TEW_{cons}(u, v; t^n) &\leq 2 C_1 t^N \left(1 + \frac{\Delta t}{\epsilon_t} \right) \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d} \right) \|\eta\|_{L^\infty(\mathbb{R})}^d (|\delta|_{var,1/2} + |\alpha|_{var,1/2}), \\ TEW_{h.o.t.}(u, v; t^n) &\leq C_1 \left\{ \frac{1}{2} \|f'(v)\| \frac{(\Delta t)^2 d |\eta|_{TV(\mathbb{R})}}{\epsilon_t \epsilon_x} t^N + 4 \left(1 + \frac{\Delta t}{\epsilon_t} \right) \Delta t \right. \\ &\quad \left. + 32 \mathbb{D}_{\Delta x} \iota_{\Delta x} (\Delta x)^2 \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \left(1 + \frac{\Delta t}{\epsilon_t} \right) t^N \right\}. \end{aligned}$$

where $C_0 = |v_0|_{TV(\mathbb{R}^d)}$ and $C_1 = C_0 \|f'(v)\|$.

To prove this result, we proceed in several steps.

First step: Relating the dual form $E_{div}^*(u, v; t)$ to the truncation error.

We have the following upper bound for $E_{div}^*(u_h, v; t^N)$.

LEMMA 3.3. *We have*

$$\begin{aligned} E_{div}^*(u_h, v; t^N) &\leq \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} F(u_K^n, v(t, x)) \cdot \nabla_x \bar{\phi}(t, x, t^{n+1}, x_K) |K| \Delta t \, dx \, dt \\ &\quad - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{e \in \partial K} |e| b_{e,K_e} [\phi(t, x, t^{n+1}, x_K) \\ &\quad \quad - \phi(t, x, t^{n+1}, x_{K_e})] F(u_K^n, v(t, x)) \cdot n_{e,K} \Delta t \, dx \, dt \\ &\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{e \in \partial K} |e| \alpha_e [\phi(t, x, t^{n+1}, x_{K_e}) \\ &\quad \quad - \phi(t, x, t^{n+1}, x_K)] \mathcal{N}_e(u_K^n, v(t, x)) \Delta t \, dx \, dt. \end{aligned}$$

To prove this result, we use the fact that v is the entropy solution and make some algebraic manipulations; see the proof of the similar result [6, Lemma 7.9].

Next, a relation between the functions $\phi(x, x_{K_e})$ and $\phi(x, x_K)$, defined by (3.1), is needed. That relation is displayed in the following lemma where, for the sake of

clarity, we drop the dependence on t and t' . This simple result is a cornerstone of the proof of Theorem 2.1.

LEMMA 3.4. *We have*

$$\phi(x, x_{K_e}) - \phi(x, x_K) = -(x_{K_e} - x_K) \cdot \nabla_x \phi(x, x_K) + \overline{H}_{e,K}(x, x_K) + \overline{\overline{H}}_{e,K}(x),$$

where

$$\begin{aligned} \overline{H}_{e,K}(x, x_K) &= \sum_{i,j=1}^d \xi_{e,K}^{ij} \partial_{x_i x_j}^2 \phi(x, x_K), \\ \xi_{e,K}^{ij} &= \frac{1}{2} \{ (x_{K_e} - x_K)_i (x_{K_e} - x_K)_j + \mathbb{I}_{K_e}^{ij} - \mathbb{I}_K^{ij} \}, \\ \mathbb{I}_\Omega^{ij} &= \frac{1}{|\Omega|} \int_\Omega (x' - x_\Omega)_i (x' - x_\Omega)_j dx', \end{aligned}$$

and

$$\begin{aligned} \overline{\overline{H}}_{e,K}(x) &= \frac{1}{|K_e|} \int_{K_e} \frac{1}{|K|^3} \int_{K^3} \int_0^1 \int_0^1 \int_0^1 \Psi(x) d\lambda d\mu d\nu dz' dy' dw' dx', \\ \Psi(x) &= \sum_{i,j,k=1}^d \zeta^{ijk} \partial_{x_i x_j x_k}^3 \varphi(x, \lambda(\mu(\nu x' + (1-\nu)w') + (1-\mu)y') + (1-\lambda)z'), \\ \zeta^{ijk} &= (x' - w')_i (\nu x' + (1-\nu)w' - y')_j (\mu(\nu x' + (1-\nu)w') + (1-\mu)y' - z')_k. \end{aligned}$$

Proof of Lemma 3.4. Since

$$\varphi(x, x') = \varphi(x, w') - \int_0^1 (x' - w') \cdot \nabla_x \varphi(x, \nu x' + (1-\nu)w') d\nu,$$

by the definition of φ (3.2), averaging the above equality over K with respect to w' yields

$$\varphi(x, x') = \phi(x, x_K) - \frac{1}{|K|} \int_K \int_0^1 (x' - w') \cdot \nabla_x \varphi(x, \nu x' + (1-\nu)w') d\nu dw',$$

by the definition of ϕ , (3.1). Using the above relation recursively twice, we easily obtain

$$\varphi(x, x') = \phi(x, x_K) + \sum_{i=1}^d A^i(x') \partial_{x_i} \phi(x, x_K) + \sum_{i,j=1}^d B^{ij}(x') \partial_{x_i x_j}^2 \phi(x, x_K) + Res(x, x'),$$

where

$$\begin{aligned} A^i(x') &= -\frac{1}{|K|} \int_K (x' - w')_i dw', \\ B^{ij}(x') &= \frac{1}{|K|} \int_K \int_0^1 (x' - w')_i \left\{ \frac{1}{|K|} \int_K (\nu x' + (1-\nu)w' - y')_j dy' \right\} d\nu dw', \\ Res(x, x') &= \frac{1}{|K|^3} \int_{K^3} \int_0^1 \int_0^1 \int_0^1 \Psi(x) d\lambda d\mu d\nu dz' dy' dw'. \end{aligned}$$

Averaging the above equation K_e with respect to x' and using the definition of ϕ , (3.1), yields

$$\phi(x, x_{K_e}) = \phi(x, x_K) + \sum_{i=1}^d \mathbb{A}^i \partial_{x_i} \phi(x, x_K) + \sum_{i,j=1}^d \mathbb{B}^{ij} \partial_{x_i x_j}^2 \phi(x, x_K) + \overline{\overline{H}}_{e,K}(x),$$

where

$$\mathbb{A}^i = \frac{1}{|K_e|} \int_{K_e} A^i(x') dx', \quad \mathbb{B}^{ij} = \frac{1}{|K_e|} \int_{K_e} B^{ij}(x') dx'.$$

The result follows if we show that $\mathbb{A}^i = -(x_{K_e} - x_K)_i$ and $\mathbb{B}^{ij} = \xi_{e,K}^{ij}$. But, since $A^i(x') = -(x' - x_K)_i$, we do have that $\mathbb{A}^i = -(x_{K_e} - x_K)_i$, and since

$$\begin{aligned} B^{ij}(x') &= \frac{1}{|K|} \int_K \int_0^1 (x' - w')_i (\nu x' + (1 - \nu)w' - x_K)_j d\nu dw' \\ &= \frac{1}{|K|} \int_K (x' - w')_i ((x' + w')/2 - x_K)_j dw' \\ &= \frac{1}{2|K|} \int_K ((x' - x_K) - (w' - x_K))_i ((x' - x_K) + (w' - x_K))_j dw' \\ &= \frac{1}{2} \{ (x' - x_K)_i (x' - x_K)_j - \mathbb{I}_K^{ij} \}, \end{aligned}$$

we have

$$\begin{aligned} \mathbb{B}^{ij} &= \frac{1}{2} \left\{ \frac{1}{|K_e|} \int_{K_e} (x' - x_K)_i (x' - x_K)_j dx' - \mathbb{I}_K^{ij} \right\} \\ &= \frac{1}{2} \left\{ (x_{K_e} - x_K)_i (x_{K_e} - x_K)_j + \frac{1}{|K_e|} \int_{K_e} (x' - x_{K_e})_i (x' - x_{K_e})_j dx' - \mathbb{I}_K^{ij} \right\} \\ &= \frac{1}{2} \{ (x_{K_e} - x_K)_i (x_{K_e} - x_K)_j + \mathbb{I}_{K_e}^{ij} - \mathbb{I}_K^{ij} \} = \xi_{e,K}^{ij}. \end{aligned}$$

This completes the proof. $\square$

With the above lemma, we can now rewrite the upper bound of $E_{div}^*(u_h, v; t^N)$ as follows.

LEMMA 3.5. *We have*

$$E_{div}^*(u_h, v; t^N) \leq TE_{visc}(u, v; t^N) + TE_{cons}(u, v; t^N) + TE_{h.o.t.}(u, v; t^N),$$

where

$$\begin{aligned}
TE_{visc}(u, v; t^N) &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} VISC_K^n(v(t, x); t, x) \, dx \, dt \, \Delta t \\
&\quad - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_{\mathbb{R}^d} F(u_K^n, v(t^N, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(t^N, x, t^{n+1}, x_K) \, dx \, |K| \, \Delta t \\
&\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_{\mathbb{R}^d} F(u_K^n, v(t^0, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(0, x, t^{n+1}, x_K) \, dx \, |K| \, \Delta t, \\
TE_{cons}(u, v; t^N) &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} CONS_K^n(v(t, x); t, x) \, dx \, dt \, \Delta t, \\
TE_{h.o.t.}(u, v; t^N) &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} HOT_K^n(v(t, x); t, x) \, dx \, dt \, \Delta t \\
&\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_{\mathbb{R}^d} F(u_K^n, v(t^N, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(t^N, x, t^{n+1}, x_K) \, dx \, |K| \, \Delta t \\
&\quad - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_{\mathbb{R}^d} F(u_K^n, v(t^0, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(0, x, t^{n+1}, x_K) \, dx \, |K| \, \Delta t.
\end{aligned}$$

The ‘viscosity’ term $VISC_K^n(c; t, x)$ is given by

$$VISC_K^n(c; t, x) = \xrightarrow{time} VISC_K^n(c; t, x) + \xrightarrow{cent} VISC_K^n(c; t, x) + \xrightarrow{visc} VISC_K^n(c; t, x),$$

where

$$\begin{aligned}
\xrightarrow{time} VISC_K^n(c; t, x) &= F(u_K^n, c) \cdot \frac{\Delta t}{2} |K| \nabla_x \phi(t, x, t^{n+1}, x_K), \\
\xrightarrow{cent} VISC_K^n(c; t, x) &= F(u_K^n, c) \cdot \sum_{e \in \partial K} |e| b_{e, K_e} \bar{H}_{e, K}(t, x, t^{n+1}, x_K) n_{e, K}, \\
\xrightarrow{visc} VISC_K^n(c; t, x) &= \sum_{e \in \partial K} |e| \alpha_e \bar{H}_{e, K}(t, x, t^{n+1}, x_K) \mathcal{N}_e(u_K^n, c).
\end{aligned}$$

The ‘consistency’ term $CONS_K^n(c; t, x)$ is given by

$$CONS_K^n(c; t, x) = \xrightarrow{cent} CONS_K^n(c; t, x) + \xrightarrow{visc} CONS_K^n(c; t, x),$$

where

$$\begin{aligned}
\xrightarrow{cent} CONS_K^n(c; t, x) &= F(u_K^n, c) \cdot \sum_{e \in \partial K} |e| \nabla_x \phi(t, x, t^{n+1}, x_K) \cdot \delta_e n_{e, K}, \\
\xrightarrow{visc} CONS_K^n(c; t, x) &= - \sum_{e \in \partial K} |e| \alpha_e (x_{K_e} - x_K) \cdot \nabla_x \phi(t, x, t^{n+1}, x_K) \mathcal{N}_e(u_K^n, c).
\end{aligned}$$

Finally, the ‘high-order’ term $HOT_K^n(c; t, x)$ is given by

$$HOT_K^n(c; t, x) = \xrightarrow{time} HOT_K^n(c; t, x) + \xrightarrow{cent} HOT_K^n(c; t, x) + \xrightarrow{visc} HOT_K^n(c; t, x),$$

where

$$\begin{aligned} \xrightarrow{time} HOT_K^n(c; t, x) &= F(u_K^n, c) \cdot \left[|K| \nabla_x \bar{\phi}(t, x, t^{n+1}, x_K) - |K| \nabla_x \phi(t, x, t^{n+1}, x_K) \right. \\ &\quad \left. - \frac{\Delta t}{2} |K| \nabla_x \phi_t(t, x, t^{n+1}, x_K) \right], \\ \xrightarrow{cent} HOT_K^n(c; t, x) &= F(u_K^n, c) \cdot \sum_{e \in \partial K} |e| b_{e, K_e} \bar{\bar{H}}_{e, K}(t, x, t^{n+1}) n_{e, K}, \\ \xrightarrow{visc} HOT_K^n(c; t, x) &= \sum_{e \in \partial K} |e| \alpha_e \bar{\bar{H}}_{e, K}(t, x, t^{n+1}) \mathcal{N}_e(u_K^n, c). \end{aligned}$$

Proof of Lemma 3.5. The result can be easily obtained by inserting the expression for $\phi(x, x_{K_e}) - \phi(x, x_K)$ of Lemma 3.4 into the upper bound for the dual form $E^*(u, v; t^N)$ in Lemma 3.3 and rearranging terms.

The only manipulation that deserves mentioning is the following. After the above mentioned rearrangement, we get for $\xrightarrow{cent} CONS_K^n(c; t, x)$ the expression

$$F(u_K^n, c) \cdot \left[|K| \nabla_x \phi(t, x, t^{n+1}, x_K) - \sum_{e \in \partial K} |e| b_{e, K_e} (x_{K_e} - x_K) \cdot \nabla_x \phi(t, x, t^{n+1}, x_K) n_{e, K} \right].$$

This expression can be rewritten as

$$F(u_K^n, c) \cdot \sum_{e \in \partial K} |e| \nabla_x \phi(t, x, t^{n+1}, x_K) \cdot [(x_e - x_K) - b_{e, K_e} (x_{K_e} - x_K)] n_{e, K},$$

or, equivalently, see the definition of δ_e , (2.7c), as

$$F(u_K^n, c) \cdot \sum_{e \in \partial K} |e| \nabla_x \phi(t, x, t^{n+1}, x_K) \cdot \delta_e n_{e, K},$$

by using the identity

$$|K| m = \sum_{e \in \partial K} |e| [m \cdot (x_e - x_K)] n_{e, K},$$

with $m = \nabla_x \phi(t, x, t^{n+1}, x_K)$. The above identity can be proved as follows:

$$\begin{aligned} |K| m &= \int_K m dx' = \int_K \nabla_{x'} [m \cdot (x' - x_K)] dx' \\ &= \sum_{e \in \partial K} \int_e [m \cdot (x' - x_K)] n_{e, K} d\Gamma(x') \\ &= \sum_{e \in \partial K} |e| [m \cdot (x_e - x_K)] n_{e, K}. \end{aligned}$$

This completes the proof. $\square$

Second step: Estimating $TE_{visc}(u, v; t^N)$. In this section, we prove the following result.

LEMMA 3.6. *We have*

$$TE_{visc}(u, v; t^N) \leq 2C_0 \frac{d|\eta|_{TV(\mathbb{R})}}{\epsilon_x} \left(1 + \frac{\Delta t}{\epsilon_t}\right) \|\nu_v\| \Delta x,$$

where $C_0 = t^N |v_0|_{TV(\mathbb{R}^d)} W(t^N)$.

Proof of Lemma 3.6. We have, by the definition of $TE_{visc}(u, v; t^N)$ in Lemma 3.5,

$$TE_{visc}(u, v; t^N) = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \Xi(t^{n+1}, x_K) |K| \Delta t,$$

where

$$\begin{aligned} \Xi(t^{n+1}, x_K) = & \int_0^{t^N} \int_{\mathbb{R}^d} \left[F(u_K^n, v(t, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(t, x, t^{n+1}, x_K) \right. \\ & + F(u_K^n, v(t, x)) \cdot \frac{1}{|K|} \sum_{e \in \partial K} |e| b_{e, K_e} \overline{H}_{e, K}(t, x, t^{n+1}, x_K) n_{e, K} \\ & + \frac{1}{|K|} \sum_{e \in \partial K} |e| \alpha_e \overline{H}_{e, K}(t, x, t^{n+1}, x_K) \mathcal{N}_e(u_K^n, v(t, x)) \Big] dx dt \\ & - \int_{\mathbb{R}^d} F(u_K^n, v(t^N, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(t^N, x, t^{n+1}, x_K) dx \\ & + \int_{\mathbb{R}^d} F(u_K^n, v(t^0, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(0, x, t^{n+1}, x_K) dx. \end{aligned}$$

A couple of integrations by parts yield

$$\begin{aligned} \Xi(t^{n+1}, x_K) = & - \int_0^{t^N} \int_{\mathbb{R}^d} \left\{ F_t(u_K^n, v(t, x)) \cdot \frac{\Delta t}{2} \nabla_x \phi(t, x, t^{n+1}, x_K) \right. \\ & + \frac{1}{|K|} \sum_{e \in \partial K} |e| b_{e, K_e} \sum_{i, j=1}^d \partial_{x_i} [F(u_K^n, v(t, x)) \cdot n_{e, K}] \xi_{e, K}^{ij} \partial_{x_j} \phi(t, x, t^{n+1}, x_K) \\ & \left. + \frac{1}{|K|} \sum_{e \in \partial K} |e| \alpha_e \sum_{i, j=1}^d \partial_{x_i} \mathcal{N}_e(u_K^n, v(t, x)) \xi_{e, K}^{ij} \partial_{x_j} \phi(t, x, t^{n+1}, x_K) \right\} dx dt. \end{aligned}$$

Then, by using the definition of F and $\mathcal{N}$, (3.4), we get

$$\Xi(t^{n+1}, x_K) = - \sum_{i, j=1}^d \int_0^{t^N} \int_{\mathbb{R}^d} U'(u_K^n - v) \partial_{x_i} v \nu_K^{ij}(v(t, x)) \partial_{x_j} \phi(t, x, t^{n+1}, x_K) dx dt,$$

where the entries of the matrix ν are

$$\nu_K^{ij}(c) = \frac{\Delta t}{2} f'_i(c) f'_j(c) - \frac{1}{|K|} \sum_{e \in \partial K} |e| [b_{e, K_e} f'(c) \cdot n_{e, K} + \alpha_e N'_e(c)] \xi_{e, K}^{ij}.$$

We have now

$$TE_{visc}(u, v; t^N) \leq T_{aux} |v_0|_{TV(\mathbb{R}^d)} t^N,$$

where

$$\begin{aligned} T_{aux} = & \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \left\{ \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \max_{1 \leq i \leq d} \sum_{j=1}^d \nu_K^{ij}(v(t, x)) \partial_{x_j} \phi(t, x, t^{n+1}, x_K) |K| \Delta t \right\} \\ \leq & \frac{d |\eta|_{TV(\mathbb{R})}}{\epsilon_x} 2 \left(1 + \frac{\Delta t}{\epsilon_t} \right) W(t^N) \|\nu_v\| \Delta x, \end{aligned}$$

where we have made use of Lemma 3.7 of [7], the definition of ϕ , (3.1), and the definition of $\|\nu_v\|$. This completes the proof. $\square$

Third step: Estimating $TE_{cons}(u, v; t^N)$.

LEMMA 3.7. *We have*

$$TE_{cons}(u, v; t^N) \leq 2C_1 \left(1 + \frac{\Delta t}{\epsilon_t}\right) \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d}\right) \|\eta\|_{L^\infty(\mathbb{R})} (|\delta|_{var, 1/2} + |\alpha|_{var, 1/2}),$$

where $C_1 = t^N |v_0|_{TV(\mathbb{R}^d)} \|f'(v)\| W(t^N)$.

Proof of Lemma 3.7. By definition of the term $TE_{cons}(u, v; t^N)$ in Lemma 3.5, we can write

$$TE_{cons}(u, v; t^N) = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \nabla_x \phi(t, x, t^{n+1}, x_K) \cdot A_K(u_K^n, v(t, x)) dx dt \Delta t,$$

where

$$A_K(u, c) = \sum_{e \in \partial K} |e| (F(u, c) \cdot n_{e,K} \delta_e - \alpha_e \mathcal{N}_e(u, c) (x_{K_e} - x_K)).$$

After a simple integration by parts, we obtain

$$\begin{aligned} TE_{cons}(u, v; t^N) &= - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \phi(t, x, t^{n+1}, x_K) \nabla_x v(t, x) \cdot \partial_v A_K(u_K^n, v(t, x)) dx dt \Delta t \\ &\leq \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \phi(t, x, t^{n+1}, x_K) \sum_{i=1}^d |\partial_{x_i} v(t, x)| B_K(v(t, x)) dx dt \Delta t, \end{aligned}$$

where

$$B_K(c) = \max_{1 \leq i \leq d} \left| \sum_{e \in \partial K} |e| (f'_i(c) \cdot n_{e,K} \delta_e - \alpha_e N'_e(c) (x_{K_e} - x_K)) \right|.$$

Hence,

$$TE_{cons}(u, v; t^N) \leq T_{aux} t^N |v_0|_{TV(\mathbb{R}^d)},$$

where

$$\begin{aligned} T_{aux} &= \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \phi(t, x, t^{n+1}, x_K) B_K(v(t, x)) \Delta t \\ &= \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \sum_{n=0}^{N-1} \sum_{K: |x - x_K| \leq \epsilon_x} \phi(t, x, t^{n+1}, x_K) B_K(v(t, x)) \Delta t \\ &\leq \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \left\{ \sum_{n=0}^{N-1} \sup_{x \in \mathbb{R}^d} \phi(t, x, t^{n+1}, x_K) \Delta t \right\} \Lambda(\epsilon_x), \end{aligned}$$

with

$$\Lambda(\epsilon_x) = \sup_{\substack{t \in (0, t^n) \\ x \in \mathbb{R}^d}} \left\{ \sum_{K: |x - x_K| \leq \epsilon_x} B_K(v(t, x)) \right\}.$$

Now, by Lemma 3.7 of [7], we have

$$\sup_{\substack{t \in (0, t^n) \\ x \in \mathbb{R}^d}} \left\{ \sum_{n=0}^{N-1} \sup_{x \in \mathbb{R}^d} \phi(t, x, t^{n+1}, x_K) \Delta t \right\} \leq \frac{\|\eta\|_{L^\infty(\mathbb{R})}^d}{\epsilon_x^d} 2 \left(1 + \frac{\Delta t}{\epsilon_t} \right) W(t^N);$$

observing also

$$\begin{aligned} \Lambda(\epsilon_x) &\leq \epsilon_x^d \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d} \right) \Lambda((\Delta x)^{1/2}) \\ &\leq \epsilon_x^d \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d} \right) \|f'(v)\| (|\delta|_{var, 1/2} + |\alpha|_{var, 1/2}), \end{aligned}$$

and using the definition of $|\delta|_{var, 1/2}$ and $|\alpha|_{var, 1/2}$, (2.9) and (2.10), we get

$$T_{aux} \leq 2 \left(1 + \frac{\Delta t}{\epsilon_t} \right) W(t^N) \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d} \right) \|\eta\|_{L^\infty(\mathbb{R})}^d (|\delta|_{var, 1/2} + |\alpha|_{var, 1/2}),$$

and the results follow. $\square$

Fourth step: Estimating $TE_{h.o.t.}(u, v; t^N)$.

LEMMA 3.8. Suppose that the conditions (2.3) are satisfied. Then,

$$\begin{aligned} TE_{h.o.t.}(u, v; t^N) &\leq C_0 \left\{ \frac{1}{2} \|f'(v)\| \frac{(\Delta t)^2 d \|\eta\|_{TV(\mathbb{R})}}{\epsilon_t \epsilon_x} t^N + 4 \left(1 + \frac{\Delta t}{\epsilon_t} \right) \Delta t \right. \\ &\quad \left. + 32 \mathbb{D}_{\Delta x} \iota_{\Delta x} (\Delta x)^2 \left\{ \frac{d^2 \|\eta\|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d \|\eta'\|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \left(1 + \frac{\Delta t}{\epsilon_t} \right) t^N \right\}, \end{aligned}$$

where $C_0 = |v_0|_{TV(\mathbb{R}^d)} \|f'(v)\| W(t^N)$.

To prove the above result, we rewrite $TE_{h.o.t.}(u, v; t^N)$ as the sum $\xrightarrow{time} TE_{h.o.t.}(u, v; t^N) + \xrightarrow{cent} TE_{h.o.t.}(u, v; t^N) + \xrightarrow{visc} TE_{h.o.t.}(u, v; t^N)$, with the obvious notation, see Lemma 3.5, and estimate each of the above three terms.

LEMMA 3.9. We have

$$\xrightarrow{time} TE_{h.o.t.}(u, v; t^N) \leq C_0 \left\{ \frac{1}{2} \|f'(v)\| \frac{(\Delta t)^2 d \|\eta\|_{TV(\mathbb{R})}}{\epsilon_t \epsilon_x} t^N + 4 \left(1 + \frac{\Delta t}{\epsilon_t} \right) \Delta t \right\},$$

where $C_0 = |v_0|_{TV(\mathbb{R}^d)} \|f'(v)\| W(t^N)$.

Proof. Integrating by parts in time and inserting the definition of ϕ , (3.1), leads to

$$\xrightarrow{time} TE_{h.o.t.}(u, v; t^N) = \sum_{n=0}^{N-1} \int_{\mathbb{R}^d} \Theta(t^{n+1}, x') dx' \Delta t,$$

where

$$\begin{aligned}\Theta(t^{n+1}, x') &= \int_0^{t^N} \int_{\mathbb{R}^d} F(u(t^n, x'), v(t, x)) \cdot \nabla_x (\bar{\varphi}(t, x, t^{n+1}, x') - \varphi(t, x, t^{n+1}, x')) \, dx \, dt \\ &\quad + \frac{\Delta t}{2} \int_0^{t^N} \int_{\mathbb{R}^d} F_t(u(t^n, x'), v(t, x)) \cdot \nabla_x \varphi(t, x, t^{n+1}, x') \, dx \, dt.\end{aligned}$$

Noticing that $\bar{\varphi}(t, t^{n+1}) - \varphi(t, t^{n+1}) = \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} (s - t^n) \varphi_t(t, s) \, ds$, one gets after integrating by parts and some elementary manipulations,

$$\begin{aligned}\Theta(t^{n+1}, x') &= \int_0^{t^N} \int_{\mathbb{R}^d} F_t(u(t^n, x'), v(t, x)) \cdot \nabla_x \int_{t^n}^{t^{n+1}} \frac{(s - t^n)^2}{2} \varphi_{t'}(t, x, s, x') \, ds \, dx \, dt \\ &\quad - \int_{\mathbb{R}^d} \nabla_x \cdot F(u(t^n, x'), v(t^N, x)) \int_{t^n}^{t^{n+1}} (s - t^n) \varphi(t^N, x, s, x') \, ds \, dx \\ &\quad + \int_{\mathbb{R}^d} \nabla_x \cdot F(u(t^n, x'), v_0(x)) \int_{t^n}^{t^{n+1}} (s - t^n) \varphi(0, x, s, x') \, ds \, dx.\end{aligned}$$

This allows us to write $\xrightarrow{time} TE_{h.o.t.}(u, v; t^N) = I - II + III$, with the obvious notation. Now we only need to estimate terms I, II , and III . First, we have

$$|I| \leq T_{aux} \|f'(v)\|^2 |v_0|_{TV(\mathbb{R}^d)} t^N,$$

where

$$\begin{aligned}T_{aux} &= \frac{(\Delta t)^2}{2} \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \int_0^{t^N} \int_{\mathbb{R}^d} |\nabla_x \varphi_{t'}(t, x, t', x')| \, dx' \, dt' \\ &\leq \frac{(\Delta t)^2}{2} \frac{d \|\eta\|_{TV(\mathbb{R})}}{\epsilon_t \epsilon_x} W(t^N),\end{aligned}$$

where we follow the lines of [6, Lemma 7.12] and [6, Lemma 7.13].

Next we estimate term II :

$$|II| \leq T_{aux} \|f'(v)\| |v_0|_{TV(\mathbb{R}^d)} \Delta t,$$

where

$$\begin{aligned}T_{aux} &= \sup_{x \in \mathbb{R}^d} \int_0^{t^N} \int_{\mathbb{R}^d} |\phi(t^N, x, t', x')| \, dx' \, dt' \\ &\leq 2 \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N),\end{aligned}$$

by [7, Lemma 3.7]. The term III can be estimated in exactly the same way. Combining the estimates of terms I, II, III completes the proof. $\square$

Before we estimate $\xrightarrow{cent} TE_{h.o.t.}(u, v; t^N)$ and $\xrightarrow{visc} TE_{h.o.t.}(u, v; t^N)$, we need to prove a couple of simple auxiliary results.

LEMMA 3.10. *We have*

$$\sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| d_e \mathcal{I}_{e,K}(\Phi) \leq 2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \|\Phi\|_{L^1(\mathbb{R}^d)},$$

where

$$\begin{aligned} \mathcal{I}_{e,K}(\Phi) &= \frac{1}{|K_e|} \int_{K_e} \frac{1}{|K|^3} \int_{K^3} \int_0^1 \int_0^1 \int_0^1 \Phi \, d\lambda \, d\mu \, d\nu \, dz' \, dy' \, dw' \, dx', \\ \Phi &\equiv \Phi(\lambda(\mu(\nu x' + (1-\nu)w') + (1-\mu)y') + (1-\lambda)z'). \end{aligned}$$

Proof. Since the point $\lambda(\mu(\nu x' + (1-\nu)w') + (1-\mu)y') + (1-\lambda)z'$ belongs to the convex hull of K and K_e , C_e , we have

$$|\mathcal{I}_{e,K}(\Phi)| \leq \frac{1}{\max\{|K|, |K_e|\}} \|\Phi\|_{L^1(C_e)}.$$

Hence,

$$\begin{aligned} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| d_e \mathcal{I}_{e,K}(\Phi) &\leq \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} \frac{|e| d_e}{\max\{|K|, |K_e|\}} \|\Phi\|_{L^1(C_e)} \\ &\leq \mathbb{D}_{\Delta x} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} \|\Phi\|_{L^1(C_e)}, \quad \text{by (2.5),} \\ &\leq 2 \mathbb{D}_{\Delta x} \sum_{e \in \mathcal{E}_{\Delta x}} \|\Phi\|_{L^1(C_e)} \\ &\leq 2 \mathbb{D}_{\Delta x} \sum_{K \in \mathcal{T}_{\Delta x}} \iota_{\Delta x} \|\Phi\|_{L^1(K)} \\ &\leq 2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \|\Phi\|_{L^1(\mathbb{R}^d)}, \quad \text{by (2.6).} \end{aligned}$$

This completes the proof. $\square$

LEMMA 3.11. *We have*

$$\Theta_{aux} \leq 4 t^N (\Delta x)^2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N) \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\},$$

where

$$\Theta_{aux} = \sum_{1 \leq j, k \leq d} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| b_{e,K_e} d_e \mathcal{I}_{e,K}(|\partial_{x_j x_k} \varphi(t, x)|) \Delta t.$$

Proof. By the definition of the operator $\mathcal{I}_{e,K}$ in Lemma 3.10 and the definition of φ , (3.2), we can write $\Theta_{aux} = \Theta_{aux,t} \cdot \Theta_{aux,x}$, where

$$\Theta_{aux,t} = \sum_{n=0}^{N-1} \omega_{\epsilon_t}(t - t^{n+1}) \Delta t \leq 2 \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N),$$

by [7, Lemma 3.7] and

$$\Theta_{aux,x} = \sum_{1 \leq j, k \leq d} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| b_{e,K_e} d_e \mathcal{I}_{e,K}(|\partial_{x_j x_k} \mathbb{R}^{jk}(x, \cdot)|),$$

where $\mathbb{R}^{jk}(x, x') = \prod_{i=1}^d \eta_{\epsilon_x}(x_i - x'_i)$. Since, by Lemma 3.10,

$$\Theta_{aux,x} \leq 2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \sum_{1 \leq j, k \leq d} \|\mathbb{R}^{jk}(x, \cdot)\|_{L^1(\mathbb{R}^d)},$$

and since

$$\sum_{1 \leq j, k \leq d} \left| \partial_{x_j x_k} \prod_{i=1}^d \eta_{\epsilon_x}(x_i - x'_i) \right| = \sum_{1 \leq j, k \leq d} \left| \prod_{\substack{i=1 \\ i \neq j \neq k}}^d \eta_{\epsilon_x}(x_i - x'_i) \eta'_{\epsilon_x}(x_k - x'_k) \eta'_{\epsilon_x}(x_j - x'_j) \right. \\ \left. + \prod_{\substack{i=1 \\ i \neq j}}^d \eta_{\epsilon_x}(x_i - x'_i) \eta''_{\epsilon_x}(x_j - x'_j) \delta_{jk} \right|,$$

we have that

$$\Theta_{aux,x} \leq 2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\}.$$

This completes the proof. $\square$

We are now ready to estimate the remaining term in $TE_{h.o.t.}$.

LEMMA 3.12. *Suppose that the conditions (2.3) are satisfied. Then,*

$$\overset{cent}{\rightarrow} TE_{h.o.t.}(u, v; t^N) \leq C_2 (\Delta x)^2 \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \left(1 + \frac{\Delta t}{\epsilon_t} \right), \\ \overset{visc}{\rightarrow} TE_{h.o.t.}(u, v; t^N) \leq C_2 (\Delta x)^2 \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \left(1 + \frac{\Delta t}{\epsilon_t} \right) \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})},$$

where $C_2 = 16 t^N |v_0|_{TV(\mathbb{R}^d)} \|f'(v)\| \mathbb{D}_{\Delta x} \iota_{\Delta x} W(t^N)$.

Proof. We only have to prove the first estimate since the second is similar. We have

$$\overset{cent}{\rightarrow} TE_{h.o.t.}(u, v; t^N) = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| b_{e,K_e} \Theta_{e,K}^n \Delta t,$$

where

$$\Theta_{e,K}^n = \sum_{1 \leq i, j, k \leq d} \int_0^{t^N} \int_{\mathbb{R}^d} F(u_K^n, v(t, x)) \cdot n_{e,K} \partial_{x_i} \mathcal{I}_{e,K}(\Xi^{ijk}(t, x)) dx dt.$$

Moreover, $\Xi^{ijk}(t, x) = \zeta^{ijk} \partial_{x_j x_k} \phi$, where ζ^{ijk} and ϕ are as in Lemma 3.4.

Performing integrations by parts with respect to x_i , we get

$$\Theta_{e,K}^n = - \sum_{1 \leq i, j, k \leq d} \int_0^{t^N} \int_{\mathbb{R}^d} (\partial_v F(u_K^n, v(t, x)) \cdot n_{e,K}) \partial_{x_i} v(t, x) \mathcal{I}_{e,K}(\Xi^{ijk}(t, x)) dx dt,$$

and hence,

$$\overset{cent}{\rightarrow} TE_{h.o.t.}(u, v; t^N) \leq T_{aux} \|f'(v)\| \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{1 \leq i \leq d} |\partial_{x_i} v(t, x)| dx dt \\ \leq T_{aux} \|f'(v)\| t^N |v_0|_{TV(\mathbb{R}^d)},$$

where

$$T_{aux} = \max_{1 \leq i \leq d} \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \sum_{1 \leq j, k \leq d} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} |e| b_{e, K_e} \mathcal{I}_{e, K}(\Xi^{ijk}(t, x)) \Delta t.$$

Since

$$|\zeta^{ijk}| \leq 4(\Delta x)^2 d_e,$$

we have that

$$|\mathcal{I}_{e, K}(\Xi^{ijk}(t, x))| \leq 4(\Delta x)^2 d_e \mathcal{I}_{e, K}(|\partial_{x_j} x_k \varphi(t, x)|),$$

and so,

$$\begin{aligned} T_{aux} &\leq 4(\Delta x)^2 \Theta_{aux} \\ &\leq 16 t^N (\Delta x)^2 \mathbb{D}_{\Delta x} \iota_{\Delta x} \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N) \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \end{aligned}$$

by Lemma 3.11. This completes the proof. $\square$

3d. Proof of the error estimate. To obtain the error estimate, we proceed exactly as in [6]. Inserting the estimates obtained in sections 3.b and 3.c into the approximation inequality in section 3.a, and taking the auxiliary function η as in (3.3), we obtain

$$\begin{aligned} e(t^N) &\leq 2e(0) + 8(\epsilon_x + \epsilon_t \|f'(v)\|) |v_0|_{TV(\mathbb{R}^d)} + 2\|f'(v)\| |v_0|_{TV(\mathbb{R}^d)} \Delta t \\ &\quad + 2C_0 t^N \left\{ 2 \frac{d |\eta|_{TV(\mathbb{R})}}{\epsilon_x} \left(1 + \frac{\Delta t}{\epsilon_t}\right) \right\} \|\nu_v\| \Delta x \\ &\quad + 4C_1 t^N \left(1 + \frac{\Delta t}{\epsilon_t}\right) \left(1 + \frac{(\Delta x)^{d/2}}{\epsilon_x^d}\right) \|\eta\|_{L^\infty(\mathbb{R})}^d (|\delta|_{var, 1/2} + |\alpha|_{var, 1/2}) \\ &\quad + 2C_1 \left\{ \frac{1}{2} \|f'(v)\| \frac{(\Delta t)^2 d |\eta|_{TV(\mathbb{R})}}{\epsilon_t \epsilon_x} t^N + 4 \left(1 + \frac{\Delta t}{\epsilon_t}\right) \Delta t \right. \\ &\quad \left. + 32 \mathbb{D}_{\Delta x} \iota_{\Delta x} (\Delta x)^2 \left\{ \frac{d^2 |\eta|_{TV(\mathbb{R})}^2}{\epsilon_x^2} + \frac{d |\eta'|_{TV(\mathbb{R})}}{\epsilon_x^2} \right\} \left(1 + \frac{\Delta t}{\epsilon_t}\right) t^N \right\}, \end{aligned}$$

where $C_0 = |v_0|_{TV(\mathbb{R}^d)}$ and $C_1 = C_0 \|f'(v)\|$. The estimate of Theorem 2.1 is then obtained by eliminating the parameter Δt through the use of the CFL condition (2.4) and then taking the very same optimal values as those in [6], namely,

$$\epsilon_x^* = \sqrt{t^N \|\nu_v\| \Delta x / 2}, \quad \epsilon_t = A_t (\Delta x)^{3/4}, \quad \epsilon = A (\Delta x)^{1/4}.$$

This concludes the proof of Theorem 2.1.

4. Proof of Theorem 2.3. In this section, we prove Theorem 2.3. The only difference between the proof of Theorem 2.3 and that of Theorem 2.1 is the following estimate of the consistency term of the truncation error, $TE_{cons}(u, v; t^N)$.

LEMMA 4.1. *We have*

$$TE_{cons}(u, v; t^N) \leq C_1 \left\{ \mathbb{D}_{\Delta x} \iota_{\Delta x} \|f'(v)\| t^N |v_0|_{TV(\mathbb{R}^d)} \right. \\ \left. + \|f'(u)\| \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} |e| |u_{K_e^+}^n - u_{K_e^-}^n| \Delta t \right\},$$

where $C_1 = 2 \frac{d|\eta|_{TV}}{\epsilon_x} (1 + \frac{\Delta t}{\epsilon_t}) W(t^N) \{ \|\delta\|_{\ell^\infty(\mathcal{E}_{\Delta x})/\mathbb{R}^d} + \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})} \}$ and $\mathbb{D}_{\Delta x}$ and $\iota_{\Delta x}$ are defined in (2.5) and (2.6), respectively.

Theorem 2.3 follows easily from this result, section 3, as well as the following observation:

$$\sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} |e| |u_{K_e^+}^n - u_{K_e^-}^n| \Delta t \leq \frac{1}{\iota_{inf}} \|u\|_{L^1(0, t^N; TV(\mathbb{R}^d))},$$

where $\iota_{inf} = \inf_{K \in \mathcal{T}_{\Delta x}} \iota_K$. We note again that beyond merely proving the above estimate, we aim at understanding the mechanism that allows it to be possible; see the fourth step of the proof that follows. This provides an explanation of the *supra-convergence* phenomenon for the schemes under consideration.

To prove Lemma 4.1, we will need the following simple auxiliary result.

LEMMA 4.2. *We have*

$$\sum_{e \in \mathcal{E}_{\Delta x}} |e| |\psi_{K_e^+} - \psi_{K_e^-}| \leq \mathbb{D}_{\Delta x} \iota_{\Delta x} |\psi|_{TV(\mathbb{R}^d)},$$

where $\psi_K = \frac{1}{|K|} \int_K \psi(x') dx'$.

Proof. Let us call Θ the left-hand side of the inequality we want to prove. Then by the proof of Lemma 3.4,

$$\Theta = \sum_{e \in \mathcal{E}_{\Delta x}} |e| \left| \frac{1}{|K_e^+| |K_e^-|} \int_{K_e^+} \int_{K_e^-} \int_0^1 (x' - w') \cdot \nabla_{x'} \psi(\nu x' + (1 - \nu)w') d\nu dw' dx' \right|.$$

Assume that $|K_e^-| = \max\{|K_e^+|, |K_e^-|\}$; then, since $|x' - w'| \leq d_e$ and the point $\nu x' + (1 - \nu)w'$ belongs to the convex hull of K and K_e , C_e , we have

$$\begin{aligned} \Theta &\leq \sum_{e \in \mathcal{E}_{\Delta x}} |e| \left| \frac{d_e}{|K_e^+| |K_e^-|} \int_{K_e^+} \int_0^1 \int_{C_e} \sum_{i=1}^d |\partial_{x_i} \psi(w')| dw' d\nu dx' \right| \\ &\leq \sum_{e \in \mathcal{E}_{\Delta x}} \frac{|e| d_e}{|K_e^-|} |\psi|_{TV(C_e)} \\ &\leq \mathbb{D}_{\Delta x} \sum_{e \in \mathcal{E}_{\Delta x}} |\psi|_{TV(C_e)}, \quad \text{by (2.5),} \\ &\leq \mathbb{D}_{\Delta x} \sum_{K \in \mathcal{T}_{\Delta x}} \iota_K |\psi|_{TV(K)} \\ &\leq \mathbb{D}_{\Delta x} \iota_{\Delta x} \sum_{K \in \mathcal{T}_{\Delta x}} |\psi|_{TV(K)}, \quad \text{by (2.6),} \\ &\leq \mathbb{D}_{\Delta x} \iota_{\Delta x} |\psi|_{TV(\mathbb{R}^d)}. \end{aligned}$$

This completes the proof. $\square$

We are now ready to prove Lemma 4.1.

Proof of Lemma 4.1. We proceed in several steps.

First step. First, we rewrite the term $TE_{cons}(u, v; t^N)$ as follows:

$$TE_{cons}(u, v; t^N) = \sum_{n=0}^{N-1} \int_0^{t^N} \int_{\mathbb{R}^d} \Theta(t, x, t^{n+1}; v(t, x)) dx dt \Delta t,$$

where

$$\Theta(t, x, t^{n+1}; c) = \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} \nabla_x \phi(t, x, t^{n+1}, x_K) \cdot \mathcal{A}_{e,K}(u_K^n, c),$$

and

$$\mathcal{A}_{e,K}(u, c) = |e| (F(u, c) \cdot n_{e,K} \delta_e - \alpha_e \mathcal{N}_e(u, c) (x_{K_e} - x_K)).$$

Note that, since $\sum_{e \in \partial K} |e| n_{e,K} = 0$, we can modify $\mathcal{A}_{e,K}(u, c)$ as follows:

$$\mathcal{A}_{e,K}(u, c) = |e| (F(u, c) \cdot n_{e,K} (\delta_e - \hat{\delta}) - \alpha_e \mathcal{N}_e(u, c) (x_{K_e} - x_K)),$$

where $\hat{\delta}$ is an arbitrary constant vector.

Second step. Next, we rewrite $\Theta(t, x, t^{n+1}; c)$ as a sum on all the faces e of finite volumes K of the triangulation, $\mathcal{E}_{\Delta x}$. To perform this discrete integration by parts, we denote by K_e^+ and K_e^- the two finite volumes sharing the face e , drop the superscript n in u_K^n , and write ϕ_K instead of $\phi(t, x, t^{n+1}, x_K)$; we get

$$\begin{aligned} \Theta(t, x, t^{n+1}; c) &= \sum_{K \in \mathcal{T}_{\Delta x}} \sum_{e \in \partial K} \nabla_x \phi_K \cdot \mathcal{A}_{e,K}(u_K, c) \\ &= \sum_{e \in \mathcal{E}_{\Delta x}} \{ \nabla_x \phi_{K_e^+} \cdot \mathcal{A}_{e,K_e^+}(u_{K_e^+}, c) + \nabla_x \phi_{K_e^-} \cdot \mathcal{A}_{e,K_e^-}(u_{K_e^-}, c) \} \\ &= \Theta_1(t, x, t^{n+1}; c) + \Theta_2(t, x, t^{n+1}; c), \end{aligned}$$

where

$$\begin{aligned} \Theta_1(t, x, t^{n+1}; c) &= \sum_{e \in \mathcal{E}_{\Delta x}} \nabla_x (\phi_{K_e^+} - \phi_{K_e^-}) \cdot \mathcal{A}_{e,K_e^+}(u_{K_e^+}, c), \\ \Theta_2(t, x, t^{n+1}; c) &= \sum_{e \in \mathcal{E}_{\Delta x}} \nabla_x \phi_{K_e^-} \cdot (\mathcal{A}_{e,K_e^+}(u_{K_e^+}, c) + \mathcal{A}_{e,K_e^-}(u_{K_e^-}, c)) \\ &\equiv \sum_{e \in \mathcal{E}_{\Delta x}} \nabla_x \phi_{K_e^-} \cdot \mathcal{B}_e(u_{K_e^-}, u_{K_e^+}, c), \end{aligned}$$

with the obvious notation for $\mathcal{B}_e$. The above results allow us to write $TE_{cons}(u, v; t^N) = TE_{1,cons} + TE_{2,cons}$.

Third step. To estimate $TE_{1,cons}$, we proceed as follows. After a simple integration by parts, we obtain

$$TE_{1,cons} = - \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} (\phi_{K_e^+} - \phi_{K_e^-}) \nabla_x v(t, x) \cdot \partial_v \mathcal{A}_{e,K_e^+}(u_{K_e^+}, v(t, x)) dx dt \Delta t,$$

and after taking absolute values,

$$TE_{1,cons} \leq \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} |\phi_{K_e^+} - \phi_{K_e^-}| \sum_{i=1}^d |\partial_{x_i} v(t, x)| \mathbb{A}_e(v(t, x)) dx dt \Delta t,$$

where

$$\mathbb{A}_e(c) = |e| \max_{1 \leq i \leq d} |f'(c) \cdot n_{e, K_e^+} (\delta_e - \hat{\delta})_i - \alpha_e N'_e(c) (x_{K_e^-} - x_{K_e^+})_i|.$$

Hence,

$$TE_{1,cons} \leq T_{aux} \mathbb{C} \|f'(v)\| t^N |v_0|_{TV(\mathbb{R}^d)},$$

where

$$\mathbb{C} = \|\delta_e - \hat{\delta}\|_{\ell^\infty(\mathcal{E}_{\Delta x})} + \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})},$$

and

$$\begin{aligned} T_{aux} &= \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} |e| |\phi_{K_e^+} - \phi_{K_e^-}| \Delta t \\ &\leq 2 \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N) \mathbb{D}_{\Delta x} \iota_{\Delta x} \frac{d|\eta|_{TV(\mathbb{R})}}{\epsilon_x}, \end{aligned}$$

by Lemma 3.7 of [7] and Lemma 4.2. This implies

$$TE_{1,cons} \leq 2 \frac{d|\eta|_{TV(\mathbb{R})}}{\epsilon_x} \left(1 + \frac{\Delta t}{\epsilon_t}\right) W(t^N) \mathbb{C} \mathbb{D}_{\Delta x} \iota_{\Delta x} \|f'(v)\| t^N |v_0|_{TV(\mathbb{R}^d)}.$$

Note that the term $TE_{1,cons}$ has been estimated without using regularity properties of the approximate solution. This is not the case for the term $TE_{2,cons}$, as we see next.

Fourth step. To estimate $TE_{2,cons}$, we simply take absolute values to obtain

$$TE_{2,cons} \leq \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} \int_0^{t^N} \int_{\mathbb{R}^d} \sum_{i=1}^d |\partial_{x_i} \phi_{K_e^-}| \mathbb{B}_e(u_{K_e^-}, u_{K_e^+}) dx dt \Delta t,$$

where

$$\begin{aligned} \mathbb{B}_e(u_{K_e^-}, u_{K_e^+}) &= \sup_{\substack{t \in (0, t^N) \\ x \in \mathbb{R}^d}} |\mathcal{B}_e(u_{K_e^-}, u_{K_e^+}, v(t, x))| \\ &\leq |e| \left\{ |f(u_{K_e^+}) - f(u_{K_e^-}) \cdot n_{e,k}| \|\delta_e - \hat{\delta}\|_{\ell^\infty(\mathcal{E}_{\Delta x})} \right. \\ &\quad \left. + |N_e(u_{K_e^+}) - N_e(u_{K_e^-}) \cdot n_{e,k}| \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})} \right\} \\ &\leq |e| \|f'(u)\| \mathbb{C} |u_{K_e^+} - u_{K_e^-}|. \end{aligned}$$

This is the crucial estimate that allows the *supraconvergence* phenomenon to take place: The estimate of $\mathbb{B}_e(u_{K_e^-}, u_{K_e^+})$ depends *only* on ℓ^∞ -norms of δ_e and $\alpha_e(x_{K_e^+} - x_{K_e^-})$. This is possible because the above quantities depend *only* on the face e and

this fact is a direct consequence of the conservativity and consistency properties of the fluxes (see section 2.c).

However, the price to pay, is that now the estimate of $TE_{2,cons}$ must depend on the smoothness properties of the approximate solution since

$$TE_{2,cons} \leq \frac{d|\eta|_{TV(\mathbb{R})}}{\epsilon_x} W(t^N) \mathbb{C} \|f'(u)\| \sum_{n=0}^{N-1} \sum_{e \in \mathcal{E}_{\Delta x}} |e| |u_{K_e^+}^n - u_{K_e^-}^n| \Delta t.$$

Fifth step. The result now follows from the estimates of $TE_{1,cons}$ and $TE_{2,cons}$, from the fact that $TE_{cons}(u, v; t^N) = TE_{2,cons} + TE_{1,cons}$, and from the fact that, since the vector $\hat{\delta}$ is arbitrary, we can take $\mathbb{C} = \|\delta\|_{\ell^\infty(\mathcal{E}_{\Delta x})/\mathbb{R}^d} + \|\alpha\|_{\ell^\infty(\mathcal{E}_{\Delta x})}$. $\square$

5. Concluding remarks. In [6], a general theory of a priori estimates for scalar conservation laws was proposed. The approach is based on a modification of the original Kuznetsov approximation theory [17]. The key feature and advantage of this new and modified method is that *no* regularity properties of the approximate solution are needed. This point is significant since establishing such properties is a well-known bottleneck in this area. This approach was applied to general multidimensional flux-splitting monotone schemes in [7]. For grids consisting of Cartesian products of possibly nonuniform one-dimensional meshes, optimal order of convergence was obtained.

In this paper, we generalize this result to general grids. It should be emphasized that, to the authors' knowledge, this is the first derivation of optimal orders of convergence for truly multidimensional numerical methods for conservation laws. By truly multidimensional, we mean that, in the spirit of the finite volume methods considered, no “dimension-by-dimension” argument is used. This fact is not a mere detail since, as shown in the introduction, standard properties like total variation diminishing (TVD) do not hold, in general, on non-Cartesian meshes, which accounts for the lack of similar results so far.

The relationship between the consistency of the method and the properties of the grid is analyzed. In particular, it is shown that first, requiring consistency imposes extremely strong restrictions on both the mesh and the numerical fluxes, second, those restrictions are not needed, neither theoretically nor practically. A much larger class of “nearly consistent” (but *not* consistent) schemes is studied. In both cases, optimal a priori estimates are obtained. It is shown how the combination of the *conservation form* of the scheme together with the *consistency of the numerical fluxes* allows this *supraconvergence* phenomenon to take place. For schemes that are nonconsistent all together, optimal error estimates are obtained as well. However, the nature of the term describing the loss of consistency does require a bound on the total variation $L^1(0, T; TV(\mathbb{R}^d))$ of the approximate solution (which is used there and only there).

In conclusion, we have shown that the approach taken in [6, 7] keeps its promises, by being able to tackle problems previously unsolved. The application to nonsplitting numerical fluxes, and to higher-order methods will be the object of future publications. The generalization to *systems* of conservation laws constitutes an exciting challenge.

Acknowledgment. The third author would like to thank Timothy Brule for providing a code to generate Figure 1.

REFERENCES

- [1] T. BOUKADIDA AND A.Y. LEROUX, *A new version of the two-dimensional Lax–Friedrichs scheme*, Math. Comp., 63 (1994), pp. 541–553.
- [2] F. COQUEL AND P. LEFLOCH, *Convergence of finite difference schemes for conservation laws in several space variables: A general theory*, SIAM J. Numer. Anal., 30 (1993), pp. 675–700.
- [3] B. COCKBURN, F. COQUEL, AND P. LEFLOCH, *Convergence of the finite volume method for multidimensional conservation laws*, SIAM J. Numer. Anal., 32 (1995), pp. 687–705.
- [4] B. COCKBURN, F. COQUEL, AND P. LEFLOCH, *An error estimate for finite volume methods for conservation laws*, Math. Comp., 64 (1994), pp. 77–103.
- [5] B. COCKBURN AND P.-A. GREMAUD, *An error estimate for finite element methods for conservation laws*, SIAM J. Numer. Anal., 33 (1996), pp. 522–554.
- [6] B. COCKBURN AND P.-A. GREMAUD, *A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach*, Math. Comp., 65 (1996), pp. 533–573.
- [7] B. COCKBURN AND P.-A. GREMAUD, *A priori error estimates for numerical methods for scalar conservation laws. Part II: Flux-splitting monotone schemes on irregular Cartesian grids*, Math. Comp., 66 (1997), pp. 547–572.
- [8] M. CRANDALL AND A. MAJDA, *Monotone difference approximations for scalar conservation laws*, Math. Comp., 34 (1980), pp. 1–21.
- [9] B. ENGQUIST AND S. OSHER, *One sided difference approximations for nonlinear conservation laws*, Math. Comp., 36 (1981), pp. 321–351.
- [10] B. GARCÍA-ARCHILLA, *A supraconvergent scheme for the Korteweg-de-Vries equation*, Numer. Math., 61 (1992), pp. 291–310.
- [11] B. GARCÍA-ARCHILLA AND J.M. SANZ-SERNA, *A finite difference formula for the discretization of d^3/dx^3 on nonuniform grids*, Math. Comp., 57 (1991), pp. 239–257.
- [12] M. GEIBEN, D. KRÖNER, AND M. ROKYTA, *A Lax–Wendroff type theorem for upwind finite volume schemes in 2-D*, East-West J. Numer. Math., 4 (1996), pp. 279–292.
- [13] J.D. HOFFMAN, *Relationship between the truncation error of centered finite difference approximations on uniform and nonuniform meshes*, J. Comput. Phys., 46 (1982), pp. 469–474.
- [14] H.-O. KREISS, T.A. MANTEUFFEL, B. SWARTZ, B. WENDROFF, AND A.B. WHITE, JR., *Supraconvergent schemes on irregular grids*, Math. Comp., 47 (1986), pp. 537–554.
- [15] D. KRÖNER AND M. ROKYTA, *Convergence of upwind finite volume schemes for scalar conservation laws in 2-D*, SIAM J. Numer. Anal., 31 (1994), pp. 324–343.
- [16] S.N. KRUZHKOV, *First order quasilinear equations in several independent variables*, Math. USSR-Sb., 10 (1970), pp. 217–243.
- [17] N.N. KUZNETSOV, *The accuracy of some approximate methods for computing the weak solutions of a first-order quasilinear equation*, USSR Comp. Math. and Math. Phys., 16 (1976), pp. 105–119.
- [18] B. J. LUCIER, *A stable adaptive scheme for hyperbolic conservation laws*, SIAM J. Numer. Anal., 22 (1985), pp. 180–203.
- [19] B. J. LUCIER, *Error bounds for the methods of Glimm, Godunov, and LeVeque*, SIAM J. Numer. Anal., 22 (1985), pp. 1074–1081.
- [20] B. J. LUCIER, *On nonlocal monotone difference schemes for scalar conservation laws*, Math. Comp., 47 (1986), pp. 19–36.
- [21] B. J. LUCIER, *A moving mesh numerical method for hyperbolic conservation laws*, Math. Comp., 46 (1986), pp. 59–69.
- [22] T.A. MANTEUFFEL AND A.B. WHITE, JR., *The numerical solution of second-order boundary value problems on nonuniform meshes*, Math. Comp., 47 (1986), pp. 511–535.
- [23] H. NESSYAHU AND E. TADMOR, *The convergence rate of approximate solutions for nonlinear scalar conservation laws*, SIAM J. Numer. Anal., 29 (1992), pp. 1505–1519.
- [24] H. NESSYAHU, E. TADMOR, AND T. TASSA, *The convergence rate of Godunov type schemes*, SIAM J. Numer. Anal., 31 (1994), pp. 1–16.
- [25] S. NÖELLE, *A note on entropy inequalities and error estimates for higher-order accurate finite volume schemes on irregular grids*, Math. Comp., 65 (1996), pp. 1155–1163.
- [26] J. PIKE, *Grid adaptive algorithms for the solution of the Euler equations on irregular grids*, J. Comput. Phys., 71 (1987), pp. 194–223.
- [27] R. SANDERS, *On convergence of monotone finite difference schemes with variable spatial differencing*, Math. Comp., 40 (1983), pp. 91–106.
- [28] G. STRANG, *Accurate partial difference methods II: Non-linear problems*, Numer. Math., 6 (1964), pp. 37–46.
- [29] E. TADMOR, *Local error estimates for discontinuous solutions of nonlinear hyperbolic equations*, SIAM J. Numer. Anal., 28 (1991), pp. 891–906.

- [30] E. TURKEL, *Accuracy of schemes with nonuniform meshes for compressible fluid flow*, Appl. Numer. Math., 2 (1986), pp. 529–550.
- [31] J.-P. VILA, *Convergence and error estimates in finite volume schemes for general multidimensional scalar conservation laws*, Model. Math. Anal. Numer., 28 (1994), pp. 267–295.
- [32] B. WENDROFF, *Supraconvergence in Two Dimensions*, Los Alamos National Laboratory Report LA-UR-95-3068, Vieweg, 1995.
- [33] B. WENDROFF AND A.B. WHITE, JR., *Some supraconvergent schemes for hyperbolic equations on irregular grids*, in Nonlinear Hyperbolic Equations—Theory, Numerical Methods and Applications, J. Ballman and R. Jeltsch, eds., Proc. Second International Conference on Hyperbolic Equations, Aachen, 1988, pp. 671–677.
- [34] B. WENDROFF AND A.B. WHITE, JR., *A supraconvergent scheme for nonlinear hyperbolic systems*, Comput. Math. Appl., 18 (1989), pp. 761–767.
- [35] J.X. YANG, *Analysis of numerical methods for nonlinear hyperbolic problems*, PhD. thesis, University of Minnesota, Minneapolis, Minnesota, in preparation.