- CULSHAW, R. J., G. J. BANCROFT, AND V. McDONALD. 1997. Gut intraepithelial lymphocytes induce immunity against *Cryptosporidium* infection through a mechanism involving gamma interferon production. *Infection and Immunity* **65**: 3074–3079.
- DAVAMI, M. H., G. J. BANCROFT, AND V. McDONALD. 1997. *Cryptosporidium* infection in major histocompatibility complex congenic strains of mice: Variation in susceptibility and the role of T-cell cytokine response. *Parasitology Research* **83**: 257–263.
- GRAHAM, M. B., D. K. DALTON, D. GILTINAN, V. L. BRACIALE, T. A. STEART, AND T. J. BRACIALE. 1993. Response to influenza infection in mice with a targeted disruption in the interferon γ gene. *Journal of Experimental Medicine* **178**: 1725–1732.
- HARP, J. A., AND R. E. SACCO. 1996. Development of cellular immune functions in neonatal to weanling mice: Relationship to *Cryptosporidium parvum* infection. *Journal of Parasitology* **82**: 245–249.
- , M. W. WANNEMUEHLER, D. B. WOODMANSEE, AND H. W. MOON. 1988. Susceptibility of germfree and antibiotic-treated adult mice. *Infection and Immunity* **56**: 2006–2010.
- , W. M. WHITMIRE, AND R. SACCO. 1994. In vitro proliferation and production of gamma interferon by murine CD4 cells in response to *Cryptosporidium parvum* antigen. *Journal of Parasitology* **80**: 67–92.
- HEINE, J., H. W. MOON, AND D. B. WOODMANSEE. 1984. Persistent *Cryptosporidium* infection in congenitally athymic (nude) mice. *Infection and Immunity* **43**: 856–859.
- McDONALD, V., AND G. J. BANCROFT. 1994. Mechanisms of innate and acquired resistance to *Cryptosporidium parvum* infection in SCID mice. *Infection and Immunity* **62**: 1474–1479.
- NOVAK, S. M., AND C. R. STERLING. 1991. Susceptibility dynamics in neonatal BALB/c mice infected with *Cryptosporidium parvum*. *Journal of Protozoology* **38**: 102S.
- O'DONOGHUE, P. J. 1995. *Cryptosporidium* and cryptosporidiosis in man and animals. *International Journal of Parasitology* **25**: 139–195.
- RASMUSSEN, K. R., AND M. C. HEALEY. 1992. Experimental *Cryptosporidium parvum* infections in immunosuppressed adult mice. *Infection and Immunity* **60**: 1648–1652.
- REHG, J. E. 1996. Effect of interferon- γ in experimental *Cryptosporidium parvum* infection. *Journal of Infectious Diseases* **174**: 232–237.
- THEODOS, C., K. L. SULLIVAN, J. K. GRIFFITHS, AND S. TZIPIRI. 1995. Profiles of healing and nonhealing *Cryptosporidium parvum* infection in C57BL/6 mice with functional B and T lymphocytes: The extent of gamma interferon modulation determines the outcome of infection. *Infection and Immunity* **65**: 4761–4769.
- TILLEY, M., V. McDONALD, AND G. J. BANCROFT. 1995. Resolution of cryptosporidial infection in mice correlates with parasite-specific lymphocyte proliferation associated with Th1 and Th2 cytokine secretion. *Parasite Immunology* **17**: 459–464.
- TZIPIRI, S., W. RAND, AND C. THEODOS. 1995. Evaluation of a two-phase scid mouse model preconditioned with anti-interferon monoclonal antibody for drug testing against *Cryptosporidium parvum*. *Journal of Infectious Diseases* **172**: 1160–1164.
- UNGAR, B. L. P., T. C. KAO, J. A. BURRIS, AND F. D. FINKELMAN. 1994. *Cryptosporidium* infection in an adult mouse model. Independent roles for IFN- γ and CD4 T lymphocytes in protective immunity. *Journal of Immunology* **153**: 1022–1024.
- URBAN, J. F., JR., R. FAYER, S.-J. CHEN, W. C. GAUSE, M. K. GATLIN, AND F. D. FINKELMAN. 1996. IL-12 protects immunodeficient neonatal mice against *Cryptosporidium parvum*. *Journal of Immunology* **156**: 236–241.
- WANG, Z. E., S. L. REINER, S. ZHENG, D. K. DALTON, AND R. M. LACKSLEY. 1994. CD4 effector cells default to the Th2 pathway in interferon gamma-deficient mice infected with *Leishmania major*. *Journal of Experimental Medicine* **179**: 1367–1371.
- WYATT, C. R., E. J. BRACKETT, L. E. PERRYMAN, A. C. RICE-FORCE, C. BROWN, AND K. I. O'ROURKE. 1997. Activation of intestinal intraepithelial T lymphocytes in calves infected with *Cryptosporidium parvum*. *Infection and Immunity* **65**: 185–190.

Seroprevalence of *Toxoplasma gondii* and *Trichinella spiralis* in North Carolina Black Bears (*Ursus americanus*)

F. B. Nutter[†]*, J. F. Levine^{††}, M. K. Stoskopf[‡], H. R. Gamble[§], and J. P. Dubey[§], *Department of Companion Animal and Special Species Medicine, College of Veterinary Medicine, North Carolina State University, 4700 Hillsborough St., Raleigh, North Carolina 27606; [†]Department of Microbiology, Pathology, and Parasitology, College of Veterinary Medicine, North Carolina State University, 4700 Hillsborough St., Raleigh, North Carolina 27606; ^{††}Environmental Medicine Consortium, College of Veterinary Medicine, North Carolina State University, 4700 Hillsborough St., Raleigh, North Carolina 27606; and [§]Parasite Biology and Epidemiology Laboratory, Livestock and Poultry Sciences Institute, Agricultural Research Service, United States Department of Agriculture, BARC-East, Beltsville, Maryland 20705.

ABSTRACT: Serum samples from 143 hunter-killed black bears were collected during the 1996 and 1997 black bear hunting seasons in eastern North Carolina. All samples were tested for antibodies to *Toxoplasma gondii* by the modified agglutination test. Antibodies to *T. gondii* were present in 120 of 143 (84%) bears. Females had significantly higher titers than males (Wilcoxon rank sums test, $P = 0.045$), and titers increased with age (Jonckheere test, $P = 0.01$). Samples collected during 1996 ($n = 79$) were tested for antibodies to *Trichinella spiralis* by enzyme-linked immunosorbent assay. No samples were positive for antibodies to *T. spiralis*.

Toxoplasma gondii and *Trichinella spiralis* can cause severe disease in humans. Both can be acquired by consuming infective stages in undercooked meat. *Toxoplasma gondii* can also be contracted transplacentally or by consumption of food or water contaminated with oocysts (Dubey and Beattie, 1988). Historically, domestic animals have been considered the primary reservoir

hosts of these parasites for humans, but wildlife also play a role as sources of both human and domestic animal infection.

Numerous point-source outbreaks of human trichinellosis have been traced to bear meat (Wand and Lyman, 1972), and recently black bears (*Ursus americanus*) from Pennsylvania were reported to have 80% seroprevalence for *T. gondii* (Brice et al., 1993). Viable *T. gondii* was isolated from the hearts of 10 of 22 seropositive bears (Dubey et al., 1995). However the epizootiology of these parasites in black bears is not well understood. Opportunistic sampling of hunter-killed black bears in eastern North Carolina was undertaken to evaluate seroprevalence of *T. gondii* and *T. spiralis*. Hunters were also surveyed to investigate the disposition of the carcasses.

Blood samples were collected from the hearts of intact carcasses or from the body cavities of dressed carcasses of bears

collected in Hyde County (35°43'N to 35°04'N, 75°45'W to 75°39'W) and Tyrrell County (36°00'N to 35°34'N, 75°55'W to 75°38'W), North Carolina. From 11 November through 16 November 1996, 79 samples were collected (14 female, 65 male). From 10 November through 15 November 1997, 64 samples were collected (26 female, 38 male). Blood samples were placed in sterile 50-ml conical polypropylene centrifuge tubes and stored on ice until transported and processed at North Carolina State University, Raleigh, North Carolina. Blood was centrifuged and serum removed and frozen in polypropylene cryopreservation tubes at -20 C within 24 hr of collection. Gender, weight (intact carcasses only), and location of kill were recorded for each bear. A first premolar tooth was collected for sectioning and analysis of cementum annuli for age determination (Craighead et al., 1970). In addition, in 1996 a brief verbal survey was conducted with each hunter to collect information on intended uses of the carcass, sharing of meat, and meat storage and cooking methods (n = 79 complete surveys).

Frozen serum samples were sent to the Parasite Biology and Epidemiology Laboratory of the United States Department of Agriculture (USDA), Beltsville, Maryland, for analysis. Samples from 1996 and 1997 (n = 143) were tested for antibodies to *T. gondii* at dilutions of 1:25, 1:50, and 1:500 by the modified agglutination test (MAT) as described by Dubey and Desmonts (1987). Based on experience with MAT testing of human and animal sera, titers of $\geq 1:25$ were considered positive (Dubey et al., 1994, 1995). Samples collected in 1996 (n = 79) were also tested for antibodies to *T. spiralis* by enzyme-linked immunosorbent assay (ELISA) (Dubey et al., 1994, 1995). Positive and negative controls were included in each test and all samples were run by a single individual.

The Wilcoxon rank sum test (Hollander and Wolfe, 1973) was used to compare the *T. gondii* titers in male and female bears. The Jonckheere test for ordered alternatives was used to assess the association between age and titer (Hollander and Wolfe, 1973).

None of the 1996 samples was positive for antibodies to *T. spiralis*, and, therefore, further testing was not pursued in 1997. Antibodies to *T. gondii* were found in 84% (120 of 143) of sera tested. Prevalence of antibodies at the various terminal dilutions was 3.5% at 1:25 (n = 5), 61% at 1:50 (n = 87), and 19.5% at 1:500 (n = 28), while 16% (n = 23) of samples were negative. A significantly greater number of females than males were seropositive at the higher dilutions (Wilcoxon rank sums test, $P = 0.045$). Titers also increased with age (Jonckheere test, $P = 0.01$).

All hunters associated with the 79 bears sampled in 1996 completed questionnaires. Ninety-six percent of the hunters butchered the carcasses themselves and none wore gloves while skinning or handling the carcasses. All of the hunters surveyed consume the bear meat or share with others who do, and 20% prepare the meat in a manner that may be inadequate to inactivate infective stages of *T. gondii* or *T. spiralis*, such as preparing jerky or not cooking the meat thoroughly. Although no questions addressed the immunocompetency of the hunters surveyed, 3 volunteered information that they were immunocompromised, 1 due to immunosuppressive therapy following organ transplantation, and 2 due to cancer chemotherapy.

in the prevalence of trichinellosis in the U.S. swine population has been documented from about 1.4% at the turn of the century to 0.125% by the 1970s (Zimmerman, 1977). During the same period, the role of wildlife, particularly bears, as a major sylvatic reservoir of the disease and important source of human infection was recognized (Bailey and Schantz, 1990). Between 1982 and 1986, 14.6% of trichinellosis cases in the United States were traced to infected bear meat (Bailey and Schantz, 1990). The reported prevalence of *T. spiralis* in North American black bears ranges from 1.8% in Pennsylvania to 15.6% in Montana (Worley et al., 1991; Dubey et al., 1994).

The lack of antibodies to *T. spiralis* in this study was not anticipated. The sample size was relatively small and limited in geographic distribution and may not have had sufficient power to detect low seroprevalence. Alternatively, the absence of antibodies in black bears may reflect the historically low prevalence of *T. spiralis* in the North Carolina swine herd and the low numbers of commercial swine operations in the region surveyed (Cowen et al., 1990). Domestic swine have long been considered the major source of infection for wildlife that scavenge improperly dumped swine carcasses or scraps and refuse at garbage dumps. However, recent surveys of wildlife in Europe and the United States suggest that a sylvatic cycle of *Trichinella* infection operates largely independent from the domestic pig cycle (Pozio et al., 1992; Snyder et al., 1993). These studies also suggest that the dominant *Trichinella* species in wildlife may be sylvatic genotypes *Trichinella* T5 and T6 (Worley et al., 1990; Zarlenga et al., 1991).

Toxoplasmosis is a serious disease of humans and livestock, although infections in immunocompetent individuals are rare. Congenital infection can cause disease in the developing fetus, resulting in hydrocephaly, mental retardation, and other birth defects (Dubey and Beattie, 1988). Reported seroprevalence, by a variety of tests, for *T. gondii* in North American black bears has ranged from 8% to 80% across different regions of the country (Binnering et al., 1980; Briscoe et al., 1993). The results of this study are comparable to those recently reported for Pennsylvania black bears using the identical serologic test (Briscoe et al., 1993) and confirm that black bears have the highest seroprevalence of any animal or human population tested to date in the United States. While no testing for parasite burdens was undertaken for this study, previous work has shown the MAT to be the most consistent and reliable serologic test in black bears (Dubey et al., 1994, 1995). Sensitivity and specificity of the MAT in bears have not been fully evaluated but are established for pigs (Dubey, 1997; Dubey et al., 1997).

As omnivores, black bears consume both forage and foraging animals, as well as carrion and garbage. This may contribute to their high *T. gondii* seroprevalence, although the prevalence in North Carolina and Pennsylvania black bears is much higher than reported in other wild omnivores. A 1990 survey of free-ranging wild hogs (*Sus scrofa*) in the Great Smokey Mountains National Park, reported 31% seroprevalence by MAT (Diederich et al., 1996). A serosurvey of raccoons (*Procyon lotor*) from 7 states reported a mean prevalence of 50%, also by MAT (Dubey et al., 1992). The positive relationship between titer and the age of the bears suggests that the high seroprevalence may be due to cumulative exposure and supports similar observations from other studies (Briscoe et al., 1993; Zarnke et al., 1997). Signif-

bias introduced by the hunters. Larger, heavier bears are favored targets, and accordingly male bears are overrepresented in the sample. The larger female bears that are also favored by hunters tend to be older and, as previously demonstrated, titers increase with age.

The hunter survey suggests that bear hunters and those who handle and consume bear meat should be educated about the potential risks of exposure to *T. gondii* and how to minimize them. Such education could be accomplished by inclusion of a pamphlet on safe carcass handling with hunting licenses or dissemination of similar material to hunting clubs or hunting supply retailers. Important points to address include: (1) wearing gloves while butchering carcasses, (2) washing hands thoroughly with soap and water after handling bear meat, (3) cooking bear meat until welldone (to an internal temperature of 66°C for at least 3 min), and (4) burying viscera and carcass scraps to reduce scavenging by other animals.

The authors thank Scott Osborne, Mike Seamster, and Gordon Warburton (North Carolina Wildlife Resource Commission), students from North Carolina State University College of Veterinary Medicine, and Nick Vukich for their assistance with sample collection.

LITERATURE CITED

- BAILEY, T. M., AND P. M. SCHANTZ. 1990. Trends in the incidence and transmission patterns of trichinosis in humans in the United States: Comparisons of the periods 1975–1981 and 1982–1986. *Reviews of Infectious Diseases* **12**: 5–11.
- BINNINGER, C. E., J. J. BEECHAM, L. A. THOMAS, AND L. D. WINWARD. 1980. A serologic survey for selected infectious diseases of black bears in Idaho. *Journal of Wildlife Diseases* **16**: 423–430.
- BRISCOE, N., J. G. HUMPHREYS, AND J. P. DUBEY. 1993. Prevalence of *Toxoplasma gondii* infections in Pennsylvania black bears, *Ursus americanus*. *Journal of Wildlife Diseases* **29**: 599–601.
- COWEN, P., S. LI, AND T. MCGINN. 1990. Survey of trichinosis in breeding and cull swine, using an enzyme-linked immunosorbent assay. *American Journal of Veterinary Research* **51**: 924–928.
- CRAIGHEAD, J. J., F. C. CRAIGHEAD, JR., AND H. E. MCCUTCHEN. 1970. Age determination of grizzly bears from fourth premolar tooth sections. *Journal of Wildlife Management* **34**: 353–363.
- DIDERRICH, V. R., J. C. NEW, AND S. PATTON. 1996. Serologic survey of *Toxoplasma gondii* antibodies in free-ranging wild hogs (*Sus scrofa*) from the Great Smoky Mountains National Park and from sites in South Carolina. *Journal of Eukaryotic Microbiology* **43**: 122S.
- DUBEY, J. P. 1997. Validation of the specificity of the modified agglutination test for toxoplasmosis in pigs. *Veterinary Parasitology* **71**: 307–310.
- , C. D. ANDREWS, P. THULLIEZ, P. LIND, AND O. C. H. KWOK. 1997. Long-term humoral antibody responses by various serologic tests in pigs orally inoculated with oocysts of four strains of *Toxoplasma gondii*. *Veterinary Parasitology* **68**: 41–50.
- , AND C. P. BEATTIE. 1988. *Toxoplasmosis of animals and man*. CRC Press, Incorporated, Boca Raton, Florida, 220 pp.
- , N. BRISCOE, R. GAMBLE, D. ZARLENGA, J. G. HUMPHREYS, AND P. THULLIEZ. 1994. Characterization of *Toxoplasma* and *Trichinella* isolates from muscles of black bears in Pennsylvania. *American Journal of Veterinary Research* **55**: 815–819.
- , AND G. DESMONTS. 1987. Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Veterinary Journal* **19**: 337–339.
- , A. N. HAMIR, C. A. HANLON, AND C. E. RUPPRECHT. 1992. Prevalence of *Toxoplasma gondii* infection in raccoons. *Journal of the American Veterinary Medical Association* **200**: 534–536.
- , J. G. HUMPHREYS, AND P. THULLIEZ. 1995. Prevalence of viable *Toxoplasma gondii* tissue cysts and antibodies to *T. gondii* by various serologic tests in black bears (*Ursus americanus*) from Pennsylvania. *Journal of Parasitology* **81**: 109–111.
- HOLLANDER, M., AND D. A. WOLFE. 1973. *Nonparametric statistical methods*. John Wiley and Sons, New York, New York, 503 p.
- POZIO, E., G. LAROSA, P. ROSSI, AND K. D. MURRELL. 1992. Biological characterization of *Trichinella* isolates from various host species and geographical regions. *Journal of Parasitology* **78**: 647–653.
- SNYDER, D. E., D. S. ZARLENGA, G. LAROSA, AND E. POZIO. 1993. Biochemical, biological, and genetic characterization of a sylvatic isolate of *Trichinella*. *Journal of Parasitology* **79**: 347–352.
- WAND, M., AND D. LYMAN. 1972. Trichinosis from bear meat. *Journal of the American Medical Association* **220**: 245–246.
- WORLEY, D. E., F. M. SEESE, AND R. H. ESPINOSA. 1991. Prevalence and geographic distribution of *Trichinella spiralis* infection in hunter-killed bears in Montana, USA (1984–1989). *Helminthologia* **28**: 53–55.
- , D. S. ZARLENGA, AND F. M. SEESE. 1990. Freezing resistance of a *Trichinella spiralis* isolate from a gray wolf, *Canis lupus*, in Montana, with observations on genetic and biological characters of the biotype. *Journal of the Helminthological Society of Washington* **57**: 57–60.
- ZARLENGA, D. S., F. AL-YAMAN, AND D. J. MICHELLA. 1991. A repetitive DNA probe specific for a North American sylvatic genotype of *Trichinella*. *Molecular and Biochemical Parasitology* **48**: 131–138.
- ZARNKE, R. L., J. P. DUBEY, O. C. H. KWOK, AND J. M. VERHOEF. 1997. Serologic survey for *Toxoplasma gondii* in grizzly bears from Alaska. *Journal of Wildlife Diseases* **33**: 267–270.
- ZIMMERMAN, W. J. 1977. Trichinosis in bears of western and northcentral United States. *American Journal of Epidemiology* **106**: 167–171.

J. Parasitol., 84(5), 1998 p. 1050–1052
© American Society of Parasitologists 1998

Hosts Manipulated by One Parasite Incur Additional Costs from Infection by Another Parasite

R. Poulin, K. Hecker*, and F. Thomas, Department of Zoology, University of Otago, P.O. Box 56, Dunedin, New Zealand; *Department of Biology, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario, Canada K1S 5B6

ABSTRACT: Manipulation of host phenotype by parasites often serves to increase the predation rate of definitive hosts on intermediate hosts. For intermediate hosts, the indirect consequences of manipulation may extend beyond the direct increase in predation, however. Metacercariae of the trematode *Curtuteria australis* encyst in the foot of New Zealand cockles, *Austrovenus stutchburyi*, and stunt

its growth, rendering cockles incapable of burrowing into the sediments. Here, we show that cockles manipulated by *C. australis* are 5 times more likely to be infected by the castrating sporocysts of another trematode than normal, nonmanipulated cockles. Our results indicate that the consequences for *C. australis*-manipulated cockles are far more important than a simple increase in the risk of predation