- W. Yoder, Jr. Ames, Iowa State University Press. pp 471-484
- KOUWENHOVEN, B., DAVELAAR, F. G., BURGER, A. G. & LUTTICKEN, D. (1983) Significance of variant infectious bronchitis serotypes in The Netherlands. Proceedings of the 66th meeting of the Australian Poultry Association. August 31 to September 2, 1983. pp 53-71
- KUSTERS, J. G., NIESTERS, H. G. M., BLEUMINK-PLUYM, N. M. C., DAVELAAR, F. G., HORZINEK, M. C. & VAN DER ZEIJST, B. A. M. (1987) Molecular epidemiology of infectious bronchitis virus in The Netherlands. *Journal of General Virology* **68**, 343-352
- MARCHI, R. & ZANELLA, A. (1984) Avian infectious bronchitis. Investigation of a new serotype and its spreading. *Clinica Veterinaria* **107** 121-125
- PICAULT, J. P., DROUIN, P., LAMANDÉ, J., ALLEC, C., TOUX, J. Y., LE COQ, H., GUITTET, M. & BENNEJCAN, G. (1995) L'épizootie récente de Bronchite infectieuse aviaire en France: importance, évolution et étiologie. *Journées de la Recherche Avicole, Mars 1995*, 177-178
- REED, L. J. & MUENCH, H. (1938) A simple method of estimating fifty percent endpoint. *American Journal of Hygiene* **27**, 493-497
- RINALDI, A., CRESPI, A., CERVIO, G. & MANDELLI, G. (1966) Isolamento di un ceppo nefropatogeno del virus della bronchite infettiva del pollo. *Selezione Veterinaria* **7**, 284-287
- WANG, L., JUNKER, D. & COLLISON, E. W. (1993) Evidence of natural recombination within the S1 gene of infectious bronchitis virus. *Virology* **192**, 710-716
- ZANELLA, A. (1977) Bronchite infettiva nei polli da carne con particolare riferimento alle forme di nefrite-nefrosi. *Clinica Veterinaria* **100**, 407-415
- ZANELLA, A., GUALLINI, L. & MORINI, M. T. (1967) Caratteristiche dei ceppi nefropatogeni del virus della bronchite infettiva aviaria. *Proceedings of the Società Italiana delle Scienze Veterinarie* **21**, 895-899
- ZANELLA, A., MARCHI, R., MELLANO, D. & PONTI, W. (1988) Avian infectious bronchitis: nephropathogenic and respiratory viruses isolated and their spreading in Italy. Proceedings of the International Symposium on Infectious Bronchitis. Giessen, Germany. June 23-26. 1988. pp 245-255
- ZANELLA, A. & MARTINO (1998) Avian infectious bronchitis: persistence of nephropathogenic strains related to serotype AZ23/74. Proceedings of the International Symposium on Infectious Bronchitis and Pneumovirus Infections in Poultry. Giessen, Germany. June 15-18, 1998. pp 189-197

FIG 1: Vesicle (white arrowhead), ulcers and crusts (black arrowheads) present on the concave aspect of the ear pinna

Novel localised variant of canine epidermolysis bullosa acquisita

T. OLIVRY, A. PETERSEN, S. M. DUNSTON, M. CHEN

Veterinary Record (2000) **146**, 193-194

T. Olivry, DrVet, PhD, DipACVD, DipECVD, **S. M. Dunston**, BS, MS, Department of Clinical Sciences, College of Veterinary Medicine, North Carolina State University, Raleigh, North Carolina 27606, USA
A. Petersen, DrMedVet, Department of Small Animal Clinical Sciences, College of Veterinary Medicine, Michigan State University, East Lansing, Michigan, USA
M. Chen, PhD, LAC, University of Southern California, Los Angeles, California, USA

IN 1957, Brunsting and Perry described seven human patients with a chronically recurring blistering and scarring dermatosis preferentially localised to the head and neck (Brunsting and Perry 1957). These authors concluded that their patients were afflicted with a benign subset of bullous pemphigoid (BP) because of the observation of microscopic subepidermal vesicles (Brunsting and Perry 1957). This localised subset of BP was subsequently referred to as 'Brunsting-Perry cicatricial pemphigoid' (CP). In the past two decades, human beings with a clinical phenotype suggestive of Brunsting-Perry CP have been shown to exhibit skin-fixed and circulating immunoglobulin G (IgG) autoantibodies targeting antigens in anchoring fibrils (Dahl 1979, Gammon and others 1984, Kurzals and others 1991, Karpouzis and others 1993). These findings established that Brunsting-Perry CP actually represented a localised variant of epidermolysis bullosa acquisita (EBA). The latter, recognised in human beings and canine species, is an autoimmune blistering dermatosis that most commonly exhibits a generalised phenotype and seems refractory to therapy. EBA is characterised microscopically by neutrophil-rich

subepidermal vesicles and immunologically by autoantibodies directed against the non-collagenous amino terminus of type VII collagen (Gammon 1991, Olivry and others 1998). The aim of this study was to describe the clinical, histopathological and immunological features of a localised variant of canine EBA which resembled Brunsting-Perry CP in human beings.

An 11-month-old male, intact, German shorthaired pointer was presented with a 12-day history of cutaneous vesicles and ulcers. Initial lesions were scattered on the head and trunk and had rapidly decreased in severity following a short course of corticosteroids at anti-inflammatory doses. At the time of presentation to the dermatologist, vesicles, ulcers and scars were restricted to the concave aspect of the ear pinnae (Fig 1). All active cutaneous lesions progressively resolved with the administration of prednisone at low-immunosuppressive dosages (1 mg/kg two times a day). The dog was lost to follow-up after the initial remission was achieved.

Histopathological examination of lesional skin biopsy specimens demonstrated that subepidermal cleavage initially occurred with minimal dermal inflammation (Fig 2a). Multiple vesicles, however, were filled with neutrophil granulocytes and appeared as subepidermal microabscesses (Fig 2b). The observed microscopic lesions were indistinguishable from those seen in the skin of dogs with EBA (Olivry and others 1998). Antigen immunomapping, performed with collagen IV-specific monoclonal antibodies (Olivry and others 1998), corroborated that dermoepidermal cleavage occurred below the lamina densa of the epidermal basement membrane.

Direct immunofluorescence testing was performed on frozen skin sections as described by Olivry and others (1998). A very thick deposit of IgG was noticed at the epidermal basement membrane zone in both blistered and non-clefted areas (Fig 3). Basement membrane-fixed immunoglobulin A, immunoglobulin M and C3 were not seen. A high titre (>1:500) of circulating IgG anti-basement membrane autoantibodies was detected in this dog's serum using normal canine lip as a substrate. When salt-split canine lip epithelium was used (Iwasaki and others 1997, Olivry and others 1998), the IgG autoantibodies were shown to recognise one or more antigen located on the dermal side of artificially clefted skin. These results suggested that

FIG 2: (a) Rare subepidermal vesicles containing few inflammatory cells, (b) while others appear as neutrophil-rich subepidermal microabscesses. Haematoxylin and eosin. Bar = 100 µm

FIG 3: Thick linear and continuous deposit of immunoglobulin G present at the basement membrane zone (arrowhead) × 100

autoantibodies of the dog recognised antigens in or below the lamina densa.

To confirm the target specificity of this dog's serum IgG autoantibodies, an enzyme-linked immunosorbent assay (ELISA) was performed with recombinant human NC1 domain, the non-collagenous amino terminus of type VII collagen. This ELISA had been used as an aid to the diagnosis in human and canine patients affected by EBA and type I bullous systemic lupus erythematosus (BSLE) (Chen and others 1997, Olivry and others 1998, 1999). Moreover, recent cloning of canine NC1 yielded a sequence highly homologous to the human peptide, thereby confirming the validity of using a heterospecific substrate for NC1 immunoreactivity (Xu and others 1998). The ELISA studies confirmed that this dog's circulating IgG autoantibodies bound human recombinant collagen VII-NC1 with optical densities similar to those of other human and canine patients with EBA or BSLE (Olivry and others 1998, 1999). Conversely, the NC1 recombinant peptide was not recognised by circulating IgG from five normal control dogs. Since immunological investigations confirmed the presence of anti-collagen VII autoimmunity in the dog in this case, the diagnosis of EBA was confirmed (Gammon 1991).

The clinical phenotype exhibited by this dog, predominating to the head region and characterised by vesicles progressing to ulcers and scars, is reminiscent of that of Brunsting-Perry type EBA in human beings (Dahl 1979, Gammon and others 1984, Kurzals and others 1991, Karpouzis and others 1993). Moreover, the lesional distribution observed is remarkably less severe than that seen in other EBA-affected dogs which exhibit generalised and severe blistering and ulcerative lesions (Olivry and others 1998, T. Olivry, unpublished observations). This difference in clinical presentation justifies the recognition of a localised variant of EBA in the canine species.

ACKNOWLEDGEMENTS

The authors are grateful to Professor R. W. Dunstan for help with the initial histopathological diagnosis.

References

- BRUNSTING, L. A. & PERRY, H. O. (1957) Benign pemphigoid: a report of seven cases with chronic scarring, herpetiform plaques about the head and neck. *Archives of Dermatology* **75**, 489-501
- CHEN, M., CHAN, L. S., CAI, X. Y., O'TOOLE, E. A., SAMPLE, J. C. & WOODLEY, D. T. (1997) Development of an ELISA for rapid detection of anti-type VII collagen autoantibodies in epidermolysis bullosa acquisita. *Journal of Investigative Dermatology* **108**, 68-72
- DAHL, M. G. C. (1979) Epidermolysis bullosa acquisita – a sign of cicatricial pemphigoid. *British Journal of Dermatology* **101**, 475-484
- GAMMON, W. R. (1991) Epidermolysis bullosa acquisita: a disease of autoimmunity to type VII collagen. *Journal of Autoimmunity* **4**, 59-71
- GAMMON, W. R., BRIGGAMAN, R. A., WOODLEY, D. T., HEALD, P. W. & WHEELER, C. E. (1984) Epidermolysis bullosa acquisita – a pemphigoid-like disease. *Journal of the American Academy of Dermatology* **11**, 830-832
- IWASAKI, T., ISAJI, M., YANAI, T., KITAGAWA, H. & SASAKI, Y. (1997) Immunomapping of basement membrane zone macromolecules in canine salt-split skin. *Journal of Veterinary Medical Science* **59**, 391-393
- KARPOUZIS, A., PROST, C., CORDOLIANI, F., VIGUIER, D., BLUM, L. & PUISANT, A. (1993) Acquired localized epidermolysis bullosa. A case with scalp involvement and immunoelectron microscopic study. *Annales de Dermatologie et Vénéréologie* **120**, 464-468
- KURZALS, G., STOLZ, W., MEUER, M., KUNZE, J., BRAUN-FALCO, O. & KRIEG, T. (1991) Acquired epidermolysis bullosa with the clinical features of Brunsting-Perry cicatricial bullous pemphigoid. *Archives of Dermatology* **127**, 391-395
- OLIVRY, T., FINE, J.-D., DUNSTON, S. M., CHASSE, D., PASCAL TENORIO, A., MONTEIRO-RIVIERE, N. A., CHEN, M. & WOODLEY, D. T. (1998) Canine epidermolysis bullosa acquisita: circulating autoantibodies target the aminoterminal non-collagenous (NC1) domain of collagen VII in anchoring fibrils. *Veterinary Dermatology* **9**, 19-31
- OLIVRY, T., SAVARY, K. C. M., MURPHY, K. M., DUNSTON, S. M. & CHEN, M. (1999) Bullous systemic lupus erythematosus (type-1) in a dog. *Veterinary Record* **145**, 165-169
- XU, L. T., CHEN, M., PENG, J., O'TOOLE, E. A., WOODLEY, D. T. & CHAN, L. S. (1998) Molecular cloning and characterization of a cDNA encoding canine type VII collagen non-collagenous (NC1) domain, the target antigen of autoimmune disease epidermolysis bullosa acquisita (EBA). *Biochimica et Biophysica Acta – Molecular Basis of Disease* **1408**, 25-34

ABSTRACT

Ultrasonography for evaluating ovine joint disease

THE elbow, carpal, fetlock and stifle joints of six normal sheep were examined with a 7.5 MHz transducer with a stand-off pad, and the ultrasonographic appearance was compared with postmortem findings. In four of them the joints were examined postmortem before and after the joint capsules had been distended with 10 ml isotonic saline; this procedure made the capsules visible as a 2 mm thick echogenic line. Five sheep with chronic arthritis/synovitis were examined in the same way. The gross thickening of the joint capsule was visible as a hyperechoic band up to 20 mm thick. It is concluded that ultrasonography is useful in the evaluation of chronic joint disease in sheep, and may be valuable in assessing the prognosis for sheep with chronic arthritis/synovitis.

MACRAE, A. I. & SCOTT, P. R. (1999) *Veterinary Journal* **158**, 135