correct decay rates for $D = 0$, because any single point moment at time t depends on all the moments at $t = 0$.

The distribution function approach obviously preserves (2). Let $P'[\Psi|\mathbf{u}]$ denote the conditional probability functional for the concentration field $\psi(\mathbf{x})$, given the velocity field $\mathbf{u}(\mathbf{x})$. Consider the $\mathbf{x}$ space discrete so that elemental volume of Ψ space exists. The method of Edwards⁷ gives the following continuity equation:

$$\frac{\partial}{\partial t} P'[\Psi|\mathbf{u}] = \int d^3x \frac{\delta}{\delta\psi(\mathbf{x})} \cdot [C\psi^2(\mathbf{x}) + \mathbf{u}(\mathbf{x}) \cdot \nabla\psi(\mathbf{x}) - D\nabla^2\psi(\mathbf{x})] P'[\Psi|\mathbf{u}], \quad (5)$$

where δ/δ denotes functional differentiation. The single point probability density function is obtained by multiplying (5) by $F(\psi)$, the probability density functional of Ψ for all $\mathbf{x}$ and t , and integrating over all ψ and over Ψ for all points but $\mathbf{x}$, resulting in

$$\begin{aligned} & \left(\frac{\partial}{\partial t} - C \frac{\partial}{\partial\psi} \psi^2 \right) P'_x(\psi) \\ &= -D \lim_{\mathbf{x}' \rightarrow \mathbf{x}} \frac{\partial}{\partial\psi} \nabla_{\mathbf{x}'}^2 \cdot \int d\psi' \psi' P'_{x,x'}(\psi, \psi') \\ & \quad - \int d^3u \mathbf{u} \cdot \nabla P'_{x,x}(\psi, \mathbf{u}) \end{aligned} \quad (6a)$$

which reduces to (2) if $D = 0$ and if Γ and $\mathbf{U}$ are jointly homogeneous. An unclosed hierarchy of equations for the multipoint probability density functions can be set up analogous to that for the ordinary turbulence problem,^{8,9} the next member being

$$\begin{aligned} & \left[\frac{\partial}{\partial t} - C \left(\frac{\partial}{\partial\psi} \psi^2 + \frac{\partial}{\partial\psi'} \psi'^2 \right) \right] P'_{x,x'}(\psi, \psi') \\ &= -D \left\{ \lim_{\mathbf{x}'' \rightarrow \mathbf{x}} \frac{\partial}{\partial\psi} + \lim_{\mathbf{x}'' \rightarrow \mathbf{x}'} \frac{\partial}{\partial\psi'} \right\} \nabla_{\mathbf{x}''}^2 \cdot \\ & \quad \cdot \int d\psi'' \psi'' P'_{x,x',x''}(\psi, \psi', \psi'') \\ & \quad - \int d^3u \mathbf{u} \cdot [\nabla_x P'_{x,x',x}(\psi, \psi', \mathbf{u}) \\ & \quad + \nabla_{x'} P'_{x,x',x}(\psi, \psi', \mathbf{u})]. \end{aligned} \quad (6b)$$

The velocity terms may be treated by considering the parallel hierarchy of joint probability density functions $P'_{x,x}(\psi, \mathbf{u})$, etc., also obtained from (5) without use of the Navier-Stokes equations and formally similar to Eqs. (6) but without a $\mathbf{u}$ integration. Closure may be obtained by parameterization of assumed higher order joint probability density functions.

Let us turn our attention to the reciprocal of the concentration of the contaminant, its specific

volume $\Upsilon(\mathbf{x}, t) \equiv \Gamma^{-1}(\mathbf{x}, t)$ (use Υ^{N-1} for N th order reactions), whose dynamical equation is

$$\Re(\mathbf{x}, t) \Upsilon(\mathbf{x}, t) = C - \frac{2D}{\Upsilon(\mathbf{x}, t)} |\nabla \Upsilon(\mathbf{x}, t)|^2. \quad (7)$$

The logarithmic nonlinearity in (7) is present in single point hierarchy equations for both (1) and (7),

$$\Re \Gamma^n = -n C \Gamma^{n+1} - (n-1) D (\nabla \log \Gamma) \cdot (\nabla \Gamma^n),$$

$$\Re \Upsilon^n = n C \Upsilon^{n-1} - (n+1) D (\nabla \log \Upsilon) \cdot (\nabla \Upsilon^n).$$

The dissipative term is usually avoided by considering the Γ hierarchy of multiple point correlations. After closure certain points are made to coincide before substitution into lower order equations. Unfortunately, the log term appears in the first equation of the Υ hierarchy and cannot be avoided. As a rough approximation one could replace Υ^{-1} by $\langle \Upsilon \rangle^{-1}$ and use a moment closure. Because the dissipative term disappears when $D = 0$, O'Brien's invariance is automatically satisfied. Furthermore, when the turbulence is also homogeneous the hierarchy of $\langle \Upsilon^n \rangle$ equations is closed at each level, giving the exact result $d\langle \Upsilon \rangle/dt = C$ and all the central moments of Υ constant in time.

The author is grateful for many beneficial consultations with Dr. J. R. Herring.

* National Research Council-National Aeronautics and Space Administration Resident Research Associate, now at Shell Development Company, Emeryville, California.

¹ E. E. O'Brien, *Phys. Fluids* **9**, 1561 (1966).

² E. E. O'Brien, *Phys. Fluids* **11**, 2087 (1968).

³ S. Corrsin, *Phys. Fluids* **1**, 42 (1958).

⁴ For further details of this note see J. C. Hill, *Goddard Space Flight Center Document X-641-69-108* (1969).

⁵ See also Eq. (42) of S. Corrsin, in *Mécanique de la Turbulence*, A. Favre, Ed. (Centre National de la Recherche Scientifique, Paris, 1962), p. 27.

⁶ E. E. O'Brien, *Phys. Fluids* **11**, 2328 (1968).

⁷ S. F. Edwards, *J. Fluid Mech.* **18**, 239 (1964).

⁸ T. S. Lundgren, *Phys. Fluids* **10**, 969 (1967).

⁹ A. S. Monin, *Prikl. Mat. Mekh.* **31**, 1057 (1967) [*J. Appl. Math. Mech.* **31**, 1057 (1967)].

Direct Correlation Function in Space and Time

F. LADO

*Department of Physics, North Carolina State University
Raleigh, North Carolina
(Received 29 August 1969)*

A new, simpler derivation is presented of the equation suggested by Percus and Yevick to define the generalized direct correlation function in space and time. In addition, an explicit, though formal, definition of this function is given.

The time-dependent generalization of the total correlation function $G(\mathbf{r}) \equiv g(\mathbf{r}) - 1$ used in the

classical equilibrium theory of fluids is the density-density correlation function¹

$$\hat{G}(\mathbf{r}, t) \equiv (nN)^{-1} \int d\mathbf{r}' \langle R(\mathbf{r}', 0) R(\mathbf{r}' + \mathbf{r}, t) \rangle, \quad (1)$$

where

$$R(\mathbf{r}, t) = \exp(-t\mathcal{L})R(\mathbf{r}, 0), \quad (2)$$

$$R(\mathbf{r}, 0) = \sum_{i=1}^N \delta(\mathbf{r} - \mathbf{r}_i) - n, \quad (3)$$

and where $\mathcal{L}$ is the Liouville operator (i.e., $\mathcal{L}F$ is the Poisson bracket of the Hamiltonian H with F), N is the number of particles in the volume V , and n is the mean density. The brackets denote an equilibrium ensemble average. The initial value of $\hat{G}(\mathbf{r}, t)$ is then, from (1),

$$\hat{G}(\mathbf{r}, 0) = G(\mathbf{r}) + n^{-1} \delta(\mathbf{r}). \quad (4)$$

Recently, Percus and Yevick² have suggested a defining equation for the time-dependent extension $\hat{C}(\mathbf{r}, t)$ of the equilibrium direct correlation function $C(\mathbf{r})$, to wit,

$$\frac{\partial \hat{G}(\mathbf{r}, t)}{\partial t} = -\frac{n}{m\beta} \nabla^2 \int_0^t dt' \cdot \int d\mathbf{r}' \hat{C}(\mathbf{r}', t') \hat{G}(\mathbf{r} - \mathbf{r}', t - t'), \quad (5)$$

where m is the mass of a particle and β is $(k_B T)^{-1}$ and where initially

$$\hat{C}(\mathbf{r}, 0) = C(\mathbf{r}) - n^{-1} \delta(\mathbf{r}). \quad (6)$$

It is the purpose of this note to present a simpler derivation of Eqs. (5) and (6) than that given in Ref. 2 and in the process to obtain an explicit, though formal, expression for the function $\hat{C}(\mathbf{r}, t)$.

The derivation is a straightforward application of Zwanzig's³ projection operator technique to the Fourier transform of Eq. (1),

$$\hat{G}_k(t) = (nN)^{-1} \langle R_{-k}(0) R_k(t) \rangle. \quad (7)$$

[We write the transform F_k of any function $F(\mathbf{r})$ defined in V such that

$$F(\mathbf{r}) = (2\pi)^{-3} \int d\mathbf{k} F_k \exp(i\mathbf{k} \cdot \mathbf{r}),$$

$$F_k = \int d\mathbf{r} F(\mathbf{r}) \exp(-i\mathbf{k} \cdot \mathbf{r}), \quad (8)$$

for infinitely large V .] To evaluate the average in Eq. (7) we do not need to know the general time behavior of $R_k(t)$ but rather only that of its projection onto the "fixed state" $R_k(0)$. To this end, we define, following Zwanzig, a projection operator $\mathcal{P}$ such that for any phase-dependent quantity Q

$$\mathcal{P}Q \equiv \frac{\langle R_{-k}(0)Q \rangle}{\langle R_{-k}(0)R_k(0) \rangle} R_k(0). \quad (9)$$

Note that we may then write

$$\hat{G}_k(t) = (nN)^{-1} \langle R_{-k}(0) \mathcal{P}R_k(t) \rangle, \quad (10)$$

and so

$$\partial \hat{G}_k(t) / \partial t = (nN)^{-1} \langle R_{-k}(0) \partial \mathcal{P}R_k(t) / \partial t \rangle. \quad (11)$$

The equation of motion of the complete $R_k(t)$,

$$\partial R_k(t) / \partial t = -\mathcal{L}R_k(t), \quad (12)$$

is obtained immediately from Eq. (2). To find the corresponding equation for just the projected part $\mathcal{P}R_k(t)$ we operate on Eq. (12) with $\mathcal{P}$ and with $(1-\mathcal{P})$, obtaining two equations linking the projected part and the "remainder" $(1-\mathcal{P})R_k(t)$. The "remainder" is eliminated, yielding a closed, formal equation for $\mathcal{P}R_k(t)$; we refer to Zwanzig³ for the details of this calculation. The result is

$$\frac{\partial \mathcal{P}R_k(t)}{\partial t} = \int_0^t dt' \mathcal{P} \mathcal{L} \exp[-t'(1-\mathcal{P})\mathcal{L}] \mathcal{L}R_k(0) \cdot \frac{\langle R_{-k}(0)R_k(t-t') \rangle}{\langle R_{-k}(0)R_k(0) \rangle}, \quad (13)$$

so that, with Eq. (11), we have

$$\begin{aligned} \frac{\partial \hat{G}_k(t)}{\partial t} &= \int_0^t dt' \frac{\langle R_{-k}(0) \mathcal{L} \exp[-t'(1-\mathcal{P})\mathcal{L}] \mathcal{L}R_k(0) \rangle}{\langle R_{-k}(0)R_k(0) \rangle} \\ &\quad \cdot \hat{G}_k(t-t') \\ &= - \int_0^t dt' K_k(t') \hat{G}_k(t-t'), \end{aligned} \quad (14)$$

where we have put

$$K_k(t) \equiv \frac{\langle \mathcal{L}R_{-k}(0) \exp[-t(1-\mathcal{P})\mathcal{L}] \mathcal{L}R_k(0) \rangle}{\langle R_{-k}(0)R_k(0) \rangle}. \quad (15)$$

The initial value of the kernel $K_k(t)$ may easily be evaluated

$$K_k(0) = \frac{k^2}{m\beta n \hat{G}_k(0)}. \quad (16)$$

But, in the equilibrium theory of fluids the direct correlation function is defined by

$$(1 + nG_k)^{-1} \equiv [n\hat{G}_k(0)]^{-1} = 1 - nC_k, \quad (17)$$

so that (16) may also be written

$$K_k(0) = -\frac{nk^2}{m\beta} \left(C_k - \frac{1}{n} \right). \quad (18)$$

In general, we define a function $\hat{C}_k(t)$ such that

$$K_k(t) = -\frac{nk^2}{m\beta} \hat{C}_k(t). \quad (19)$$

Then (14) becomes

$$\frac{\partial \hat{G}_k(t)}{\partial t} = \frac{nk^2}{m\beta} \int_0^t dt' \hat{C}_k(t') \hat{G}_k(t - t'), \quad (20)$$

the inverse transform of which is just Eq. (5). Thus, we have verified Percus and Yevick's suggestion that Eq. (5) can serve to define a time-dependent direct correlation function which reduces to the equilibrium function at the initial time. In addition, we are able to give the explicit definition of this function, namely,

$$\hat{C}_k(t) \equiv -\frac{1}{n^2 \hat{G}_k(0)} \frac{\langle \mathcal{L}R_{-k}(0) \exp[-t(1 - \mathcal{P})\mathcal{L}] \mathcal{L}R_k(0) \rangle}{\langle \mathcal{L}R_{-k}(0) \mathcal{L}R_k(0) \rangle}. \quad (21)$$

And finally, it is evident that this result is but a special case of the general formalism for time correlation functions developed by Zwanzig.³

This work was supported by the North Carolina Engineering Foundation.

¹ L. van Hove, Phys. Rev. **95**, 249 (1954).

² J. K. Percus and G. J. Yevick, Phys. Fluids **11**, 2050 (1968).

³ R. Zwanzig, in *Lectures in Theoretical Physics*, W. E. Brittin, W. B. Downs, and J. Downs, Eds. (Interscience Publishers, Inc., New York, 1961) Vol. 3, p. 135.

Blast Waves and Scaling Laws

DONALD L. JONES

General Electric Company, Philadelphia, Pennsylvania
(Received 11 August 1969; final manuscript received
3 December 1969)

Spherical blast wave scaling is studied with the intermediate strength formalism. Simple scaling relations rarely exist, but the intermediate strength equations permit a direct comparison of arrival times and front overpressures for different explosions.

In the evaluation of the effects of explosions under one set of conditions, it is often convenient to compare them with the measured effects of a known explosion under different conditions. The need to compare, rather than compute, explosion effects arises from the complexity of the fluid equations of motion, which are nonlinear. Only for the case of strong blast waves ($\Delta P/P_0 > 10$) is there an analytical, closed form solution.¹ Since the region $\Delta P/P_0 < 10$ is quite important (some shock damage occurs for overpressures as low as 1 psi), and since the shock spends over 95% of its lifetime in the overpressure region between 10 atm and 1 psi, a scaling approach developed by Sachs² has been widely used. Although Sachs

scaling is a general procedure, its principal application has been in relating the distance from the origin for the occurrence of a given overpressure³

$$\frac{D}{D_1} = \left(\frac{W}{W_1} \right)^{1/3}, \quad (1)$$

where D is the distance from the origin to the shock front and W is the energy. Other important relationships are the variation of the overpressure at a particular location or the arrival time ratio. Unfortunately, such relations are not easily found with scaling relations.

It is the purpose of this note to set down explicitly certain "scaling" relations derived from the intermediate strength blast relations.⁴ For brevity we limit this discussion to spherical blasts, although both the cylindrical and plane cases could be handled in a similar manner.

From Jones⁴ we write the equation for the trajectory of the shock front for a spherical blast

$$\tau = 0.543[(1 + 4.61X^{5/2})^{2/5} - 1], \quad (2)$$

and the overpressure at the front is

$$\frac{\Delta P}{P_0} = \frac{2\gamma}{\gamma + 1} \frac{4}{25} \{0.611[(1 + 5.89X^3)^{5/18} - 1]\}^{-1}, \quad (3)$$

where $\tau = a_0 t/R_0$, $X = D/R_0$, a_0 is the speed of sound in the undisturbed medium, R_0 is the characteristic radius $[2.5(1/B)(1/\gamma)(W/P_0)]^{1/3}$, D is the distance from the shock front to the source, B is a constant,⁵ γ is the specific heat ratio, P_0 is the ambient pressure, and W is the energy input to the shock from the explosion. For strong blast waves $\tau = X^{5/2}$ and $\Delta P/P_0 = [2\gamma/(\gamma + 1)]^{4/5} X^{-3}$.

As an example, we derive the scaling relation for the distance at which a given overpressure will occur, assuming different energy releases. Utilizing (3) and some algebra we find $\Delta P/\Delta P_1 = X_1^3/X^3$. But $\Delta P = \Delta P_1$; $X_1 = X$. From the definition of X , we now write $D/D_1 = R_0/R_{01}$. This is equivalent to $D/D_1 = (W/W_1)^{1/3}$, which is in accord with Ref. 2 [Eq. (3.56.1)]. Following the same procedure leads to the relations shown in Table I. In addition to the relationship between the given variables and the intermediate formalism, the limiting cases of strong and weak shocks (columns 4 and 5) are provided for comparison.

From Table I, it is clear that only relation (a) is constant for all shocks. The remaining situations have scaling factors that depend on the shock strength; however, Eqs. (2) and (3) provide a means of comparing shock front effects throughout the range of overpressures to as low as 1 psi.

¹ G. I. Taylor, Proc. Roy. Soc. (London) **A201**, 159 (1950).

² U. Ericsson and K. Edin, Phys. Fluids **3**, 893 (1960).

³ S. Glasstone, Ed., *The Effects of Nuclear Weapons* (U. S.