MOMENTS OF ORDER STATISTICS FROM A NORMAL POPULATION

by

R. C. Bose and Shanti S. Gupta.

University of North Carolina.

This research was supported by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command under contract No. AF 18 (600) - 83.

Institute of Statistics
Mimeograph Series No. 154
July, 1956

UN CLASSIFIED

Security Information

Bibliographical Control Sheet

1. University of North Carolina, Chapel Hill, N. C.
2. Mathematics Division, Air Force Office of Scientific Research
3. AFOSR-TN-56-501
4. ASTIA AD 110 316
5. MOMENTS OF ORDER STATISTICS FROM A NORMAL POPULATION
6. R. C. Bose and Shanti S. Gupta
7. October, 1956
8. AF 18(600)-83
9. File 3.3

MOMENTS OF ORDER STATISTICS FROM A NORMAL POPULATION

By

R. C. Bose and Shanti S. Gupta

University of North Carolina

1. Introduction and summary.

Statistics based on ordered observations have been called systematic statistics by Mosteller [12]. They are now being increasingly used in new statistical procedures [1, 2, 3, 4, 7, 8, 13, 14, 15, 16, 17, 20, 22]. The present paper deals with the problem of obtaining the moments of $X_{(k)}$, the k -th order statistic for a sample of size n from a normal population $N(0, 1)$. This problem has been considered among others by Cole [5], Godwin [6], Hastings, Mosteller, Tukey and Winsor [9], Jones [11], Ruben [18] and Tippet [23].

It has been shown that $\mu_t'(n, k)$, the t -th moment of $X_{(k)}$, can be expressed in terms of lower moments of order $t - 2i$ ($i = 1, 2, \dots, t/2$ or $(t - 1)/2$) and the integral

$$(1.1) \quad \int_{-\infty}^{+\infty} P_{t+1}(x) e^{-(t+1)x^2/2} dx$$

where $P_{t+1}(x)$ for $t > 0$, is defined by

$$(1.2) \quad P_{t+1}(x) = k \binom{n}{k} \frac{d^t}{d\phi^t} [\phi^{k-1} (1 - \phi)^{n-k}] ,$$

it being understood that in (1.2), ϕ is replaced after differentiation by $\Phi(x)$, the c. d. f. of $N(0, 1)$. $P_t(x)$ is thus a polynomial of degree $(n - t)$ in $\Phi(x)$ if $n \leq t$ and is zero if $n > t$. Exact values of all odd order moments can be derived when $n \leq 5$, and the exact values of all even order moments can be derived when $n \leq 6$. Godwin [6] and Jones [11] have given tables of exact moments $\mu_t'(n, k)$ for $t = 1$ and 2 . The corresponding tables for $t = 3$ and 4 are provided

in this paper. In general the numerical evaluation of the integral (1.1) can be expeditiously done by using the Gauss-Jacobi method of mechanical quadrature [21] based on the zeros and the weight factors of the Hermite-polynomials for which tables have been provided by Salzer, Zucker and Capuano [19]. It is believed that the formulae derived here are better suited for numerical computation than those given elsewhere.

2. The function $P_t(n, k, x)$.

Let $x_1, x_2, \dots, x_n$ be n independent observations from a normal population $N(0, 1)$ with zero mean and unit variance, and let

$$(2.1) \quad x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}$$

be the n ranked observations among $x_1, x_2, \dots, x_n$. Then the cumulative distribution function of $X_{(k)}$, the random variable corresponding to $x_{(k)}$, ($1 \leq k \leq n$), is given by

$$(2.2) \quad P_0(n, k, x) = \text{Prob} \left\{ X_{(k)} \leq x \right\} \\ = \frac{C}{(2\pi)^{1/2}} \int_{-\infty}^x [\Phi(x)]^{k-1} [1 - \Phi(x)]^{n-k} e^{-x^2/2} dx$$

where $\Phi(x)$ is defined as

$$(2.3) \quad \Phi(x) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^x e^{-x^2/2} dx$$

and C is the constant

$$(2.4) \quad C = \frac{n!}{(k-1)! (n-k)!} .$$

UN CLASSIFIED

Security Information

Bibliographical Control Sheet

1. University of North Carolina, Chapel Hill, N. C.
2. Mathematics Division, Air Force Office of Scientific Research
3. AFOSR-TN-56-501
4. ASTIA AD 110 316
5. MOMENTS OF ORDER STATISTICS FROM A NORMAL POPULATION
6. R. C. Bose and Shanti S. Gupta
7. October, 1956
8. AF 18(600)-83
9. File 3.3

Let us now define the function $P_t(n, k, x)$, which we shall abbreviate to $P_t(x)$ for convenience, by the relation

$$(2.5) \quad P_{t+1}(x) = (2\pi)^{1/2} e^{x^2/2} \frac{dP_t}{dx}$$

where $P_0(x)$ is given by (2.2). Then

$$(2.6) \quad P_1(x) = C \int \phi(x) \int^{k-1} \int^{1-\phi(x)} \int^{n-k}.$$

It is clear that $P_t(x)$ is a polynomial of degree $n-t$ in $\phi(x)$ if $1 \leq t \leq n$, and is zero for $t > n$. In fact, we can write

$$(2.7) \quad P_{t+1}(x) = C \frac{d^t}{d\phi^t} \int \phi^{k-1} (1-\phi)^{n-k} \int$$

where ϕ is replaced by $\phi(x)$ after the differentiation. It follows that for given t, n, k , $\phi_t(x)$ is a bounded function of x . The functions $P_2(x)$, $P_3(x)$, $P_4(x)$ and $P_5(x)$ are given below explicitly, where ϕ is written for $\phi(x)$.

$$(2.8) \quad P_2(x) = C \phi^{k-2} (1-\phi)^{n-k-1} \int (k-1) - (n-1) \phi \int$$

$$(2.9) \quad P_3(x) = C \phi^{k-3} (1-\phi)^{n-k-2} \int (k-1)(k-2) - 2(k-1)(n-2)\phi + (n-1)(n-2)\phi^2 \int$$

$$(2.10) \quad P_4(x) = C \phi^{k-4} (1-\phi)^{n-k-3} \int (k-1)(k-2)(k-3) - 3(k-1)(k-2)(n-3)\phi \\ + 3(k-1)(n-2)(n-3)\phi^2 - (n-1)(n-2)(n-3)\phi^3 \int$$

$$(2.11) \quad P_5(x) = C \phi^{k-5} (1-\phi)^{n-k-4} \int (k-1)(k-2)(k-3)(k-4) - 4(k-1)(k-2)(k-3)(k-4)\phi \\ + 6(k-1)(k-2)(n-3)(n-4)\phi^2 \\ - 4(k-1)(n-2)(n-3)(n-4)\phi^3 + (n-1)(n-2)(n-3)(n-4)\phi^4 \int$$

3. A system of differential equations satisfied by $P_0(x)$.

From (2.5)

$$(3.1) \quad P_1(x) = (2\pi)^{1/2} e^{x^2/2} \frac{dP_0}{dx}$$

$$(3.2) \quad \begin{aligned} P_2(x) &= (2\pi)^{1/2} e^{x^2/2} \frac{d}{dx} \left[(2\pi)^{1/2} e^{x^2/2} \frac{dP_0}{dx} \right] \\ &= (2\pi) e^{x^2} \left[\frac{d^2 P_0}{dx^2} + x \frac{dP_0}{dx} \right]. \end{aligned}$$

In general let us assume

$$(3.3) \quad P_t(x) = (2\pi)^{t/2} e^{tx^2/2} \sum_{r=0}^{t-1} g_{r,t}(x) \frac{d^{t-r} P_0}{dy^{t-r}}$$

where

$$(3.4) \quad g_{0,t}(u) = 0$$

and $g_{r,t}(x)$ is a polynomial in x of the r -th degree. Differentiating (3.3) and using (2.5), we have

$$(3.5) \quad P_{t+1}(x) = (2\pi)^{(t+1)/2} e^{(t+1)x^2/2} \sum_{r=0}^{t-1} \left[g_{r,t}(x) \frac{d^{t-r+1} P_0}{dx^{t-r+1}} + \left\{ tx g_{r,t}(x) + g'_{r,t}(x) \right\} \frac{d^{t-r} P_0}{dx^{t-r}} \right].$$

This leads to the recurrence relation

$$(3.6) \quad g_{r,t+1}(x) = g_{r,t}(x) + \left\{ tx + \frac{d}{dx} \right\} g_{r-1,t}(x)$$

where $g_{r,r}(x)$ should be interpreted as zero. This together with (3.4) determines all the polynomials $g_{r,t}(x)$. Starting from

$$(3.7) \quad g_{0,1}(x) = 1$$

we can successively calculate

$$(3.8) \quad g_{0,2}(x) = 1, \quad g_{1,2}(x) = x$$

$$(3.9) \quad g_{0,3}(x) = 1, \quad g_{1,3}(x) = 3x, \quad g_{2,3}(x) = 2x^2 + 1$$

$$(3.10) \quad g_{0,4}(x) = 1, \quad g_{1,4}(x) = 6x, \quad g_{2,4}(x) = 11x^2 + 4, \quad g_{3,4}(x) = 6x^3 + 7x$$

$$(3.11) \quad g_{0,5}(x) = 1, \quad g_{1,5}(x) = 10x, \quad g_{2,5}(x) = 35x^2 + 10, \quad g_{3,5}(x) = 50x^3 + 45x$$

$$g_{4,5}(x) = 24x^4 + 46x^2 + 7.$$

Hence we have the set of equations

$$(3.12) \quad \frac{dP_0}{dx} = \frac{1}{(2\pi)^{1/2}} e^{-x^2/2} P_1(x)$$

$$(3.13) \quad \frac{d^2 P_0}{dx^2} + x \frac{dP_0}{dx} = \frac{1}{(2\pi)} e^{-x^2} P_2(x)$$

$$(3.14) \quad \frac{d^3 P_0}{dx^3} + 3x \frac{d^2 P_0}{dx^2} + (2x^2 + 1) \frac{dP_0}{dx} = \frac{1}{(2\pi)^{3/2}} e^{-3x^2/2} P_3(x)$$

$$(3.15) \quad \frac{d^4 P_0}{dx^4} + 6x \frac{d^3 P_0}{dx^3} + (11x^2 + 4) \frac{d^2 P_0}{dx^2} + (6x^3 + 7x) \frac{dP_0}{dx} = \frac{1}{(2\pi)^2} e^{-2x^2} P_4(x)$$

$$(3.16) \quad \frac{d^5 P_0}{dx^5} + 10x \frac{d^4 P_0}{dx^4} + (35x^2 + 10) \frac{d^3 P_0}{dx^3} + (50x^3 + 45x) \frac{d^2 P_0}{dx^2} + (24x^4 + 46x^2 + 7) \frac{dP_0}{dx} \\ = \frac{1}{(2\pi)^{5/2}} e^{-5x^2/2} P_5(x).$$

We can proceed in this manner up to any order but it should be noted that $P_n(x)$ is a constant and $P_t(x) = 0$ if $t > n$. The general equation is

$$(3.17) \quad g_{0,t}(x) \frac{d^t P_0}{dx^t} + g_{1,t}(x) \frac{d^{t-1} P_0}{dx^{t-1}} + \dots + g_{t-1,t}(x) \frac{dP_0}{dx} = \frac{1}{(2\pi)^{t/2}} e^{-tx^2/2} P_t(x).$$

4. Moments of $X_{(k)}$.

We shall first prove the following Lemma:

Lemma. If α and r are non-negative integers, then

$$(4.1) \quad \int_{-\infty}^{+\infty} x^{\alpha} \frac{d^{r+1}P_0}{dx^{r+1}} dx = (-1)^r \alpha(\alpha-1) \dots (\alpha-r+1) \mu'_{\alpha-r} \quad \text{or } 0$$

according as

$$r \leq \alpha \quad \text{or} \quad r > \alpha$$

where $\mu'_{\alpha-r}$ is the $(\alpha-r)$ -th order moment of $X_{(k)}$ about the origin.

It should be noted that by definition

$$(4.2) \quad \int_{-\infty}^{+\infty} x^{\alpha} \frac{dP_0}{dx} dx = \mu'_{\alpha}.$$

From (3.12) and (3.13)

$$(4.3) \quad \frac{dP_0}{dx} = \frac{1}{(2\pi)^{1/2}} e^{-x^2/2} P_1(x)$$

$$(4.4) \quad \frac{d^2P_0}{dx^2} = -\frac{x}{(2\pi)^{1/2}} e^{-x^2/2} P_1(x) + \frac{1}{2\pi} e^{-x^2/2} P_2(x)$$

and in general using the system of equations (3.12)-(3.17) we can write

$$(4.5) \quad \frac{d^r P_0}{dx^r} = \sum_{i=1}^r h_i(x) e^{-x^2/2} P_i(x)$$

where $h_i(x)$ is a polynomial in x . Now

$$(4.6) \quad \int_{-\infty}^{+\infty} x^{\alpha} \frac{d^{r+1}P_0}{dx^{r+1}} dx = \left| x^{\alpha} \frac{d^r P_0}{dx^r} \right|_{-\infty}^{+\infty} - \int_{-\infty}^{+\infty} \alpha x^{\alpha-1} \frac{d^r P_0}{dx^r} dx.$$

Since $P_t(x)$ for any non-negative integer t is a bounded function of x , it follows from (4.5) that the first part on the right hand side of (4.6) vanishes.

Repeating this process we get if $r \leq \alpha$

$$\int_{-\infty}^{+\infty} x^\alpha \frac{d^{r+1}P_0}{dx^{r+1}} dx = (-1)^r \alpha(\alpha-1)\dots(\alpha-r+1) \int_{-\infty}^{+\infty} x^{\alpha-r} \frac{dP_0}{dx} = (-1)^r \alpha(\alpha-1)\dots(\alpha-r+1) \mu'_{\alpha-r}.$$

If $r > \alpha$, we get on repeating the process α times,

$$\begin{aligned} \int_{-\infty}^{+\infty} x^\alpha \frac{d^{r+1}P_0}{dx^\alpha} dx &= (-1)^\alpha \alpha(\alpha-1)\dots 3.2.1 \int_{-\infty}^{+\infty} \frac{d^{r+1-\alpha}P_0}{dx^{r+1-\alpha}} dx \\ &= (-1)^\alpha \alpha(\alpha-1)\dots 3.2.1 \left| \frac{d^{r-\alpha}P_0}{dx^{r-\alpha}} \right|_{-\infty}^{+\infty} \\ &= 0. \end{aligned}$$

This proves the Lemma.

On applying the Lemma and integrating the equations (3.13) ... (3.16) we get

$$(4.7) \quad \mu'_1 = \frac{1}{2\pi} \int_{-\infty}^{+\infty} P_2(x) e^{-x^2} dx$$

$$(4.8) \quad -3 + (2\mu'_2 + 1) = \frac{1}{(2\pi)^{3/2}} \int_{-\infty}^{+\infty} P_3(x) e^{-3x^2/2} dx$$

$$(4.9) \quad -2\mu'_1 + (6\mu'_3 + 7\mu'_1) = \frac{1}{(2\pi)^2} \int_{-\infty}^{+\infty} P_4(x) e^{-2x^2} dx$$

$$(4.10) \quad 70 - 150\mu'_2 - 45 + (24\mu'_4 + 46\mu'_2 + 7) = \frac{1}{(2\pi)^{5/2}} \int_{-\infty}^{+\infty} P_5(x) e^{-5x^2/2} dx.$$

We may write $\mu'_{\alpha}(n,k)$ instead of μ'_{α} to denote the fact that we have the α -th moment about the origin of the k -th order statistic out of a sample of n observations from $N(0,1)$. We then have

$$(4.11) \quad \mu_1'(n, k) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} P_2(x) e^{-x^2} dx$$

$$(4.12) \quad \mu_2'(n, k) = 1 + \frac{1}{2!(2\pi)^{3/2}} \int_{-\infty}^{+\infty} P_3(x) e^{-3x^2/2} dx$$

$$(4.13) \quad \mu_3'(n, k) = \frac{5}{2} \mu_1'(n, k) + \frac{1}{3!(2\pi)^2} \int_{-\infty}^{+\infty} P_4(x) e^{-4x^2/2} dx$$

$$(4.14) \quad \mu_4'(n, k) = -\frac{4}{3} + \frac{13}{3} \mu_2'(n, k) + \frac{1}{4!(2\pi)^{5/2}} \int_{-\infty}^{+\infty} P_5(x) e^{-5x^2/2} dx$$

In general applying the Lemma to (3.17) we can express $\mu_t'(n, k)$ in terms of lower moments of even (odd) order when t is even (odd) and the integral

$$(4.15) \quad \int_{-\infty}^{+\infty} P_{t+1}(x) e^{-(t+1)x^2/2} dx$$

where the polynomials $P_2(x) \dots P_5(x)$ are given by (2.8) ... (2.11). In the particular case when $n = k$, i.e., $x_{(n)}$ is the largest of $x_1, x_2, \dots, x_n$, $P_t(x)$ assumes the very simple form

$$(4.16) \quad P_t(x) = n(n-1)(n-2) \dots (n-t+1) [\Phi(x)]^{n-t}.$$

Hence we get

$$(4.17) \quad \mu_1'(n, n) = \frac{2 \binom{n}{2}}{2\pi} \int_{-\infty}^{+\infty} [\Phi(x)]^{n-2} e^{-x^2} dx$$

$$(4.18) \quad \mu_2'(n, n) = 1 + \frac{3 \binom{n}{3}}{(2\pi)^{3/2}} \int_{-\infty}^{+\infty} [\Phi(x)]^{n-3} e^{-3x^2/2} dx$$

$$(4.19) \quad \mu_3'(n,n) = \frac{5}{2} \mu_1'(n,n) + \frac{4 \binom{n}{4}}{(2\pi)^2} \int_{-\infty}^{+\infty} [\bar{\Phi}(x)]^{n-4} e^{-4x^2/2} dx$$

$$(4.20) \quad \mu_4'(n,n) = -\frac{4}{3} + \frac{13}{3} \mu_2'(n,n) + \frac{5 \binom{n}{5}}{(2\pi)^{5/2}} \int_{-\infty}^{+\infty} [\bar{\Phi}(x)]^{n-5} e^{-5x^2/2} dx.$$

It should be noted that in the formulae (4.17) through (4.20)

$$[\bar{\Phi}(x)]^{n-t}$$

should be interpreted as zero if $t > n$.

Some integrals of the type occurring in (4.17) through (4.20) have been numerically evaluated by Hojo [10].

5. Exact values of some moments.

Let

$$(5.1) \quad I_n(a) = \int_{-\infty}^{+\infty} [\bar{\Phi}(ax)]^n e^{-x^2} dx,$$

then

$$(5.2) \quad I_0(a) = \pi^{1/2}.$$

Now

$$(5.3) \quad \int_{-\infty}^{+\infty} [\bar{\Phi}(ax) - \frac{1}{2}]^{2m+1} e^{-x^2} dx = 0,$$

since the integrand is an odd function of x . Hence

$$(5.4) \quad I_{2m+1}(a) = \sum_{r=1}^{2m+1} (-1)^{r+1} \binom{2m+1}{r} I_{2m-r+1}(2) / 2^r.$$

In particular

$$(5.5) \quad I_1(a) = \frac{1}{2} I_0(a) = \frac{1}{2} \pi^{1/2},$$

and

$$(5.6) \quad I_3(a) = \frac{3}{2} I_2(a) - \frac{3}{4} I_1(a) + \frac{1}{8} I_0(a) = \frac{3}{2} I_2(a) - \frac{1}{4} I_0(a).$$

In general, $I_{2m+1}(a)$ can be expressed as a linear function of $I_{2m}(a)$, $I_{2m-2}(a)$, ..., $I_0(a)$.

Differentiating (5.1) with respect to a , (this is justified in virtue of the uniform convergence of the integrals with respect to a , $-\infty < a < \infty$, and the continuity of the integrands), we get for $n = 2$,

$$(5.7) \quad \frac{dI_2}{da} = \frac{1}{\pi^{1/2}} \frac{a}{(a^2+2)(a^2+1)^{1/2}},$$

so that

$$(5.8) \quad I_2(a) = \frac{1}{\pi^{1/2}} \arctan(\sqrt{a^2+1}).$$

Using (5.6)

$$(5.9) \quad I_3(a) = \frac{3}{2\pi^{1/2}} \arctan(\sqrt{a^2+1}) - \frac{\pi^{1/2}}{4}.$$

Since $P_{t+1}(x)$ is a polynomial in $\Phi(x)$ of degree $n-t-1$, by using (5.2), (5.5), (5.8) and (5.9), we can exactly evaluate (4.15) if $n \leq t+4$. Hence we can exactly evaluate $\mu_t^i(n,k)$ for all odd values of t , if $n \leq 5$ and all even values of t if $n \leq 6$. Godwin [6] and Jones [11] have given tables of exact moments μ_t^i for $t = 1$ and 2 . The corresponding tables for $t = 3$ and 4 are given below.

Table I.

$\mu_3^i(n,k)$			
$n \quad k$	$k = n$	$k = n - 1$	$k = n - 2$
2	$2A$		
3	$3A$	0	
4	$2B_1 + 2C$	$12A - 6B_1 - 6C$	
5	$-5A + 5B_1 + 5B_2$	$20A - 10B_1 - 10C$	0

Here

$$A = \frac{5}{4\pi^{1/2}}$$

$$B_1 = \frac{1}{2\sqrt{2}\pi^{3/2}}$$

$$B_2 = \frac{15}{2\pi^{3/2}}$$

$$C = \frac{15}{2\pi^{3/2}} \arctan(\sqrt{2})$$

Table II

Here

$\mu_k^1(n, k)$			
k			
n			
2	3		
3	3+a	3-2a	
4	3+2a	3-2a	
5	3+b+c	3+10a-4b-4c	3-20a+6b+6c
6	3-5a+3b+3c	3+25a-9b-9c	3-20a+6b+6c

$$a = \frac{13}{\sqrt{3}(2\pi)}$$

$$b = \frac{65}{\sqrt{3}\pi^2} \arctan(\sqrt{5}/3)$$

$$c = \frac{\sqrt{5}}{4\pi^2}$$

REFERENCES

- [1] Bahadur, Raghu Raj, "On a problem in the theory of k populations," Ann. Math. Stat., Vol. 21 (1950), pp. 362-375.
- [2] Bechhofer, Robert E., "A single-sample multiple decision procedure for ranking means of normal populations with known variances," Ann. Math. Stat., Vol. 25 (1954), pp. 16-29.
- [3] Bechhofer, R. E., Dunnett, C. W. and Sobel, M., "A two-sample multiple decision procedure for ranking means of normal populations with a common unknown variance," Biometrika, Vol. 41 (1954), pp. 170-176.
- [4] Bechhofer, R. E. and Sobel, M., "A sequential multiple decision procedure for selecting the population with the largest mean from k normal populations with a common unknown variance (preliminary Report)," abstract, Ann. Math. Stat., Vol. 27 (1956), pp. 218-219.
- [5] Cole, R. H., "Relations between moments of order statistics," Ann. Math. Stat. Vol. 22 (1951), pp. 308-310.
- [6] Godwin, H. J., "Some low moments of order statistics," Ann. Math. Stat. Vol. 20 (1949), pp. 279-285.

- [7] Godwin, H. J., "On the estimation of dispersion by linear systematic statistics," Biometrika, Vol. 36 (1949), pp. 92-100.
- [8] Gupta, Shanti S., "On a decision rule for a problem in ranking means," Ph. D. Thesis (1956), University of North Carolina, Chapel Hill.
- [9] Hastings, C., Jr.; Mosteller, F.; Tukey, J. W. and Winsor, C. P., "Low moments for small samples: a comparative study of order statistics," Ann. Math. Stat., Vol. 18 (1947), pp. 413-26.
- [10] Hojo, T., "Distribution of median, quartiles, and interquartile distance in samples from a normal population," Biometrika, Vol. 23 (1931), pp. 315-360.
- [11] Jones, H. L., "Exact lower moments of order statistics in small samples from a normal distribution," Ann. Math. Stat., Vol. 19 (1948), pp. 270-273.
- [12] Mosteller, Frederick, "On some useful inefficient statistics," Ann. Math. Stat., Vol. 17 (1946), pp. 377-408.
- [13] Mosteller, Frederick, "A k-sample slippage test for an extreme population," Ann. Math. Stat., Vol. 19 (1948), pp. 58-65.
- [14] Nair, K. R., "The distribution of the extreme deviate from the sample mean and its studentized form," Biometrika, Vol. 35 (1948), pp. 118-144.
- [15] Paulson, Edward, "A multiple decision procedure for certain problems in the analysis of variance," Ann. Math. Stat., Vol. 20 (1949), pp. 95-98.
- [16] Paulson, Edward, "On the comparison of several experimental categories with a control," Ann. Math. Stat., Vol. 23 (1952), pp. 239-246.
- [17] Paulson, Edward, "An optimum solution to the k-sample slippage problem for the normal distribution," Ann. Math. Stat., Vol. 23 (1952) pp. 610, 616.

- [18] Ruben, H., "On the moments of order statistics in samples from normal populations," Biometrika, Vol. 41 (1954), pp. 200-227.
- [19] Salzer, H. E.; Zucker, Ruth; and Capuano, Ruth, "Table of the zeros and weight factors of the first twenty Hermite polynomials," Journal of Research of the National Bureau of Standards, Vol. 48 (1952), pp. 111-116.
- [20] Seal, K. C., "On a class of decision procedures for ranking means of normal populations," Ann. Math. Stat., Vol. 26 (1955), pp. 387-398.
- [21] Szego, Gabor, Orthogonal Polynomials, New York City, American Mathematical Society Colloquium Publications, Vol. 23 (1939).
- [22] Teichroew, Daniel, Probabilities Associated with Order Statistics in Samples from Two Normal Populations with Equal Variance, Chemical Corps Engineering Agency Statistical Tables, Army Chemical Center, Maryland, 1955.
- [23] Tippett, L. H. C., "The extreme individuals and the range of samples taken from a normal population," Biometrika, Vol. 17 (1925), pp. 364-387.