UNIVERSITY OF NORTH CAROLINA
Department of Statistics
Chapel Hill, N. C.

Mathematical Sciences Directorate
Air Force Office of Scientific Research
Washington 25, D. C.

AFOSR Report No.

AN ALGORITHM FOR A MINIMUM COVER
OF
AN ABSTRACT COMPLEX *

by

D. K. Ray-Chaudhuri

February, 1961

Contract No. AF 49(638)-213

Algorithms for a minimum cover and maximum matching of an abstract complex are obtained. These algorithms are useful for many combinatorial problems.

* This research was supported in part by the United States Air Force Office of Scientific Research of the Air Research and Development Command, under Contract No. AF(638)-213. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Qualified requestors may obtain copies of this report from the ASTIA Document Service Center, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their "need-to-know" certified by the cognizant military agency of their project or contract.

Mimeo Series 275
Institute of Statistics

AN ALGORITHM FOR A MINIMUM COVER OF AN ABSTRACT COMPLEX

D. K. RAY - CHAUDHURI
University of North Carolina

Introduction. Let $X = \{x_1, x_2, \dots, x_m\}$ be a finite set of m points and $\mathcal{A} = \{A_1, A_2, \dots, A_n\}$ be a class of n subsets of X . Such a system of points and sets is called a complex $(X, \mathcal{A})$. If every set of the class $\mathcal{A}$ contains two points, the complex is a graph with m points $x_1, x_2, \dots, x_m$ and n edges $A_1, A_2, \dots, A_n$. For any subclass $\mathcal{A}_1$ of $\mathcal{A}$, $\mathcal{A}_1(x)$ denotes the class consisting of the sets which belong to $\mathcal{A}_1$ and contain the point x . For a subset A , $\mathcal{A}_1(A)$ denotes the class consisting of those sets which belong to $\mathcal{A}_1$ and contain at least one point of A . A complex $(X, \mathcal{A})$ in which every set has the same number of points is called a regular complex. $|A|$ and $|\mathcal{A}_1|$ denote respectively the number of points in A and the number of sets in the class $\mathcal{A}_1$. Let $\underline{c}$ be a m -vector of positive integers $c(x_1), c(x_2), \dots, c(x_m)$. A subclass $\mathcal{A}_1$ is called a $\underline{c}$ -cover if for every point x , $|\mathcal{A}_1(x)| \geq c(x)$. A subclass $\mathcal{A}_1$ is a $\underline{c}$ -matching if for every point x , $|\mathcal{A}_1(x)| \leq c(x)$. Covers with minimum number of sets and matchings with maximum number of sets are respectively called minimum covers and maximum matchings. For a minimum $\underline{c}$ -cover $\mathcal{A}_1$, let X_1 be the set of those point x for which $|\mathcal{A}_1(x)| = c(x)$. Following Fulkerson and Ryser $\lfloor \frac{1}{2} \rfloor |X_1|$ is called the $\underline{c}$ -height of $\mathcal{A}_1$. Minimum possible $\underline{c}$ -height of a minimum $\underline{c}$ -cover is called the $\underline{c}$ -height of the complex $(X, \mathcal{A})$. Similarly for a $\underline{c}$ -matching $\mathcal{A}_1$, let X_2 denote the set of those points x for which $|\mathcal{A}_1(x)| < c(x)$. $|X_2|$

is called the $\underline{c}$ - depth of the $\underline{c}$ - matching A_1 . Maximum possible $\underline{c}$ - depth of a $\underline{c}$ - matching is called the $\underline{c}$ - depth of the complex.

To a complex (X, A) we can associate an incidence matrix $A = ((a_{ij}))$ with m rows and n columns where $a_{ij} = 1$ if $x_i \in A_j$, and $a_{ij} = 0$, otherwise, $i = 1, 2, \dots, m$, $j = 1, 2, \dots, n$. Fulkerson and Ryser [2] considers the minimum cover problem in terms of the matrix A . A set of ε columns of the matrix A is called an α -set of representatives if in the submatrix of A consisting of the m rows and the ε columns, each row sum is not less than α . Let $\varepsilon(\alpha)$ be the minimum number of columns of A that form an α -set of representatives. The number $\varepsilon(\alpha)$ is called the α -width of the matrix A . Obviously $\varepsilon(\alpha)$ is the cardinality of the minimum $\underline{c}$ - cover of the complex (X, A) where $\underline{c} = (\alpha, \alpha, \dots, \alpha)$. Fulkerson and Ryser comments, "Very little is known concerning good computational methods for determining widths and heights of $(0, 1)$ -matrices. Efficient algorithms in this domain would be of great interest".

Petersen [4] introduced the concept of alternating path which led to algorithms for minimum cover and maximum matching of graphs. Berge [1] first gave an inductive proof of the maximum matching theorem for graphs. Norman and Rabin [3] gave an ingenious proof of the maximum matching and minimum cover theorems for graphs. In this paper the idea of alternating path is extended to general complexes and certain theorems are proved about minimum covers and maximum matchings of general complexes. The proofs of this theorem follow the line of proof of Norman and Rabin [3]. Section 1 proves the main theorem about the minimum cover of a complex and gives the corresponding algorithm. A 'level transformation' is defined. By applying this transformation on a particular minimum cover all minimum covers can be obtained. In section 3 we prove a theorem about the minimum cover of a regular complex. This theorem gives a shorter algorithm for the minimum cover of a regular

complex. In section 3 is proved a theorem about the height of a minimum cover which gives an algorithm for the height of a complex. A 'parallel transformation' is defined. By applying ^{this transformation} on a minimum cover with minimum height, the family of all minimum covers with minimum heights ^{can be obtained.} Section 4 gives a simpler algorithm for the height of a regular complex. Section 5 establishes a close relationship between a matching problem and a cover problem. It is shown that a maximum $\underline{c}$ - matching can be obtained from a certain minimum cover. Also the $\underline{c}$ - depth of a complex can be determined from the $\underline{d}$ - height of the complex for a certain vector $\underline{d}$.

Extending the ideas of the present paper, it is possible to get an algorithm for any integer programming problem. This will be developed in a subsequent paper. There are many important practical applications of the minimum cover and maximum matching algorithm. For some special applications it is possible to obtain an algorithm sharper than the general algorithm. These applications also will be discussed in the subsequent paper.

1. Minimum $\underline{c}$ - cover. Let A_1 and B_1 be two classes of sets. Consider a finite sequence C of distinct sets $A_1, B_1, A_2, B_2, \dots$ where $A_i \in A_1$ and $B_i \in B_1$, $i = 1, 2, \dots$. Let H_i be the union of the sets $A_1, B_2, \dots, A_i, B_i$, $i = 1, 2, \dots$. D_i and F_i respectively denote the classes consisting of the sets $A_1, A_2, \dots, A_i$ and $B_1, B_2, \dots, B_i$, $i = 1, 2, \dots$. F_0 denotes the null class.

$$\text{Let } D_i = \left\{ x \mid x \in H_i, |A_1(x)| + |F_{i-1}(x)| - |D_i(x)| < c(x) \right\}$$

$$\text{and } F_i = \left\{ x \mid x \in G_i, |D_i(x)| < |F_i(x)| \right\}, i = 1, 2, \dots$$

We shall use these notations throughout sections 1 and 2. Let A_1 be a $\underline{c}$ - cover and $B_1 = A - A_1$. A sequence $C = \{A_1, B_1, A_2, B_2, \dots\}$

is an alternating chain if

$$(1.1) \quad \begin{aligned} & A_1 \in \mathcal{A}_1 \\ & \dots B_i \in (\mathcal{B}_1 - \mathcal{F}_{i-1})(D_i) \\ & \text{and } A_{i+1} \in (\mathcal{A}_1 - \mathcal{D}_i)(F_i), \end{aligned}$$

$$i = 1, 2, \dots$$

An alternating chain $C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}$ is called a reducible chain if D_{p+1} is a null set. A single set A_1 is a reducible chain if D_1 is a null set, i.e. if for every point x of A_1 , $|A_1(x)| > c(x)$. The set A_1 of the reducible chain is called the leader of the chain. A $\underline{c}$ -cover A_1 is an irreducible $\underline{c}$ -cover if there exists no reducible chain with respect to (w.r.t) A_1 .

It is easy to see that if A_1 is a $\underline{c}$ -cover and $C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}$ is a reducible chain w.r.t A_1 , then $A_2 = A_1 - \mathcal{D}_{p+1} \cup \mathcal{F}_p$ is a $\underline{c}$ -cover with cardinality less than that of A_1 .

Let $\mathcal{B}_1$ be a $\underline{c}$ -cover and $A_1 = A - \mathcal{B}_1$. Consider a sequence C of sets $A_1, B_1, A_2, B_2, \dots, A_p, B_p$ where $A_i \in \mathcal{A}_1$ and $B_i \in \mathcal{B}_1$, $i = 1, 2, \dots, p$.

$$\text{Let } D_i' = \left\{ x \mid x \in H_i, |\mathcal{F}_{i-1}(x)| < |\mathcal{D}_i(x)| \right\}, \\ i = 1, 2, \dots$$

Such a sequence is called a level chain if

$$\begin{aligned} B_i & \in (\mathcal{B}_1 - \mathcal{F}_{i-1})(D_i'), \\ A_{i+1} & \in (\mathcal{A}_1 - \mathcal{D}_i)(F_i) \end{aligned}$$

$$\text{and } |\mathcal{B}_1(x)| + |\mathcal{D}_p(x)| - |\mathcal{F}_p(x)| \geq c(x)$$

for every x in H_p and $i = 1, 2, \dots$. Consider the transformation which takes the $\underline{c}$ -cover $\mathcal{B}_1$ to $\mathcal{B}_2$ where

$$\mathcal{B}_2 = \mathcal{B}_1 - \mathcal{F}_p \cup \mathcal{D}_p$$

Obviously $\mathcal{B}_2$ is also a $\underline{c}$ -cover. Since the cardinality of $\mathcal{B}_2$ is same as that of $\mathcal{B}_1$, the above transformation is called a 'level transformation'.

Let $L(\mathcal{B}_1)$ be the family of all classes which can be obtained from $\mathcal{B}_1$ by applying the level transformation a finite number of times. For two covers

A_1 and A_0 , $d(A_1, A_0)$ denotes the number of sets of the class

A_1 which do not belong to A_0 . Let $\mathcal{B}_1$ be a cover in the family $L(A_0)$ such that

$$d(A_1, \mathcal{B}_1) \leq d(A_1, A'), A' \in L(A_0).$$

Then $\mathcal{B}_1$ is said to be a cover of the family $L(A_0)$ nearest to A_1 .

Lemma 1. Let A_1 and A_0 be two $\underline{c}$ -covers. Let $\mathcal{B}_1$ be a cover in the family $L(A_0)$ nearest to A_1 . If there exists a set A_1 in $A_1 - \mathcal{B}_1$, then there exists a reducible chain w.r.t. A_1

$$C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}$$

where $A_{i+1} \in A_1 - \mathcal{B}_1$ and $B_i \in \mathcal{B}_1 - A_1$, $i = 1, 2, \dots, p$.

Proof. Assume that there exists no such reducible chain (w.r.t. A_1) with A_1 as the leader. Consider $(\mathcal{B}_1 - \mathcal{F}_0)(D_1)$. Since there is no reducible chain with A_1 as leader D_1 is not empty. If possible, suppose

$$(1.1) \dots \dots \dots (\mathcal{B}_1 - \mathcal{F}_0)(D_1) \subset A_1 - D_1.$$

Since $\mathcal{B}_1$ is a $\underline{c}$ -cover, it follows from (1.1) that for every x in D_1

$$(1.2) \dots \dots \dots |A_1(x)| - |D_1(x)| + |\mathcal{F}_0(x)| \geq c(x).$$

This contradicts the definition of D_1 . Hence (1.1) is not true and we can choose B_1 where

$$B_1 \in (B_1 - F_0)(D_1) - (A_1 - D_1).$$

Next we consider $(A_1 - D_1)(F_1)$. If possible, suppose

$$(1.3) \dots\dots\dots (A_1 - D_1)(F_1) \subset B_1 - F_1$$

Since A_1 is a $\underline{c}$ -cover, from (1.3) it follows that for every x in F_1

$$(1.4) \dots\dots\dots |B_1(x)| - |F_1(x)| + |D_1(x)| \geq c(x).$$

From (1.4) we can easily see that $C = \{A_1, B_1\}$ is a level chain w.r.t. B_1 and

$$B_2 = B_1 - F_1 \cup D_1 \in L(A_0).$$

Also we have

$$(1.5) \dots\dots\dots d(A_1, B_2) < d(A_1, B_1)$$

which contradicts the assumption that B_1 is nearest to A_1 . Hence

(1.3) is not true and we can choose A_2 where

$$A_2 \in (A_1 - D_1)(F_1) - (B_1 - F_1)$$

Using the same arguments by induction we can show that there exists an infinite sequence of distinct sets which is a contradiction. Hence there must exist a reducible chain with A_1 as the leader with the required property.

Theorem 1. A $\underline{c}$ -cover A_1 of a complex (X, A) is a minimum $\underline{c}$ -cover if and only if A_1 is irreducible.

Proof. Necessity is obvious. To prove sufficiency, assume that A_1 is irreducible. Let A_0 be a minimum $\underline{c}$ -cover. Let B_1 be a cover in the family $L(A_0)$ nearest to A_1 . Since B_1 is also a minimum $\underline{c}$ -cover, it is sufficient to show that $A_1 \subset B_1$. If possible, suppose there exists a set A_1 contained in $A_1 - B_1$. Then by lemma 1, there is a reducible chain w.r.t. A_1 which is a contradiction. Hence $A_1 - B_1$ is empty. This completes the proof of the theorem.

Theorem 2. If A_1 is a minimum $\underline{c}$ - cover, then any other minimum $\underline{c}$ - cover A_2 belongs to the family $L(A_1)$.

Proof. Let B_1 be the cover in the family $L(A_1)$ nearest to A_2 . Since A_2 is a minimum $\underline{c}$ - cover, it is irreducible. Now by the same arguments as used in the proof of theorem 1, we can show that $B_1 = A_2$.

Theorem 1 gives the following algorithm for the minimum cover of a complex. We start with a $\underline{c}$ - cover A_1 . If there is no reducible chain w.r.t. A_1 , A_1 is a minimum $\underline{c}$ - cover. So the algorithm consists in looking for reducible chains. Whenever a reducible chain is obtained, we get a new $\underline{c}$ - cover whose cardinality is one less than that of the original $\underline{c}$ - cover. In this way, finally we get a $\underline{c}$ - cover which is irreducible. To test if there is a reducible chain w.r.t. a $\underline{c}$ - cover A_1 with a given set A_1 of A_1 as the leader, we can proceed as follows. Let $B_1 = A - A_1$. We start with A_1 and test if A_1 is a reducible chain w.r.t. A_1 . If not, if possible we choose $B_1 \in \mathcal{B}_1(D_1)$. Next if possible we find

$$A_2 \in (A_1 - D_1) (F_1)$$

and test whether D_2 is a null set. If D_2 is a null set, $D_2 \cup F_1$ is a reducible chain w.r.t. A_1 and the test is completed. If not, if possible we find

$$B_2 \in (B_1 - F_1) (D_2).$$

In this manner, we proceed to build a chain until we get a reducible chain or until no further addition of sets to the chain satisfying the conditions (1.1)

is possible. For instance the chain will terminate after the selection of $A_1, B_1, \dots, A_p, B_p$ if $(A_1 - D_p) (F_p)$ is a null class. The chain will terminate after the selection of $A_1, B_1, \dots, A_p, B_p, A_{p+1}$ if $D_{p+1} \cup F_p$ is a reducible chain w.r.t. A_1 or $(B_1 - F_p) (D_{p+1})$ is a null class. At the terminating stage, we have a maximal chain. By varying the choices of the sets $B_1, A_2 \dots$ we can get all possible maximal chains and

find out if there is a reducible chain w.r.t. A_1 with A_1 as the leader. Theorem 2 gives an algorithm for finding out all minimum $\underline{c}$ - covers starting from a given minimum $\underline{c}$ - cover A_1 . The algorithm consists in looking for level chains. First we find out all possible level chains w.r.t. A_1 and find out minimum $\underline{c}$ - covers $A_2, A_3, \dots, A_k$. Next we find level chains w.r.t. each of $A_2, A_3, \dots, A_k$ and find new minimum $\underline{c}$ - covers $A_{k+1}, \dots, A_l$ and so on. Finally we shall reach a stage when we do not find any more minimum $\underline{c}$ - covers.

2. Minimum covers of regular complexes. Let A_1 be a $\underline{c}$ - cover of a regular complex (X, A) . A set A of A_1 is called a heavy set if for at least one point x of A , $|A_1(x)| > c(x)$. A reducible chain whose leader is a heavy chain is called a heavy reducible chain. In this section we consider only regular complexes.

Lemma 2. A reducible chain w.r.t. a $\underline{c}$ - cover A_1 contains at least one heavy set.

Proof. Suppose $C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}$ is a reducible chain w.r.t. A_1 . Let H_{p+1} be the union of the sets in the chain C . Let H' be the union of the sets $A_1, A_2, \dots, A_{p+1}$ and $H^* = H_{p+1} - H'$. If possible, suppose there is no point x in H' for which $|A_1(x)| > c(x)$. Let q be the number of points in every set of A . Then it follows easily that

$$(2.1) \dots \sum_{x \in H_{p+1}} |D_{p+1}(x)| = (p+1)q > pq = \sum_{x \in H_{p+1}} |F_p(x)|.$$

Let $A' = A_1 - D_{p+1}$. Since C is a reducible chain $A_2 = A' \cup F_p$ is a $\underline{c}$ - cover. For every point x in H' ,

$$|A_1(x)| = |A'(x)| + |D_{p+1}(x)| = c(x). \text{ Also } |A_2(x)| = |A'(x)| + |F_p(x)|$$

Hence for x in H'

$$(2.2) \dots \dots \dots |A'(x)| + |D_{p+1}(x)| - c(x) \leq |A'(x)| + |F_p(x)| - c(x),$$

For x in H^* , $|D_{p+1}(x)| = 0$. Hence for $x \in H^*$,

$$(2.3) \dots \dots \dots |A'(x)| + |D_{p+1}(x)| - c(x) \leq |A'(x)| + |F_p(x)| - c(x).$$

From (2.2) and (2.3), we get

$$(2.4) \dots \dots \dots \sum_{x \in H_{p+1}} (|A'(x)| + |D_{p+1}(x)| - c(x)) \leq$$

$$\sum_{x \in H_{p+1}} (|A'(x)| + |F_p(x)| - c(x))$$

The inequality (2.4) contradicts (2.2). Hence there must exist a point x in H' for which $|A_1(x)| > c(x)$. In other words there is a heavy set in the reducible chain.

Lemma 3. Let A_1 be any $\underline{c}$ -cover and A_0 be a minimum $\underline{c}$ -cover. Let

B_1 be a $\underline{c}$ -cover in the family $L(A_0)$ nearest to A_1 . If all heavy sets of A_1 are contained in B_1 , then A_1 is a minimum - cover.

Proof. It is sufficient to show that $A_1 \subset B_1$. If not, suppose $A_1 -$

B_1 contains a set A_1 . Then by lemma 1, there exists a reducible chain $C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}$ where $A_{i+1} \in A_1 - B_1$ and $B_i \in B_1 - A_1$, $i = 1, 2, \dots, p$. By lemma 2 there is a heavy set A in the reducible chain C . This contradicts our assumption. Hence the lemma follows.

Theorem 3. If a $\underline{c}$ -cover A_1 is such that there is no heavy reducible chain, w.r.t. A_1 , then A_1 is a minimum $\underline{c}$ -cover.

Proof. Let A_0 be a minimum $\underline{c}$ -cover and B_1 be a $\underline{c}$ -cover in the family $L(A_0)$ nearest to A_1 . By lemma 3, it is sufficient to show that all heavy sets of A_1 are contained in B_1 . If not, suppose $A_1 - B_1$ contains a heavy set A_1 . Then by lemma 1 there exists a reducible chain w.r.t. A_1 .

$$C = \{A_1, B_1, \dots, A_p, B_p, A_{p+1}\}.$$

By definition C is a heavy reducible chain. This contradicts our hypothesis. Hence the theorem is true.

Theorem 3 gives a shorter algorithm for minimum covers of regular complexes. It shows that in the case of regular complexes it is necessary to look for only heavy reducible chains.

3. Height of a complex. Let A_1 and B_1 be two subclasses of A . Consider a sequence E of sets $E_1, E_2, \dots, E_{2p-1}, E_{2p}$ belonging to A . Let $P_i = \bigcup_{j=1}^i E_j$, $i=1, 2, \dots, 2p$. Let R_i and S_i respectively denote the classes consisting of those sets of A_1 and B_1 which occur among $E_1, E_2, \dots, E_i$, $i = 1, 2, \dots, 2p$.

Let

$$R_i = \{x \mid x \in P_i, |R_i(x)| > |S_i(x)|\}$$

(3.1)

$$\text{and } S_i = \{x \mid x \in P_i, |R_i(x)| < |S_i(x)|\}$$

$$i = 1, 2, \dots, 2p.$$

We shall use the above notations throughout sections 3 and 4.

Now we assume that A_1 is a $\underline{c}$ -cover and $B_1 = A - A_1$.

A sequence E is called a properly connected sequence (w.r.t. A_1)

if the following four conditions are satisfied for $k = 1, 2, \dots, p$.

(i) If $E_{2k} \in B_1$ and $(A_1 - R_{2k})(S_{2k})$ is not empty,

$$E_{2k+1} \in (A_1 - R_{2k})(S_{2k})$$

$$\text{and } E_{2k+2} \in (B_1 - S_{2k+1})(R_{2k+1}).$$

(ii) If $E_{2k} \in B_1$ and $(A_1 - R_{2k}) (S_{2k})$ is empty,

$$E_{2k+1} \in (B_1 - I_{2k}) (R_{2k})$$

$$\text{and } E_{2k+2} \in (A_1 - R_{2k+1}) (S_{2k+1})$$

(iii) If $E_{2k} \in A_1$ and $(B_1 - I_{2k}) (R_{2k})$ is not empty,

$$E_{2k+1} \in (B_1 - I_{2k}) (R_{2k})$$

$$\text{and } E_{2k+2} \in (A_1 - R_{2k+1}) (S_{2k+1}).$$

(iv) If $E_{2k} \in A_1$ and $(B_1 - I_{2k}) (R_{2k})$ is empty,

$$E_{2k+1} \in (A_1 - R_{2k}) (S_{2k})$$

$$\text{and } E_{2k+2} \in (B_1 - I_{2k+1}) (R_{2k+1}).$$

A properly connected sequence E is called an exchange sequence if

$A_2 = A_1 - R_{2p} \cup I_{2p}$ is a $\underline{c}$ -cover. An exchange sequence E (w.r.t. A_1) is called a low sequence (w.r.t. A_1) if

$E_1 \in A_1$ and $E_2 \in B_1(R_1)$ and

$$(3.2) \dots \dots \dots \left| \left\{ x \mid \begin{array}{l} |A_1(x)| = c(x) \\ |A_2(x)| = c(x) \end{array} \right\} \right| > \left| \{ x \mid \right|$$

An exchange sequence E (w.r.t. A_1) is called a parallel sequence

(w.r.t. A_1) if $E_1 = B_1$ and $E_2 \in A_1(S_1)$ and

$$(3.3) \dots \dots \dots \left| \left\{ x \mid \begin{array}{l} |A_1(x)| = c(x) \\ |A_2(x)| = c(x) \end{array} \right\} \right| = \left| \{ x \mid \right|$$

Obviously if A_1 is a $\underline{c}$ -cover and E is a low sequence w.r.t.

A_1 , A_2 is a $\underline{c}$ -cover with height less than that of A_1 .

Similarly if E is a parallel sequence, A_2 is a $\underline{c}$ -cover with

height equal to that of A_1 . In this case for convenience we say that

A_2 is obtained from A_1 by applying a 'parallel transformation'.

Let $M(A_1)$ denote the family of all $\underline{c}$ -covers which can be obtained from A_1 by applying parallel transformation a finite number of times.

Lemma 4. Suppose A_1 and B_1 are two minimum $\underline{c}$ -covers, B_1 being a minimum $\underline{c}$ -cover with minimum height. Let $E = \{E_1, E_2, \dots, E_{2p}\}$ be a properly connected sequence w.r.t. both A_1 and B_1 where $E_1 \in$

$A_1 - B_1$ and $E_2 \in B_1 - A_1$ and each E_i belongs to either $A_1 - B_1$ or $B_1 - A_1$. If

$$(A_1 - R_{2p})^{(S_{2p})} \subset (B_1 - I_{2p})^{(S_{2p})}$$

(3.4)

$$\text{and } (B_1 - I_{2p})^{(R_{2p})} \subset (A_1 - R_{2p})^{(R_{2p})},$$

then either E is a low sequence w.r.t. A_1 or E is a parallel sequence w.r.t. B_1 .

Proof. First we show that E is an exchange sequence w.r.t. both A_1 and B_1 . For this we have to show that both $A_2 = A_1 - R_{2p} \cup$

I_{2p} and $B_2 = B_1 - I_{2p} \cup R_{2p}$ are $\underline{c}$ -covers. To show that A_2 is a $\underline{c}$ -cover, it is sufficient to show that for every $x \in R_{2p}$, $|A_2(x)| \geq c(x)$. Using (3.4) and the fact that A_1 is a $\underline{c}$ -cover, we get for $x \in R_{2p}$

$$\begin{aligned} |A_2(x)| &= |A_2(x)| = |(A_1 - R_{2p})(x)| + |I_{2p}(x)| \\ &\geq |(B_1 - I_{2p})(x)| + |I_{2p}(x)| \\ &= |B_1(x)| \geq c(x). \end{aligned}$$

Similarly we can show that B_2 is a $\underline{c}$ -cover. From (3.4), we can easily obtain

$$(3.5) \dots \dots \dots \left\{ x \mid x \in R_{2p}, |A_2(x)| = c(x) \right\} \subset \left\{ x \mid x \in R_{2p}, |A_1(x)| = c(x) \right\}$$

$$\text{and } \left\{ x \mid x \in S_{2p}, |B_2(x)| = c(x) \right\} \subset \left\{ x \mid x \in S_{2p}, |A_1(x)| = c(x) \right\}.$$

If E is not a low sequence w.r.t. A_1 , we have

$$(3.6) \dots \dots \dots \left| \left\{ x \mid |A_1(x)| > c(x) \right\} \right| \geq \left| \left\{ x \mid |A_2(x)| > c(x) \right\} \right|$$

If x does not belong to $R_{2p} \cup S_{2p}$, $|A_1(x)| = |A_2(x)|$.

Hence from 3.6, it follows that

$$(3.7) \dots \dots \dots \left| \left\{ x \mid x \in R_{2p}, |A_1(x)| > c(x) \right\} \right| - \left| \left\{ x \mid x \in R_{2p}, |A_2(x)| > c(x) \right\} \right| \geq \left| \left\{ x \mid x \in S_{2p}, |A_2(x)| > c(x) \right\} \right| - \left| \left\{ x \mid x \in S_{2p}, |A_1(x)| > c(x) \right\} \right|$$

It is easy to check that

$$\left\{ x \mid x \in R_{2p}, |A_2(x)| = c(x) \right\} = \left\{ x \mid x \in R_{2p}, |A_1(x)| > c(x) \right\} - \left\{ x \mid x \in R_{2p}, |A_2(x)| > c(x) \right\}$$

$$(3.8) \dots \dots \dots \text{and } \left\{ x \mid x \in S_{2p}, |A_1(x)| = c(x) \right\} = \left\{ x \mid x \in S_{2p}, |A_2(x)| > c(x) \right\} - \left\{ x \mid x \in S_{2p}, |A_1(x)| > c(x) \right\}$$

Also we have

$$(3.9) \quad \left\{ x \mid x \in R_{2p}, |A_1(x)| > c(x) \right\} \supset \left\{ x \mid x \in R_{2p}, |A_2(x)| > c(x) \right\}$$

and

$$\left\{ x \mid x \in S_{2p}, |A_2(x)| > c(x) \right\} \supset \left\{ x \mid x \in S_{2p}, |A_1(x)| > c(x) \right\}$$

From (3.6), (3.7), (3.8), and (3.9), we get that if E is not a low sequence w.r.t. A_1

$$(3.10) \quad \left| \left\{ x \mid x \in R_{2p}, |A_2(x)| = c(x) \right\} \right| \geq \left| \left\{ x \mid x \in S_{2p}, |A_1(x)| = c(x) \right\} \right|$$

Similarly using the fact that B_1 is a $\underline{c}$ -cover with least height, we can prove that if E is not a level sequence w.r.t. B_1

$$(3.11) \quad \left| \left\{ x \mid x \in S_{2p}, |B_2(x)| = c(x) \right\} \right| > \left| \left\{ x \mid x \in R_{2p}, |B_1(x)| = c(x) \right\} \right|$$

Suppose the lemma is not true, then from (3.10) and (3.11) we have

$$(3.12) \quad \left| \left\{ x \mid x \in R_{2p}, |A_2(x)| = c(x) \right\} \right| + \left| \left\{ x \mid x \in S_{2p}, |B_2(x)| = c(x) \right\} \right| > \left| \left\{ x \mid x \in S_{2p}, |A_1(x)| = c(x) \right\} \right| + \left| \left\{ x \mid x \in R_{2p}, |B_1(x)| = c(x) \right\} \right|$$

which contradicts (3.5). Hence the lemma is true.

Lemma 5. Let A_1 be a minimum $\underline{c}$ -cover and A_0 be a minimum $\underline{c}$ -cover with least height. Let B_1 be a $\underline{c}$ -cover in the family $M(A_0)$ nearest to A_1 . If $A_1 - B_1$ contains a set E_1 , there exists a low sequence $E = \{E_1, E_2, \dots, E_{2p}\}$ w.r.t. A_1 where any set in the sequence E

belonging to A_1 belongs to $A_1 - R_1$.

Proof. If possible, suppose there is no such low sequence w.r.t. A_1 with E_1 as the leader. Consider $R_1(R_1)$. If $R_1(R_1) \subset A_1 - R_1$, we can show that $A_1 - R_1$ is a $\underline{c}$ -cover. This contradicts the assumption that A_1 is a minimum $\underline{c}$ -cover. So we can choose E_2 where

$$E_2 \in R_1(R_1) - (A_1 - R_1)(R_1)$$

Next we consider $(A_1 - R_2)(S_2)$. There are two possible cases.

Case 1. In this case

$$(A_1 - R_2)(S_2) \not\subset (R_1 - I_2)(S_2).$$

So we choose E_3 where

(3.13)

$$E_3 \in (A_1 - R_2)(S_2) - (R_1 - I_2)(S_2).$$

Then consider $(R_1 - I_3)(R_3)$. If possible, suppose

(3.14)

$$(R_1 - I_3)(R_3) \subset (A_1 - R_3)(R_3).$$

In this case we can show that $A_1 - R_3 \cup I_3$ is a $\underline{c}$ -cover

with cardinality less than that of A_1 which is a contradiction. Hence

(3.14) is not true and we can choose

(3.15)

$$E_4 \in (R_1 - I_3)(R_3) - (A_1 - R_3)(R_3)$$

Case 2. In this case

(3.16)

$$(A_1 - R_2)(S_2) \subset (R_1 - I_2)(S_2)$$

Now we consider $(R_1 - I_2)(R_2)$. If possible, suppose

(3.17)

$$(R_1 - I_2)(R_2) \subset (A_1 - R_2)(R_2)$$

Then it is easily checked that the sequence $E = \{E_1, E_2\}$ satisfies the conditions of lemma 4. Hence either E is a low sequence w.r.t. A_1

or E is a parallel sequence w.r.t. B_1 . Since by our assumption there is no low sequence w.r.t. A_1 , E must be a parallel sequence w.r.t. B_1 . Then $B_2 = B_1 - \mathcal{I}_2 \cup \mathcal{R}_2$ belongs to the family $M(A_0)$ and is nearer to A_1 . This contradicts the assumption that B_1 is a $\underline{c}$ -cover in the family $M(A_0)$ nearest to A_1 . Hence (3.17) is not true and we can choose

$$(3.18) \dots \dots \dots E_3 \in (B_1 - \mathcal{I}_2)_{(R_2)} - (A_1 - \mathcal{R}_2)_{(R_2)}$$

Next consider $(A_1 - \mathcal{R}_3)_{(S_3)}$. If possible, suppose

$$(3.19) \dots \dots \dots (A_1 - \mathcal{R}_3)_{(S_3)} \subset (B_1 - \mathcal{I}_3)_{(S_3)}.$$

From (3.19), we can show that $A_2 = A_1 - \mathcal{R}_3 \cup \mathcal{I}_3$

is a $\underline{c}$ -cover with cardinality less than that of A_1 . This contradicts the assumption that A_1 is a minimum $\underline{c}$ -cover. Hence (3.19) is not true and we can choose

$$E_4 \in (A_1 - \mathcal{R}_3)_{(S_3)} - (B_1 - \mathcal{I}_3)_{(S_3)}$$

Using the arguments given above by induction we can show that if there is no low sequence w.r.t. A_1 with E_1 as the leader there exists an infinite sequence of distinct sets which is a contradiction. This completes the proof of the lemma.

Theorem 4. A minimum $\underline{c}$ -cover A_1 has minimum height if and only if there is no low sequence w.r.t. A_1 .

Proof. Necessity is obvious. To prove sufficiency, assume that there is no low sequence w.r.t. the minimum cover A_1 . Let A_0 be a minimum cover with minimum height. Let B_1 be the cover in the family $M(A_0)$ nearest to A_1 . It is sufficient to show that $A_1 \subset B_1$. If possible, suppose $A_1 - B_1$ contains a set E_1 . Then by lemma 5, there is a low sequence w.r.t. A_1 . This contradicts our hypothesis. Hence $A_1 \subset B_1$.

Theorem 5. Let A_0 be a minimum cover with minimum height. Then any other minimum cover A_1 belongs to the family $M(A_0)$.

Proof. Proof follows from lemma 5 and the fact that there is no low sequence w.r.t. A_1 .

Theorem 4 gives an algorithm for the height of a complex. We start with a minimum c -cover A_1 and look for low sequences w.r.t. A_1 . Whenever we get a low sequence, we get a new minimum cover with height less than that of the original minimum cover. In this way finally we get a minimum cover w.r.t. which there are no low sequences. To test if there is any low sequence w.r.t.

A_1 with a set E_1 of A_1 as the leader, we can proceed as follows. If possible, we choose E_2 belonging to $B_1(R_1)$ where $B_1 = A - A_1$. After choosing E_2 , we examine if $E = \{E_1, E_2\}$ is a low sequence. For this is necessary to compute $|A_2(x)| = |A_1(x)| - |R_2(x)| + |S_2(x)|$ for $x \in R_2$. For E to be an exchange sequence, it is necessary that for $x \in R_2$, $|A_2(x)| \geq c(x)$. If E is an exchange sequence, E will be a low sequence w.r.t. A_1 if and only if

$$\left| \left\{ x \mid x \in R_2, |A_2(x)| = c(x) \right\} \right| < \left| \left\{ x \mid x \in S_2, |A_1(x)| = c(x) \right\} \right|.$$

If E is a low sequence, the test terminates. If not, we consider $(A_1 - R_2)(S_2)$. There will be two possible cases (i) $(A_1 - R_2)(S_2)$ is not empty and (ii) $(A_1 - R_2)(S_2)$ is empty. If (i) is the case, we choose E_3 belonging to the class $(A_1 - R_2)(S_2)$. Next, if possible, we choose E_4 from the class $(B_1 - S_3)(R_3)$. If (ii) is the case, if possible we choose E_3 from the class $(B_1 - S_2)(R_2)$. Next if possible, we choose E_4 from the class $(A_1 - R_3)(S_3)$. After

choosing E_4 , we examine if $E = \{E_1, E_2, E_3, E_4\}$ is a low sequence or not. In this manner we proceed to build a sequence $E = \{E_1, E_2, \dots, E_{2p}\}$. The sequence terminates at an even stage if we get a low sequence or if at some stage (say $2p$ th stage) both $(A_1 - R_{2p})$ (S_{2p}) and $(R_{2p} - \mathcal{Y}_{2p})$ (R_{2p}) are empty classes. By varying the choices of the sets $E_2, E_3, E_4, \dots$ we can find out if there is a low sequence with E_2 as the leader or not.

Theorem 5 gives an algorithm for finding out all minimum covers with minimum height.

4. Height of a regular complex. Let A_1 be a minimum $\underline{c}$ -cover of a regular complex (X, A) : A set E_2 of A_1 is called a loaded set if for some point x in E_2 , $|A_1(x)| > c(x) + 1$. A low sequence $E = \{E_1, E_2, \dots, E_{2p}\}$ w.r.t. A_1 whose leader E_1 is a loaded set is called a loaded low sequence. In this section we consider only regular complexes.

Lemma 6. A low sequence w.r.t. a minimum $\underline{c}$ -cover A_1 contains at least one loaded set.

Proof. Let $E = \{E_1, E_2, \dots, E_{2p}\}$ be a low sequence w.r.t. A_1 . Let P_{2p} be the union of the sets $E_1, E_2, \dots, E_{2p}$. Let $P^!$ be the union of those sets of A_1 which occur in the sequence E and $P^* = P_{2p} - P^!$. Let q be the number of points in each set of the complex (X, A) . It follows easily that

$$(4.1) \quad \sum_{x \in P_{2p}} |R_{2p}(x)| = \sum_{x \in P_{2p}} |\mathcal{Y}_{2p}(x)| = pq$$

If possible, suppose for no point x in $P^!$, $|A_1(x)| > c(x) + 1$.

Let

$$A' = A_1 - R_{2p}$$

$$A_2 = A' \cup \mathcal{Y}_{2p}$$

Since E is a low sequence w.r.t. A_1 , we have

(4.2)

$$\left| \left\{ x \mid x \in P_{2p}, |A_2(x)| > c(x) \right\} \right| > \left| \left\{ x \mid x \in P_{2p}, |A_1(x)| > c(x) \right\} \right|$$

Let

(4.3)

$$\left\{ x \mid x \in P_{2p}, |A_1(x)| > c(x) \right\} = G_1,$$

$$\left\{ x \mid x \in P_{2p}, |A_2(x)| > c(x) \right\} = G_2,$$

$$\left\{ x \mid x \in P^1, |A_1(x)| - c(x) = 1 \right\} = G_3,$$

$$\left\{ x \mid x \in P^*, |A_1(x) - c(x) \geq 1 \right\} = G_4,$$

$$\left\{ x \mid x \in P^1, |A_2(x)| - c(x) \geq 1 \right\} = G_5$$

and $\left\{ x \mid x \in P^* - G_4, |A_2(x)| - c(x) \geq 1 \right\} = G_6$

Since by our assumption for no point $x \in P^1$, $|A_1(x)| > c(x)+1$, it follows that G_1 is the union of the disjoint sets G_3 and G_4 .

Also it is easily checked that G_2 is the union of the disjoint sets

G_4 , G_5 and G_6 . From (4.2), we have

(4.4)

$$|G_5| + |G_6| > |G_3|.$$

Using (4.3) and (4.4), we get

(4.5)

$$\begin{aligned} \sum_{x \in P_{2p}} \{ |A_1(x)| + |A_2(x)| - c(x) \} &= \sum_{x \in P_{2p}} \{ |A_1(x)| - c(x) \} \\ &= \sum_{x \in G_4} \{ |A_2(x)| - c(x) \} + \sum_{x \in G_5} \{ |A_2(x)| - c(x) \} \\ &\quad + \sum_{x \in G_6} \{ |A_2(x)| - c(x) \} \end{aligned}$$

$$\begin{aligned}
&\geq \sum_{x \in G_4} \{ |A_1(x)| - c(x) \} + |G_5| + |G_6| \\
&> \sum_{x \in G_4} \{ |A_1(x)| - c(x) \} + |G_3| \\
&= \sum_{x \in G_4} \{ |A_1(x)| - c(x) \} + \sum_{x \in G_3} \{ |A_1(x)| - c(x) \} \\
&= \sum_{x \in P_{2p}} \{ |A_1(x)| - c(x) \} \\
&= \sum_{x \in P_{2p}} \{ |A_1(x)| + |R_{2p}(x)| - c(x) \}
\end{aligned}$$

Obviously (4.5) contradicts (4.1). Hence the low sequence E must contain at least one loaded set.

Lemma 7. Let A_1 be a minimum $\underline{c}$ -cover and A_0 be a minimum $\underline{c}$ -cover with minimum height. Let B_1 be a minimum $\underline{c}$ -cover in the family $M(A_0)$ nearest to A_1 . If all loaded sets of A_1 are contained in B_1 , A_1 is a minimum $\underline{c}$ -cover with minimum height.

Proof. It is sufficient to show that $A_1 \subseteq B_1$. If not, suppose $A_1 - B_1$ contains a set E_1 . Then by lemma 5 there exists a low sequence $E = \{E_1, E_2, \dots, E_{2p}\}$ w.r.t. A_1 where any set of the sequence E belonging to A_1 belongs to $A_1 - B_1$. By lemma 6, the sequence E must contain at least one loaded set of A_1 . This set also belongs to $A_1 - B_1$ which contradicts our hypothesis.

Theorem 6. A minimum $\underline{c}$ -cover A_1 of a regular complex (X, A) has minimum height if and only if there is no loaded low sequence w.r.t. A_1 .

Proof. Necessity is obvious. To prove sufficiency, assume that there is no loaded low sequence w.r.t. the minimum $\underline{c}$ -cover A_1 . Let A_0 be a minimum $\underline{c}$ -cover with minimum height and B_1 be a minimum $\underline{c}$ -cover in the

family $M(A_0)$ nearest to A_1 . By lemma 7 it is sufficient to show that all loaded sets of A_1 are contained in B_1 . If not, suppose $A_1 - B_1$ contains a loaded set E_1 . By lemma 5, there exists a low sequence $E = \{E_1, E_2, \dots, E_{2p}\}$ w.r.t. A_1 . Since E_1 is a loaded set, E is a loaded low sequence w.r.t. A_1 . This contradicts our hypothesis. Hence all loaded sets of A_1 are contained in B_1 . This completes the proof of the theorem.

Theorem 6 gives a simpler algorithm for the height of a regular complex. For a regular complex (X, A) , when searching for low sequences w.r.t. a minimum $\underline{c}$ -cover A_1 , it is necessary to start with loaded sets of A_1 only.

5. Relationship between maximum matching and minimum cover. Let (X, A) be a complex and $\underline{c} = (c(x_1), c(x_2), \dots, c(x_m))$ be a m -vector of positive integers. Let $\underline{d} = (d(x_1), d(x_2), \dots, d(x_m))$ be an associated m -vector where $d(x_i) = |A(x_i)| - c(x_i)$, $i = 1, 2, \dots, m$. Let A_1 be a $\underline{c}$ -matching of the complex.

Theorem 7. A $\underline{c}$ -matching A_1 is a maximum $\underline{c}$ -matching if and only if $B_1 = A - A_1$ is a minimum $\underline{d}$ -cover.

Proof. It is easily checked that if B_1 is a $\underline{d}$ -cover, $A_1 = A - B_1$ is a $\underline{c}$ -matching and vice, versa. If possible, suppose B_1 is a minimum $\underline{d}$ -cover and A_1 is not a maximum $\underline{c}$ -matching. Let A_0 be a maximum $\underline{c}$ -matching. Then $B_0 = A - A_0$ is a $\underline{d}$ -cover. Also we have

$$|A_1| + |B_1| = |A| = |A_0| + |B_0|$$

Since A_1 is not a maximum $\underline{c}$ -matching, $|A_1| < |A_0|$. Hence

$|B_1|$ must be greater than $|B_0|$ which contradicts the assumption that B_1 is a minimum $\underline{d}$ -cover. This completes the proof of the

theorem.

Similarly it can be proved that if $\mathcal{B}_1$ is a minimum $\underline{d}$ - cover with minimum height, $\mathcal{A}_1 = \mathcal{A} - \mathcal{B}_1$ is a maximum $\underline{c}$ - matching with maximum depth.

Acknowledgement. I wish to express my sincere thanks to Professor M. P. Schützenberger who taught me graph theory for suggesting the problem to me and for many stimulating discussions.

REFERENCES

1. C. Berge, "Two Theorems in Graph Theory", Proc. Nat. Acad. Sci. U.S.A. Vol. 43 (1957) pp. 842-844.
2. D. R. Fulkerson and H. J. Ryser, "Traces, Term Ranks, Widths and Heights," IBM Journal of Research and Development Vol 4 (1960), No. 5 pp. 455-459.
3. R. Z. Norman and M. O. Rabin, "An Algorithm for a Minimum Cover of a Graph," Proc. Amer. Math. Soc. Vol 10 (1959) pp 315-319.
4. J. Petersen, "Die Theorie der regulären Graphen," Acta Math. Vol 51 (1891) pp. 193-220.
5. J. P. Roth, "Algebraic Topological Methods for the Synthesis of Switching Systems I," Trans. Amer. Math. Soc. Vol. 88 (1958) pp. 302-326.