

* This research was supported by the Air Force Office of Scientific Research under Grant AFOSR-68-1415.

THE MEASURABILITY OF A STOCHASTIC PROCESS
OF SECOND ORDER AND ITS LINEAR SPACE

Stamatis Cambanis*

*Department of Statistics
University of North Carolina at Chapel Hill*

Institute of Statistics Mimeo Series No. 832

June, 1972

THE MEASURABILITY OF A STOCHASTIC PROCESS
OF SECOND ORDER AND ITS LINEAR SPACE

Stamatis Cambanis*

1. THE MEASURABILITY OF A STOCHASTIC
PROCESS OF SECOND ORDER

Let T be a separable metric space and $\mathcal{B}(T)$ the σ -algebra of Borel sets of T , and let $X_t, t \in T$, be a real stochastic process on the probability space (Ω, \mathcal{F}, P) . $X_t, t \in T$, is called measurable if the map $(t, \omega) \rightarrow X_t(\omega)$ is $\mathcal{B}(T) \times \mathcal{F}$ -measurable. A process $Y_t, t \in T$, on (Ω, \mathcal{F}, P) is called a modification of $X_t, t \in T$, if $P\{X_t = Y_t\} = 1$ for all t in T . $X_t, t \in T$, is of second order if $E(X_t^2) < +\infty$ for all t in T , and then its autocorrelation R is defined by $R(t, s) = E(X_t X_s)$ for all t, s in T . It is clear from Fubini's theorem that if a second order process $X_t, t \in T$, has a measurable modification then R is $\mathcal{B}(T) \times \mathcal{B}(T)$ -measurable. That the measurability of R is not sufficient for the existence of a measurable modification of $X_t, t \in T$, is demonstrated in Remark 2. It is thus of interest to find a condition which along with the measurability of R would imply the existence of a measurable modification of $X_t, t \in T$. This question is answered in Theorem 1, where in fact necessary and sufficient conditions are given for a second order process to have a measurable modification. A remarkable consequence of these conditions is that the existence of a measurable modification of a second order process is a second order property.

* This research was supported by the Air Force Office of Scientific Research under Grant AFOSR-68-1415.

The proof of Theorem 1 is based on the necessary and sufficient conditions for a process (not necessarily of second order) to have a measurable modification given in [5], which are expressed as follows (here the terminology of [6] is followed). Let M be the space of all real random variables on (Ω, \mathcal{F}, P) with the topology of convergence in probability, where random variables that are equal a.e. $[P]$ are considered identical. If ξ is a real random variable, its class in M is denoted by $[\xi]$. Then $X_t, t \in T$, has a measurable modification if and only if the map from T to M taking t to $[X_t]$ is measurable and has separable range [5,6]. Moreover, the measurable modification can be taken to be separable and also progressively measurable, the latter if T is an interval and a nondecreasing family $\mathcal{F}_t, t \in T$, of sub- σ -algebras of \mathcal{F} is given.

For a second order process $X_t, t \in T$, we denote by $H(X)$ the closure in $L_2(\Omega, \mathcal{F}, P)$ of the linear space of the random variables $\{X_t, t \in T\}$ and we call it the linear space of the process. We also denote by $R(K)$ the reproducing kernel Hilbert space of a real, symmetric, nonnegative definite function K on $T \times T$. It is well known that $R(K)$ consists of all functions f on T of the form $f(t) = E(\xi X_t)$, $t \in T$, for some $\xi \in H(X)$, and that the map $\xi \rightarrow E(\xi X_t)$ defines an inner product preserving isomorphism between $H(X)$ and $R(K)$ [16, p.302].

THEOREM 1. Let $X_t, t \in T$, be a real, second order process with autocorrelation R . The following are equivalent.

- (i) $X_t, t \in T$, has a measurable modification.
- (ii) R is $\mathcal{B}(T) \times \mathcal{B}(T)$ -measurable and $H(X)$ (or $R(K)$) is separable.

PROOF. (a) We first show that (ii) implies (i). It suffices to verify the conditions of [5,6]; the construction of a measurable modi-

fication is the same as in [5] or in [6].

Since convergence in $L_2(\Omega, \mathcal{F}, P)$ implies convergence in probability, the separability of $H(X)$ as a subset of $L_2(\Omega, \mathcal{F}, P)$ implies its separability as a subset of M . Thus its subset $\{[X_t], t \in T\}$ is separable in M . To complete the proof it suffices to show that the map $X: T \rightarrow M$ defined by $X(t) = [X_t]$ is measurable. The metric ρ on M defined by $\rho(\xi, \eta) = E \left(\frac{|\xi - \eta|}{1 + |\xi - \eta|} \right)$, $\xi, \eta \in M$, metrizes the topology of convergence in probability. Thus for the measurability of X it suffices to show that $X^{-1}(B) \in \mathcal{F}$ for every set B in M of the form $B = \{Y \in M: \rho(Y, Y_0) \leq r\}$, where $Y_0 \in M$ and $r > 0$. Since $X^{-1}(B) = \{t \in T: \rho([X_t], Y_0) \leq r\}$, it suffices to prove that the real function $\rho([X_t], Y_0)$ on T is $\mathcal{B}(T)$ -measurable for all $Y_0 \in M$.

Let $\{\xi_n\}_{n=1}^{\infty}$ be a complete orthonormal sequence in $H(X)$ (which exists because $H(X)$ is separable). Then for all $t \in T$ we have

$$X_t = \sum_{n=1}^{\infty} a_n(t) \xi_n$$

in $L_2(\Omega, \mathcal{F}, P)$, where $a_n(t) = E(\xi_n X_t)$. Thus $a_n \in R(R)$, and in fact $\{a_n\}_{n=1}^{\infty}$ is a complete orthonormal sequence in $R(R)$. If for every $t \in T$ we let $X_t^{(N)} = \sum_{n=1}^N a_n(t) \xi_n$, then $X_t^{(N)}$ converges to X_t in $L_2(\Omega, \mathcal{F}, P)$ and thus in probability. Let $Y_t = X_t - Y_0$ and $Y_t^{(N)} = X_t^{(N)} - Y_0$ for all $t \in T$. Then $Y_t^{(N)}$ converges to Y_t in probability, i.e., $\rho([Y_t^{(N)}], [Y_t]) \rightarrow 0$ as $N \rightarrow \infty$. Dropping the index t for simplicity we have

$$\left| \frac{|Y^{(N)}|}{1 + |Y^{(N)}|} - \frac{|Y|}{1 + |Y|} \right| = \frac{||Y^{(N)}| - |Y||}{(1 + |Y^{(N)}|)(1 + |Y|)} \leq \frac{|Y^{(N)} - Y|}{1 + |Y^{(N)} - Y|}$$

Thus

$$E \left| \frac{|Y^{(N)}|}{1 + |Y^{(N)}|} - \frac{|Y|}{1 + |Y|} \right| \leq \rho([Y^{(N)}], [Y]) \xrightarrow{N \rightarrow \infty} 0.$$

It follows that for all $t \in T$,

$$\rho([X_t], Y_0) = \lim_{N \rightarrow \infty} E \left[\frac{|X_t^{(N)} - Y_0|}{1 + |X_t^{(N)} - Y_0|} \right].$$

Note that every function in $R(R)$ is either a finite linear combination of the functions $\{R(t, \cdot), t \in T\}$ or a pointwise limit on T of such functions. Hence, since R is $B(T) \times B(T)$ - measurable, $R(t, \cdot)$ is $B(T)$ - measurable for all $t \in T$, and it follows that every f in $R(R)$ is $B(T)$ - measurable. Consequently $Y_t^{(N)}(\omega)$ is $B(T) \times F$ - measurable. By Fubini's theorem

$E \left[\frac{|X_t^{(N)} - Y_0|}{1 + |X_t^{(N)} - Y_0|} \right]$ is $B(T)$ - measurable, and thus so is $\rho([X_t], Y_0)$, which completes the proof.

(b) We now show that (i) implies (ii). The measurability of R follows from Fubini's theorem and (i). In order to prove the separability of $H(X)$ we first assume that R is uniformly bounded on T :

$$R(t, t) \leq C < +\infty \text{ for all } t \text{ in } T.$$

We will show that this implies the uniform integrability of the family of random variables $\{X_t, t \in T\}$. Indeed we have for all $a > 0$,

$$\begin{aligned} \int_{|X_t| > a} |X_t| \, dP &= \int_{\Omega} I_{\{|X_t| > a\}} |X_t| \, dP \\ &\leq [P\{|X_t| > a\} \cdot R(t, t)]^{\frac{1}{2}} \\ &\leq \frac{R(t, t)}{a}. \end{aligned}$$

Thus

$$\lim_{a \rightarrow \infty} \sup_{t \in T} \int_{|X_t| > a} |X_t| \, dP \leq \lim_{a \rightarrow \infty} \frac{C}{a} = 0$$

and $\{X_t, t \in T\}$ is uniformly integrable.

Now (i) implies that $\{[X_t], t \in T\}$ is separable in M . Thus there exists a countable subset M' of $\{[X_t], t \in T\}$ such that for every t in T , $[X_t]$ is the limit in probability of a sequence in M' , and hence also in $L_2(\Omega, \mathcal{F}, P)$, since M' is uniformly integrable [14, p. 57]. It follows that $H(X)$ equals the $L_2(\Omega, \mathcal{F}, P)$ closure of the linear span of M' and, since M' is countable, $H(X)$ is separable.

We now consider the general case and define for $N = 1, 2, \dots$,

$$T_N = \{t \in T: R(t, t) \leq N\}.$$

Since R is measurable, $T_N \in \mathcal{B}(T)$ and by (i) $\{X_t, t \in T_N\}$ has a measurable modification. It follows by what has been proven that the $L_2(\Omega, \mathcal{F}, P)$ closure of the linear span of the random variables $\{X_t, t \in T_N\}$, $H_N(X)$, is separable. Since X_t is of second order, R is finite valued and thus $T_N \uparrow T$. It follows that $H(X)$ is the $L_2(\Omega, \mathcal{F}, P)$ closure of $\bigcup_{N=1}^{\infty} H_N(X)$ and thus $H(X)$ is separable. \square

Thus a $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable, symmetric, nonnegative definite, real function R on $T \times T$ is the autocorrelation of a measurable process if and only if $R(R)$ is separable.

REMARK 1. The mean m and the covariance C of a real second order process $X_t, t \in T$, are defined by $m(t) = E(X_t)$ and $C(t, s) = E([X_t - m(t)][X_s - m(s)])$ for all t, s , in T . Then $R(t, s) = m(t)m(s) + C(t, s)$. In connection with (ii) of Theorem 1 it should be noted that

R is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable if and only if m is $\mathcal{B}(T)$ - measurable and C is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable.

The "if" part is obvious. The "only if" part is shown as follows. We

have $m(t) = E(X_t I_\Omega)$ for all t in T , where I is the indicator function. Denote by ξ the projection of $I_\Omega \in L_2(\Omega, \mathcal{F}, P)$ onto the subspace $H(X)$. Then $m(t) = E(X_t \xi)$ for all t in T and $\xi \in H(X)$, and thus $m \in \mathcal{R}(R)$. Since R is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable, m is $\mathcal{B}(T)$ - measurable (see part (a) of the proof of Theorem 1) and $C(t, s) = R(t, s) - m(t)m(s)$ is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable.

REMARK 2. Let $T = [0, 1]$ and $R(t, s) = 1$ for $t = s$ in T and $R(t, s) = 0$ for $t \neq s$ in T . Since R is symmetric and nonnegative definite, there exists a probability space (Ω, \mathcal{F}, P) and a real process $X_t, t \in T$, on it with autocorrelation R . R is clearly $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable, but since the values of X_t are orthogonal in $L_2(\Omega, \mathcal{F}, P)$, $E(X_t X_s) = 0$ for $t \neq s$ in T , $H(X)$ is not separable and by Theorem 1, $X_t, t \in T$, does not have a measurable modification. This can be also shown without using Theorem 1. Indeed, assume that $X_t, t \in T$, has a measurable modification $Y_t, t \in T$. Then

$$E\left(\int_0^1 Y_t^2 dt\right) = \int_0^1 R(t, t) dt = 1 < +\infty$$

implies that $\int_0^1 Y_t^2 dt < +\infty$ a.e. $[P]$. If $\{\phi_n\}_{n=1}^\infty$ is a complete orthonormal set in $L_2(T) = L_2(T, \mathcal{B}(T), \text{Leb})$ then

$$Y_t = \sum_{n=1}^{\infty} \xi_n \phi_n(t)$$

in $L_2(T)$ a.e. $[P]$, where $\xi_n = \int_0^1 Y_t \phi_n(t) dt$ a.e. $[P]$. Then

$$E(\xi_n^2) = \int_0^1 \int_0^1 R(t, s) \phi_n(t) \phi_n(s) dt ds = 0$$

i.e., $\xi_n = 0$ a.e. $[P]$, and thus $\int_0^1 Y_t^2 dt = \sum_{n=1}^{\infty} \xi_n^2 = 0$ a.e. $[P]$ which contradicts $E\left(\int_0^1 Y_t^2 dt\right) = 1$. It follows that $X_t, t \in T$, does not have a measurable modification.

REMARK 3. For Gaussian processes it can be easily shown that (ii) implies (i) without relying on the results of [5]; this is done in [15, p. 44].

COROLLARY 1. Let R be a symmetric, nonnegative definite, real function on $T \times T$. If $R(R)$ is separable the following are equivalent.

- (i) $R(t, \cdot)$ is $\mathcal{B}(T)$ - measurable for all t in T .
- (ii) R is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable.

PROOF. It suffices to show that (i) implies (ii). Since R symmetric, nonnegative definite and real, there exists a probability space (Ω, \mathcal{F}, P) and a real process $X_t, t \in T$, on it with autocorrelation R . It is clear from part (a) of the proof of Theorem 1 that the separability of $R(R)$ and (i) imply the existence of a measurable modification of $X_t, t \in T$, and thus (ii). This result can be shown in a simpler way without using an associated process. Indeed, if $\{a_n\}_{n=1}^{\infty}$ is a complete orthonormal set in $R(R)$, then it is easily seen that $R(t, s) = \sum_{n=1}^{\infty} a_n(t) a_n(s)$ for all t, s in T . Now (i) implies as in part (a) of the proof of Theorem 1 that every a_n is $\mathcal{B}(T)$ - measurable and thus (ii) holds. \square

COROLLARY 2. A second order process $X_t, t \in T$, which satisfies any of the following conditions has a measurable modification (in (iii) also progressively measurable).

- (i) $X_t, t \in T$, is weakly continuous on T .
- (ii) T is an arbitrary interval and $X_t, t \in T$, has orthogonal increments.
- (iii) T is an arbitrary interval and $X_t, t \in T$, is a martingale.

PROOF. (i) Since T is separable and X_t weakly continuous on T , $H(X)$ is separable [16, p. 272]. By the weak continuity of $X_t, R(t, \cdot)$

is continuous, hence $\mathcal{B}(T)$ - measurable, for all t in T . The conclusion follows from Corollary 1 and Theorem 1.

(ii) It is known that $H(X)$ is separable [8, p. 110]. Also, that X_t has left and right $L_2(\Omega, \mathcal{F}, P)$ limits on T and that except on a countable subset of T , $X_{t-} = X_t = X_{t+}$. This implies the measurability of R and the result follows from Theorem 1.

(iii) Define the function F by $F(t) = E(X_t^2)$ for all t in T . By the martingale property, with respect to the nondecreasing family \mathcal{F}_t , $t \in T$, of sub- σ -algebras of \mathcal{F} , we have for all $s \leq t$ in T , $E(X_t X_s) = E[E(X_t X_s / \mathcal{F}_s)] = E[X_s E(X_t / \mathcal{F}_s)] = E(X_s^2)$ and thus

$$E\{(X_t - X_s)^2\} = F(t) - F(s).$$

It follows from this relationship, as in [8, p. 110] and in (ii), that $H(X)$ is separable and R is $\mathcal{B}(T) \times \mathcal{B}(T)$ - measurable. \square

REMARK 4. Let X_t , $t \in T$, T an arbitrary interval, be a real separable process of second order with autocorrelation R . If X_t is mean square differentiable on T and $\frac{\partial R(t,s)}{\partial t}$, $\frac{\partial^2 R(t,s)}{\partial t \partial s}$ are locally Lebesgue integrable in t and in t, s respectively, then with probability one the paths of X_t , $t \in T$, are absolutely continuous on every compact subinterval of T . This is shown in [10, pp. 186-187] with the additional assumption that the mean square derivative X_t' of X_t has a measurable modification, which is always satisfied because of Theorem 1. Indeed, since X_t is mean square differentiable on T , it is mean square continuous on T . Thus $H(X)$ is separable and the continuity of R implies the measurability of $\frac{\partial^2 R(t,s)}{\partial t \partial s}$. Since $\frac{\partial^2 R(t,s)}{\partial t \partial s}$ is the autocorrelation of X_t' and since $H(X') \subseteq H(X)$, the conclusion follows from Theorem 1.

We conclude this section with a property which is useful in connection with problems involving conditional probabilities; such as for instance the existence of jointly measurable densities (see [9, pp. 616-617]) and properties related to metric transitivity (see [17, Ch. IV. 8]). A σ -algebra is called separable if it is generated by a countable class of sets. A sub- σ -algebra F' of F is said to coincide mod 0 with the σ -algebra F if for every set E in F there is a set E' in F' such that $P(E \Delta E') = 0$. Let $F(X)$ be the sub- σ -algebra of F generated by the random variables $\{X_t, t \in T\}$. It is known that if X_t is continuous in probability on T , $F(X)$ coincides mod 0 with a separable σ -algebra. Corollary 3 generalizes this result (and in fact, as it is clear from [6], it is valid for any process with values in a compact metric space).

COROLLARY 3. If a real process $X_t, t \in T$, has a measurable modification, then $F(X)$ coincides mod 0 with a separable σ -algebra.

PROOF. Since $X_t, t \in T$, has a measurable modification, $\{[X_t], t \in T\}$ is a separable subset of M . Thus there exists a countable subset $M' = \{[X_t], t \in S\}$ of $\{[X_t], t \in T\}$ (S is a countable subset of T) such that for every t in T , $[X_t]$ is the limit in probability of a sequence from M' , and thus X_t is the a.e.[P] limit of a sequence from $\{X_t, t \in S\}$. If F' is the sub- σ -algebra of F generated by the random variables $\{X_t, t \in S\}$, then $F' \subseteq F(X)$, F' is separable and $F(X)$ coincides with F' mod 0. \square

2. ON THE SEPARABILITY OF THE LINEAR SPACE OF A SECOND ORDER PROCESS

The linear space $H(X)$ of a second order process $X_t, t \in T$, plays an important role in the structure of such processes and in a variety of problems in statistical inference. If $H(X)$ is separable then X_t admits

series representations and also integral representations (Theorem 2) that can be effectively used in problems such as linear mean square estimation. Also the separability of $H(X)$ is the only condition needed for a second order process to have the Hida-Cramér representation (see for instance [11]). It is thus of interest that a measurable second order process has a separable linear space. $H(X)$ is known to be separable when the process X_t , $t \in T$, is weakly continuous [16, p. 272], has orthogonal increments [8, p. 110], or is a martingale (Corollary 2.(iii)). In Theorem 2 necessary and sufficient conditions are given for $H(X)$ to be separable in terms of integral representations of X_t .

Before stating the theorem we mention a few basic facts about random measures, that can be found for instance in [7, 16]. Let (V, \mathcal{V}) be a measurable space. A random measure Z on (V, \mathcal{V}) is a countably additive map from \mathcal{V} to $L_2(\Omega, \mathcal{F}, P)$; i.e., whenever A is the disjoint union of the sets $A_n \in \mathcal{V}$, $Z(A) = \sum_{n=1}^{\infty} Z(A_n)$ in $L_2(\Omega, \mathcal{F}, P)$. (Here we consider the case where Z is defined on the entire σ -algebra \mathcal{V}). To each random measure Z on V there corresponds a finite signed measure μ on $V \times V$ by $\mu(A \times B) = E[Z(A)Z(B)]$, $A, B \in \mathcal{V}$. μ is symmetric and nonnegative definite on the measurable rectangles of $V \times V$. A random measure Z is called orthogonal if $\mu(A \times B) = 0$ whenever A and B are disjoint, and to each orthogonal random measure there corresponds a finite nonnegative measure ν on V by $\nu(A) = E[Z^2(A)]$, $A \in \mathcal{V}$. Let $H(Z)$ be the closure in $L_2(\Omega, \mathcal{F}, P)$ of the linear span of $\{Z(A), A \in \mathcal{V}\}$, and let $\Lambda_2(\mu)$ be the Hilbert space of real, V -measurable functions on V with inner product $\langle f, g \rangle_{\Lambda_2(\mu)} = \int \int f(u)g(v) d\mu(u, v)$ (of course $\Lambda_2(\mu)$ consists of equivalence classes of functions, two functions f and g considered identical if $\langle f-g, f-g \rangle_{\Lambda_2(\mu)} = 0$). There is an inner product preserving isomorphism between $\Lambda_2(\mu)$ and $H(Z)$,

denoted by \leftrightarrow , such that $I_A \leftrightarrow Z(A)$, $A \in \mathcal{V}$, and integration of functions in $\Lambda_2(\mu)$ with respect to Z is defined by $\xi = \int_{\mathcal{V}} f(u) dZ(u)$, where $f \leftrightarrow \xi$. If Z is orthogonal, there is an inner product preserving isomorphism between $L_2(\nu) = L_2(\mathcal{V}, \mathcal{V}, \nu)$ and $H(Z)$, denoted again by \leftrightarrow , such that $I_A \leftrightarrow Z(A)$, $A \in \mathcal{V}$, and integration of functions in $L_2(\nu)$ with respect to Z is defined by $\xi = \int_{\mathcal{V}} f(u) dZ(u)$, where $f \leftrightarrow \xi$.

THEOREM 2. Let X_t , $t \in T$ be a second order process.

(i) If $H(X)$ is separable then for every finite measure space $(\mathcal{V}, \mathcal{V}, \nu)$ such that $L_2(\nu) = L_2(\mathcal{V}, \mathcal{V}, \nu)$ is separable and infinite dimensional, X_t has a representation

$$X_t = \int_{\mathcal{V}} f(t, u) dZ(u) \quad \text{for all } t \text{ in } T$$

where Z is an orthogonal measure on \mathcal{V} with corresponding measure ν and $f(t, \cdot) \in L_2(\nu)$ for all t in T . Conversely, if X_t has such a representation, $H(X)$ is separable.

(ii) If $H(X)$ is separable, then for every measurable space $(\mathcal{V}, \mathcal{V})$ and every finite signed measure μ on $\mathcal{V} \times \mathcal{V}$ which is symmetric and non-negative definite on the measurable rectangles of $\mathcal{V} \times \mathcal{V}$, and such that $\Lambda_2(\mu)$ is separable and infinite dimensional, X_t has a representation

$$X_t = \int_{\mathcal{V}} f(t, u) dZ(u) \quad \text{for all } t \text{ in } T$$

where Z is a random measure on \mathcal{V} with corresponding measure μ and $f(t, \cdot) \in \Lambda_2(\mu)$ for all t in T . Conversely, if X_t has such a representation, $H(X)$ is separable.

PROOF. (i) being a particular case of (ii), we will prove only (ii). We start with the second claim. If X_t has such a representation then $X_t \in H(Z)$ for all t in T , hence $H(X) \subseteq H(Z)$ and the conclusion follows

from the isomorphism between $H(Z)$ and $\Lambda_2(\mu)$ and the separability of the latter. We now prove the first claim. Assume that $H(X)$ is separable and let $\{\xi_n\}_{n=1}^{\infty}$ be a complete orthonormal set. Then for all t in T

$$X_t = \sum_{n=1}^{\infty} a_n(t) \xi_n$$

in $L_2(\Omega, \mathcal{F}, P)$, where $a_n(t) = E(X_t \xi_n)$. Let $\{f_n\}_{n=1}^{\infty}$ be a complete orthonormal set in $\Lambda_2(\mu)$. Since μ is finite, $I_A \in \Lambda_2(\mu)$ for all $A \in \mathcal{V}$.

Then

$$I_A = \sum_{n=1}^{\infty} \lambda_n(A) f_n$$

in $\Lambda_2(\mu)$, where

$$\lambda_n(A) = \langle I_A, f_n \rangle = \int_A \int_V f_n(v) d\mu(u, v).$$

Throughout the proof we will write $\langle \cdot, \cdot \rangle$ for $\langle \cdot, \cdot \rangle_{\Lambda_2(\mu)}$. Thus for all n , λ_n is a finite signed measure on (V, \mathcal{V}) . We also have

$$\sum_{n=1}^{\infty} \lambda_n^2(A) = \langle I_A, I_A \rangle = \mu(A \times A) < +\infty.$$

Hence

$$Z(A) = \sum_{n=1}^{\infty} \lambda_n(A) \xi_n$$

defines a function from \mathcal{V} to $L_2(\Omega, \mathcal{F}, P)$ (the convergence being in $L_2(\Omega, \mathcal{F}, P)$).

We will show that Z is a random measure with corresponding measure μ .

The latter is clear since for all $A, B \in \mathcal{V}$ we have

$$E[Z(A)Z(B)] = \sum_{n=1}^{\infty} \lambda_n(A) \lambda_n(B) = \langle I_A, I_B \rangle = \mu(A \times B).$$

For the countable additivity of Z let $A = \bigcup_{k=1}^{\infty} A_k$, where $\{A_k\}_{k=1}^{\infty}$ is a

disjoint sequence of sets in V . Then

$$\begin{aligned}
 E\left[\left\{Z(A) - \sum_{k=1}^K Z(A_k)\right\}^2\right] &= \sum_{n=1}^{\infty} \left\{\lambda_n(A) - \sum_{k=1}^K \lambda_n(A_k)\right\}^2 \\
 &= \sum_{n=1}^{\infty} \left\{\sum_{k=K}^{\infty} \lambda_n(A_k)\right\}^2 \\
 &= \sum_{n=1}^{\infty} \lambda_n^2\left(\bigcup_{k=K}^{\infty} A_k\right) \\
 &= \mu\left(\bigcup_{k=K}^{\infty} A_k \times \bigcup_{k=K}^{\infty} A_k\right) \xrightarrow{K \rightarrow \infty} 0
 \end{aligned}$$

since $\bigcup_{k=K}^{\infty} A_k \downarrow \emptyset$ as $K \rightarrow \infty$. Thus $Z(A) = \sum_{k=1}^{\infty} Z(A_k)$.

We now show that for every g in $\Lambda_2(\mu)$,

$$\int_V g dZ = \sum_{n=1}^{\infty} \langle g, f_n \rangle \xi_n$$

in $L_2(\Omega, \mathcal{F}, P)$. This is true for indicator functions by definition of Z , and therefore also for simple functions. Since $H(Z)$ is defined as the $L_2(\Omega, \mathcal{F}, P)$ closure of the linear space of $\{Z(A), A \in V\}$, it follows by the isomorphism between $\Lambda_2(\mu)$ and $H(Z)$ that the linear span of $\{I_A, A \in V\}$ is dense in $\Lambda_2(\mu)$. Thus every g in $\Lambda_2(\mu)$ is the $\Lambda_2(\mu)$ - limit of a sequence of simple functions $\{g_k\}_{k=1}^{\infty}$. Thus

$$\begin{aligned}
 \int_V g dZ &= \lim_{k \rightarrow \infty} \int_V g_k dZ \\
 &= \lim_{k \rightarrow \infty} \sum_{n=1}^{\infty} \langle g_k, f_n \rangle \xi_n
 \end{aligned}$$

where $\lim_{k \rightarrow \infty}$ is in $L_2(\Omega, \mathcal{F}, P)$ and the result follows from

$$E\left[\left\{\sum_{n=1}^{\infty} \langle g, f_n \rangle \xi_n - \sum_{n=1}^{\infty} \langle g_k, f_n \rangle \xi_n\right\}^2\right]$$

$$\begin{aligned}
&= E\left[\left\{\sum_{n=1}^{\infty} \langle g-g_k, f_n \rangle \xi_n\right\}^2\right] \\
&= \sum_{n=1}^{\infty} \langle g-g_n, f_n \rangle^2 \\
&= \langle g-g_k, g-g_k \rangle \xrightarrow{k \rightarrow \infty} 0.
\end{aligned}$$

In particular we have $\int_V f_n dZ = \xi_n$ which implies that $H(Z) = H(X)$.

Now since $\sum_{n=1}^{\infty} a_n^2(t) = R(t,t) < +\infty$ for all t in T , we can define $f(t, \cdot)$ in $L_2(\mu)$ for all t in T by

$$f(t, u) = \sum_{n=1}^{\infty} a_n(t) f_n(u)$$

where the convergence is in $L_2(\mu)$. It follows from the property of the integral just proven that for all t in T we have the following equality in $L_2(\Omega, F, P)$,

$$\int_V f(t, u) dZ(u) = \sum_{n=1}^{\infty} a_n(t) \xi_n = X_t$$

which concludes the proof. \square

REMARK 5. We assume throughout this remark that $H(X)$ is separable. Then it is clear that the first claim in (i) and (ii) is valid provided the dimensionality of $L_2(\nu)$ and $L_2(\mu)$ is no less than the dimensionality of the integers. Also, one can take $(V, \mathcal{V}) = (T, \mathcal{B}(T))$ or as V any interval and \mathcal{V} its Borel sets; in the latter case ν may be taken the Lebesgue measure or one absolutely continuous to it, and μ may be taken absolutely continuous to the Lebesgue measure on $V \times V$. If a series (respectively, integral) representation of X_t is known then one can obtain integral (respectively, series) representations of X_t as indicated in the proof of Theorem 2. These representations will be explicitly obtained if one can find complete orthonormal sets in the spaces

$L_2(\nu)$ and $\Lambda_2(\mu)$. If V is an interval and \mathcal{V} its Borel sets, complete orthonormal sets in $L_2(\nu)$ are given in [13] (see also [2]), and complete sets in $\Lambda_2(\mu)$ are given in [3] (In [3] the case where V is the entire real line is treated and the case where V is an interval can be treated similarly). If neither an integral nor a series representation of X_t is available, the problem arises how to obtain explicitly such a representation (in terms of the process X_t $t \in T$, and its autocorrelation R). This problem is solved in [4] for weakly continuous processes X_t $t \in T$, and T an arbitrary interval.

REMARK 6. Theorem 2 may also be stated in terms of integral representation of the autocorrelation R , which for (i) and (ii) are respectively

$$R(t,s) = \int_V f(t,u) f(s,u) d\nu(u) \quad \text{for all } t,s \text{ in } T.$$

$$R(t,s) = \int_V \int_V f(t,u) f(s,v) d\mu(u,v)$$

REMARK 7. In [12] a second order process X_t , $t \in \mathbb{R}^1 = (-\infty, +\infty)$ is called oscillatory if it has a representation

$$X_t = \int_{-\infty}^{\infty} e^{itu} a_t(u) dZ(u) \quad \text{for all } t \text{ in } \mathbb{R}^1$$

where Z is an orthogonal random measure on $(\mathbb{R}^1, \mathcal{B}(\mathbb{R}^1))$ with corresponding measure ν and $a_t(\cdot) \in L_2(\nu)$ for all t in T (this is a generalization of a concept introduced by Priestley). If X_t , $t \in \mathbb{R}^1$, is oscillatory then $H(X)$ is separable, since $L_2(\mathbb{R}^1, \mathcal{B}(\mathbb{R}^1), \nu)$ is separable. Conversely, if $H(X)$ is separable it follows by Theorem 1. (i) that for any finite measure ν on $(\mathbb{R}^1, \mathcal{B}(\mathbb{R}^1))$ we have $X_t = \int_{-\infty}^{\infty} f(t,u) dZ(u)$ for all t in T , where Z is an orthogonal random measure on $(\mathbb{R}^1, \mathcal{B}(\mathbb{R}^1))$ with corresponding

measure ν and $f(t, \cdot) \in L_2(\nu)$ for all t in \hat{T} . If we define $a_t(u) = e^{-itu} f(t, u)$, it becomes clear that $X_t, t \in \mathbb{R}^1$, is oscillatory. Thus a second order process is oscillatory if and only if its linear space is separable.

REMARK 8. Some simple sufficient conditions for $H(X)$ to be separable are as follows. If $X_t, t \in T$, is a linear operation on a second order process $Y_s, s \in S$, with separable linear space, then $H(X) \subseteq H(Y)$ and the separability of $H(X)$ follows from that of $H(Y)$. Also, because of the isomorphism between $H(X)$ and $\mathcal{R}(R)$, $H(X)$ is separable if there is a symmetric, nonnegative definite function K on $T \times T$ such that $\mathcal{R}(R) \subseteq \mathcal{R}(K)$ and $\mathcal{R}(K)$ is separable. A sufficient condition for $\mathcal{R}(R) \subseteq \mathcal{R}(K)$ is that $K-R$ be nonnegative definite [1, p. 354].

REFERENCES

1. Aronszajn, N.: Theory of reproducing kernels. Trans. Amer. Math. Soc. 68, 337-404 (1950).
2. Cambanis, S.: Bases in L_2 spaces with applications to stochastic processes with orthogonal increments. Proc. Amer. Math. Soc. 29, 284-290 (1971).
3. Cambanis, S., Liu, B.: On harmonizable stochastic processes. Information and Control 17, 183-202 (1970).
4. Cambanis, S., Masry, E.: On the representation of weakly continuous processes. Information Sci. 3, 277-290 (1971).
5. Chung, K.L., Doob, J.L.: Fields, optionality and measurability. Amer. J. Math. 87, 397-424 (1965).
6. Cohn, D.L.: Measurable choice of limit points and the existence of separable and measurable processes. Z. Wahrscheinlichkeitstheorie verw. Geb. 22, 161-165 (1972).
7. Cramér, H.: A contribution to the theory of stochastic processes. Proc. Second Berkeley Symp. Math. Statist. Prob., 329-339. Berkeley: Univ. California Press 1951.
8. Cramér, H., Leadbetter, M.R.: Stationary and Related Stochastic Processes. New York: Wiley 1967.
9. Doob, J.L.: Stochastic Processes. New York: Wiley 1953.
10. Gikhman, I.I., Skorokhod, A.V.: Introduction to the Theory of Random Process. Philadelphia: Saunders 1969.
11. Kallianpur, G., Mandrekar, V.: Multiplicity and representation theory of purely non-deterministic stochastic processes. Theor. Probability Appl. 10, 553-581 (1965).

12. Mandrekar, V.: A characterization of oscillatory processes and their prediction. Proc. Amer. Math. Soc. 32, 280-284 (1972).
13. Masry, E., Liu, B., Steiglitz, K.: Series expansion of wide-sense stationary random processes. IEEE Trans. Information Theory 14, 792-796 (1968).
14. Neveu, J.: Mathematical Foundations of the Calculus of Probability. San Francisco: Holden-Day 1965.
15. Neveu, J.: Processus Aléatoires Gaussiens. Montréal: Presses Univ. Montréal 1968.
16. Parzen, E.: Time Series Analysis Papers. San Francisco: Holden-Day 1967.
17. Rasanov, Yu. A.: Stationary Random Processes. San Francisco: Holden-Day 1967.

Stamatis Cambanis
Department of Statistics
University of North Carolina
Chapel Hill, N.C. 27514