

Reference List

- Allen, G. C., Hall, G. E., Childs, L. C., Weissinger, A. K., Spiker, S. and Thompson, W. F. (1993) Scaffold attachment regions increase reporter gene expression in stably transformed plant cells. *Plant Cell* 5: 603-613.
- Allen, G. C., Hall, G., Michalowski, S., Newman, W., Spiker, S., Weissinger, A. K. and Thompson, W. F. (1996) High-level transgene expression in plant cells: Effects of a strong scaffold attachment region from tobacco. *Plant Cell* 8: 899-913.
- Allen, G. C., Spiker, S. and Thompson, W. F. (2000) Use of matrix attachment regions (MARs) to minimize transgene silencing. *Plant Molecular Biology* 43: 361-376.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389-3402.
- Avramova, Z. and Bennetzen, J.L. (1993) Isolation of matrices from maize leaf nuclei: identification of a matrix-binding site adjacent to the *Adh1* gene. *Plant Molecular Biology* 22: 1135-1143.
- Balasubramaniam U. and Oleinick, N.L. (1995) Preferential cross-linking of matrix-attachment region (MAR) containing DNA fragments to the isolated nuclear matrix by ionizing radiation. *Biochemistry* 34: 12790-12802.
- Banan, M., I.C., Rojas, W.-H., Lee, H.L., King, J.V., Harriss, R., Kobayashi, Webb, C.F. and Gottlieb, P.D. (1997) Interaction of the nuclear matrix-associated region (MAR)-binding proteins, SATB1 and CDP/Cux, with a MAR element (L2a) in an upstream regulatory region of the mouse CD8a gene. *Journal of Biological Chemistry* 272: 18440-18452.
- Bednar, J., Horowitz, R.A., Grigoryev, S. A., Carruthers, L.M., Hansen, J.C., Koster, A.J. and Woodcock, C.L. (1998) Nucleosomes, linker DNA, and linker histone form a unique structural motif that directs the higher-order folding and compaction of chromatin. *Proceedings of the National Academy of Sciences* 95: 14173-14178.
- Belgrader, P., Siegel, A.J. and Berezney, R. (1991) A comprehensive study on the isolation and characterization of the HeLa S3 nuclear matrix. *Journal of Cell Science* 98: 281-291.
- Benham, C., Kohwi-Shigematsu, T. and Bode, J. (1997) Stress-induced duplex DNA destabilization in scaffold/matrix attachment regions. *Journal of Molecular Biology* 274: 181-196.
- Berezney, R. and Coffey, D. S. (1974) Identification of a nuclear protein matrix. *Biochemical and Biophysical Research Communications* 60: 1410-1417.
- Berezney, R. Mortyillaro, Ma, J., Wei, X. and Samarabandu, J. (1995) The nuclear matrix: a structural milieu for genomic function. *International Review of Cytology* 162A: 1-65.
- Berrios M., Osheroff, N. and Fisher, P.A. (1985) In situ localization of DNA topoisomerase II, a major polypeptide component of the *Drosophila* nuclear matrix fraction. *Proceedings of the National Academy of Sciences* 82: 4142-4146.

- Beven, A.F., Lee, R., Razaz, M., Leader, D.J., Brown, J.W.S. and Shaw, P.J. (1996) The organization of ribosomal RNA processing correlates with the distribution of nucleolar snRNAs. *Journal of Cell Science* 109:1241-1251.
- Beven, A.F., Guan, Y., Peart, J., Cooper, C. and Shaw, P.J. (1991) Monoclonal antibodies to plant nuclear matrix reveal intermediate filament-related components within the nucleus. *Journal of Cell Science* 98: 293-302.
- Bisotto, S., Lauriault, P., Duval, M. and Vincent, M. (1995) Colocalization of a high molecular mass phosphoprotein of the nuclear matrix (p255) with spliceosomes. *Journal of Cell Science* 108: 1873-1882.
- Blencowe, B.J., Nickerson, J.A., Issner, R., Penman, S. and Sharp, P.A. (1994) Association of nuclear matrix antigens with exon-containing splicing complexes. *Journal of Cell Biology* 127: 593-607.
- Blencowe, B.J., Bauren G., Eldridge A.G., Issner R., Nickerson J.A., Rosonina, E. and Sharp, P.A. (2000) The SRm160/300 splicing coactivator subunits. *RNA* 6:111-120.
- Bode, J., Kohwi, Y., Dickinson, L., Joh, T., Klehr, D., Mielke, C. and Kohwi-Shigematsu, T. (1992) Biological significance of unwinding capability of nuclear matrix-associating DNAs. *Science* 255: 195-197.
- Bode, J., Schlake, T., Rios-Ramirez, M., Meilke, C., Stengert, M., Kay, V., and Klehr-Wirth, D. (1995) Scaffold/ matrix-attached regions: Structural properties creating transcriptionally active loci. *International Review of Cytology* 162A: 389-453.
- Boudonck, K., Dolan, L. and Shaw, P.J. (1998) Coiled body numbers in the Arabidopsis root epidermis are regulated by cell type, developmental stage and cell cycle parameters. *Journal of Cell Science* 111: 3687-3694.
- Bode, J., Stengert-Iber, M., Schlake, T., Kay, V. and Dietz-Pfeilstetter, A. (1996) Scaffold/matrix-attached regions: Topological switches with multiple regulatory functions. *Critical Reviews in Eukaryotic Gene Expression* 6: 115-138.
- Bonifer, C., Vidal, M., Grosveld, F. and Sippel, A. E. (1990) Tissue specific and position independent expression of the complete gene domain for chicken lysozyme in transgenic mice. *EMBO Journal* 9: 2843-2848.
- Bonifer, C., Yannoutsos, N., Kruger, G., Grosveld, F. and Sippel, A. E. (1994) Dissection of the locus control function located on the chicken lysozyme gene domain in transgenic mice. *Nucleic Acids Research* 22: 4202-4210.
- Boulikas, T. (1993) Nature of DNA sequences at the attachment regions of genes to the nuclear matrix. *Journal of Cellular Biochemistry* 52: 14-22.
- Boulikas, T. (1995) Chromatin domains and prediction of MAR sequences. *International Review of Cytology* 162A: 279-388.
- Brancolini, C. and Schneider, C. (1991) Change in the expression of a nuclear matrix-associated protein is correlated with cellular transformation. *Proceedings of the National Academy of Sciences* 88: 6936-6940.

- Breyne, P., Van Montague, M., Depicker, A. and Gheysen, G. (1992) Characterization of a plant scaffold attachment region in a DNA fragment that normalizes transgene expression in tobacco. *Plant Cell* 4: 463-471.
- Breyne, P., Van Montagu, M. and Gheysen, G. (1994) The role of scaffold attachment regions in the structural and functional organization of plant chromatin. *Transgenic Research* 3: 195-202.
- Brown, J.W. S. and Shaw, P.J. (1998) Small nucleolar RNAs and pre-rRNA processing in plants. *Plant Cell* 10: 649-658.
- Capco, D. G., Wan, K. M. and Penman, S. (1982) The nuclear matrix: Three-dimensional architecture and protein composition. *Cell* 29: 847-858.
- Chattopadhyay, S., Whitehurst, C.E. and Chen, J. (1998) A nuclear matrix attachment region upstream of the T cell receptor beta gene binds Cux/CDP and SatB1 and modulates enhancer-dependent reporter gene expression but not endogenous gene expression. *Journal of Biological Chemistry* 273:29838-29846.
- Chattopadhyay, S., Kaul, R., Housman, C.A. and Chen, J.(2000)SMAR1, a novel, alternatively spliced gene product, binds the Scaffold/Matrix-associated region at the T cell receptor beta locus. *Genomics* 68: 93-96.
- Cockerill, P. N. and Garrard, W. T. (1986) Chromosomal loop anchorage of the kappa immunoglobulin gene occurs next to the enhancer in a region containing topoisomerase II sites. *Cell* 44: 273-282.
- Conkling, M. A., Cheng, C.-L., Yamamoto, Y. T. and Goodman, H. M. (1990) Isolation of transcriptionally regulated root-specific genes from tobacco. *Plant Physiology* 93: 1203-1211.
- deBelle, I., Cai, S.T.,KohwiShigematsu, T. (1998) The genomic sequences bound to special AT-rich sequence-binding protein 1 (SATB1) in vivo in Jurkat T cells are tightly associated with the nuclear matrix at the bases of the chromatin loops. *Journal of Cell Biology* 141: 335-348.
- Dickinson, L. A., Joh, T., Kohwi, Y. and Kohwi-Shigematsu, T. (1992) A tissue-specific MAR/SAR DNA-binding protein with unusual binding site recognition. *Cell* 70: 631-645.
- Dickinson, L.A. and Kohwi-Shigematsu,T. (1995) Nucleolin is a MAR/SAR DNA-binding protein specifically recognizing a region with high base-unpairing potential. *Molecular and Cellular Biology* 15: 456-465.
- Dragon, F. and Filipowicz, W. (2000) In vitro assembly of human H/ACA small nucleolar RNPs reveals unique features of U17 and telomerase RNAs. *Molecular and Cellular Biology* 20: 3037-3048.
- Du Q., Melnikova, I.N. and Gardner P.D. (1998) Differential effects of heterogeneous nuclear ribonucleoprotein K on Sp1- and Sp3-mediated transcriptional activation of a neuronal nicotinic acetylcholine receptor promoter. *Journal of Biological Chemistry* 273: 19877-19883.
- Dwarakanath, B.S., Wallen, C.A., St Clair, D.K., Wheeler, K.T. (1991) Characterization of nuclear matrices prepared without salt extraction. *Analytical Biochemistry* 198: 68-74.

- Dworetzky, S. I., Fey, E. G., Penman, S., Lian, J. B., Stein, J. L. and Stein, G. S. (1990) Progressive changes in the protein composition of the nuclear matrix during rat osteoblast differentiation. *Proceedings of the National Academy of Sciences* 87: 4605-4609.
- Earnshaw, W. C., Halligan, B., Cooke, C. A., Heck, M. M. S. and Liu, L.F. (1985) Topoisomerase II in mitotic chromosomes. *Journal of Cell Biology* 100: 1716-1725.
- Eggert M., Michel J., Schneider S., Bornfleth H., Baniahmad A., Fackelmayer F.O., Schmidt S., and Renkawitz R. (1997) The glucocorticoid receptor is associated with the RNA-binding nuclear matrix protein hnRNP U. *Journal of Biological Chemistry* 272: 28471-28478.
- Ellenberg J., Siggia, E.D., Moreira, J.E., Smith, C.L., Presley, J.F., Worman, H.J. and Lippincott-Schwartz, J. (1997) Nuclear membrane dynamics and reassembly in living cells: targeting of an inner nuclear membrane protein in interphase and mitosis. *Journal of Cell Biology* 138: 1193-1206.
- Eversole, A. and Maizels, N. (2000) In vitro properties of the conserved mammalian protein hnRNP D suggest a role in telomere maintenance. *Molecular and Cellular Biology* 20: 5425-5432.
- Fackelmayer, F. O. and Richter, A. (1994a) hnRNP-U/SAF-A is encoded by two differentially polyadenylated mRNAs in human cells. *Biochimica Biophysica Acta* 1217: 232-234.
- Fackelmayer, F. O. and Richter, A. (1994b) Purification of two isoforms of hnRNP-U and characterization of their nucleic acid binding activity. *Biochemistry* 33: 10416-10422.
- Fey, E.G., Krochmalnic, G. and Penman, S. (1986) The non chromatin substructures of the nucleus. The ribonucleoprotein (RNP)-containing and RNA-depleted matrices analyzed by sequential fractionation and resinless section electron microscopy. *Journal of Cell Biology* 102: 1654-1665.
- Fey, E. and Penman, S. (1988) Nuclear matrix proteins reflect cell type of origin in cultured human cells. *Proceedings of the National Academy of Sciences* 85: 121-125.
- Finch, J.T. and Klug A. (1976) Solenoidal model for superstructure in chromatin. *Proceedings of the National Academy of Sciences* 73: 1897-1901.
- Fishel, B.R., Sperry, A.O. and Garrard, W.T. (1993) Yeast calmodulin and a conserved nuclear protein participate in the in vivo binding of a matrix association region. *Proceedings of the National Academy of Sciences* 790: 5623-5627.
- Frederick, S.E., Mangan M.E., Carey J.B., Gruber P.J. (1992) Intermediate filament antigens of 60 and 65 kDa in the nuclear matrix of plants: their detection and localization. *Experimental Cell Research* 199: 213-222.
- Gall J.G., Bellini M., Wu Z. and Murphy C. (1999) Assembly of the nuclear transcription and processing machinery: Cajal bodies (coiled bodies) and transcriptosomes. *Molecular Biology of the Cell* 10: 4385-4402.
- Gasser, S.M. and Laemmli, U.K. (1987) A glimpse at chromosomal order. *Trends in Genetics* 3:16-22.
- Gasser S.M., Amati B.B., Cardenas M.E., Hofmann J.F. (1989) Studies on scaffold attachment sites and their relation to genome function. *International Review of Cytology* 119: 57-96.

- Gerdes, M. G., K. C. Carter, Moen, P.T. and Lawrence, J.B. (1994) Dynamic changes in the higher-level chromatin organization of specific sequences revealed by in situ hybridization to nuclear halos. *Journal of Cellular Biology* 126: 289-304.
- Gerner C., Holzmann K., Grimm R. and Sauer mann G. (1998) Similarity between nuclear matrix proteins of various cells revealed by an improved isolation method. *Journal of Cellular Biochemistry* 71: 363-374.
- Gindullis, F. and Meier, I. (1999) Matrix attachment region binding protein MFP1 is localized in discrete domains at the nuclear envelope. *Plant Cell* 11: 1117-1128.
- Gindullis, F., Peffer, N. J. and Meier, I. (1999) MAF1, a novel plant protein interacting with matrix attachment region binding protein MFP1, is located at the nuclear envelope. *Plant Cell* 11: 1755-1768.
- Ginisty H., Sicard H., Roger B. and Bouvet P. (1999) Structure and functions of nucleolin. *Journal of Cell Science* 112: 761-72.
- Gohring, F., Schwab, B.L., Nicotera, P., Leist, M. and Fackelmayer, F.O. (1997) The novel SAR-binding domain of scaffold attachment factor A (SAF-A) is a target in apoptotic nuclear breakdown. *EMBO Journal* 16: 7361-7371.
- Gonzalez-Melendi P., Beven A., Boudonck, K., Abranches, R., Wells B., Dolan L. and Shaw P. (2000) The nucleus: a highly organized but dynamic structure. *Journal of Microscopy* 198: 199-207.
- Grasser, K.D. (1995) Plant chromosomal high mobility group (HMG) proteins. *Plant Journal* 7: 185-192.
- Gromova, I., Nielsen, O.F. and Razin, S.V. (1995a) Long-range fragmentation of the eukaryotic genome by exogenous and endogenous nucleases proceeds in a specific fashion via preferential DNA cleavage at matrix attachment sites. *Journal of Biological Chemistry* 270: 18685-18690.
- Gromova, I., Thomsen, B. and Razin, S.V. (1995b) Different topoisomerase II antitumor drugs direct similar specific long-range fragmentation of an amplified c-MYC gene locus in living cells and in high-salt-extracted nuclei. *Proceedings of the National Academy of Sciences* 92: 102-106.
- Grondin, B., Bazinet, M. and Aubry, M. (1996) The KRAB zinc finger gene ZNF74 encodes an RNA binding protein tightly associated with the nuclear matrix. *Journal of Biological Chemistry* 271: 15458-15467.
- Gautier, T., Berges, T., Tollrvey, D. and Hurt, E. (1997) Nucleolar KKE/D repeat proteins Nop56p and Nop58p interact with Nop1p and are required for ribosome biogenesis. *Molecular and Cellular Biology* 17: 7088-7098.
- Guo, B., Odgren, P. R., van Wijnen, A. J., Last, T. J., Fey, E. G., Penman, S., Stein, J. L., Lian, J. B. and Stein, G. S. (1995) *Proceedings of the National Academy of Sciences* 92: 10526-10530.
- Hakes, D.J. and Berezney, R. (1991) DNA binding properties of the nuclear matrix and individual nuclear matrix proteins: Evidence for salt resistant DNA binding sites. *Journal of Biological Chemistry* 266: 11131-11140.

- Hall, G., Jr., Allen, G. C., Loer, D. S., Thompson, W. F. and Spiker, S. (1991) Nuclear scaffolds and scaffold-attachment regions in higher plants. *Proceedings of the National Academy of Sciences* 88: 9320-9324.
- Hall, G. E., Jr. and Spiker, S. (1994) Isolation and characterization of nuclear scaffolds. *Plant Molecular Biology Manual*. S. B. Gelvin and R. A. Schilperoort, ed (Belgium: Kluwer Academic Publishers) D1-12.
- Hancock, R., (2000) A new look at the nuclear matrix. *Chromosoma* 109: 219-225.
- Harborth, J., Wang, J., Gueth-Hallonet, C., Weber, K. and Osborn, M.. (1999) Self assembly of NuMA: multiarm oligomers as structural units of a nuclear lattice. *EMBO Journal* 18: 1689-1700.
- Harder, P. A., Silverstein, R. A. and Meier, I. (2000) Conservation of matrix attachment region-binding filament-like protein 1 among higher plants. *Plant Physiology* 122: 225-234.
- Hatton, D. and Gray, J. C. (1999) Two MAR DNA-binding proteins of the pea nuclear matrix identify a new class of DNA-binding proteins. *Plant Journal* 18: 417-429.
- He, D., Nickerson, J.A. and Penman, S. (1990) Core filaments of the nuclear matrix. *Journal of Cell Biology* 110: 569-580.
- Hendzel, M.J., Delcuve G.P. and Davie J.R. (1991) Histone deacetylase is a component of the internal nuclear matrix. *Journal of Biological Chemistry* 266: 21936-21942.
- Henras A., Henry Y., Bousquet-Antonelli C., Noaillac-Depeyre J., Gélugne J-P. and Caizergues-Ferrer M. (1998) Nhp2p and Nop10p are essential for the function of H/ACA snoRNPs. *EMBO Journal* 17:7078-7090.
- Herrscher, R. F., M. H. Kaplan, D. L. Lelsz, C. Das, R. H. Scheuermann and Tucker., P. W. (1995) The immunoglobulin heavy-chain matrix-associating regions are bound by Bright: a B cell-specific trans-activator that describes a new DNA-binding protein family. *Genes and Development* 9: 3067-3082
- Hibino, Y., Tsukada, S. and Sugan0, N. (1993) Properties of a DNA-binding protein from rat nuclear scaffold fraction. *Biochemical and Biophysical Research Communications* 197: 336-342.
- Holzmann, K. Gerner, c., Korosec, T., Poltl, A., Grimm, R. and Sauer mann, G. (1998) Identification and characterization of the ubiquitously occurring nuclear matrix protein NMP 238. *Biochemical and Biophysical Research Communications* 252: 39-45.
- Hong Z., Mann, P., Brown, N.H., Tran, L.E., Shaw, K.J., Hare, R.S. and DiDomenico, B. (1994) Cloning and characterization of KNR4, a yeast gene involved in (1,3)-beta-glucan synthesis. *Molecular and Cellular Biology* 14: 1017-1025.
- Ingram, J.L. (2000) High Mobility Group I/Y type proteins in *Nicotiana tabacum* and their interactions with matrix attachment regions and the nuclear matrix. (Doctoral dissertation) North Carolina State University Department of Genetics, Raleigh, NC.

- Irie, S. and Sezaki, M. (1983) A quantitative determination of the relative amount of histones in polyacrylamide gel by silver stain. *Analytical Biochemistry* 134: 471-478.
- Isaac, C., Yang, Y. and Meier, U.T. (1998) Nopp140 functions as a molecular link between the nucleolus and the coiled bodies. *Journal of Cell Biology* 142: 319-329.
- Ivanchenko, M. and Avramova, Z. (1992) Interaction of MAR-sequences with nuclear matrix proteins. *Journal of Cellular Biochemistry* 50: 190-200.
- Ivanchenko M., Tasheva, B., Stoilov, L., Christova, R. and Zlatanova, J. (1993) Characterization of some nuclear matrix proteins in maize. *Plant Science* 91:35-43.
- Izuarralde, E., Käs, E. and Laemmli U.K. (1989) Highly preferential nucleation of histone H1 assembly on scaffold-associated regions. *Journal of Molecular Biology* 210: 573-85.
- Jackson, D.A. Dolle, A., Robertson, G. and Cook, P.R. (1992) The attachments of chromatin loops to the nucleoskeleton. *Cell Biology International Reports* 16: 687-696.
- Jady B.E. and Kiss T. (2001) A small nucleolar guide RNA functions both in 2'-O-ribose methylation and pseudouridylation of the U5 spliceosomal RNA. *EMBO Journal* 20: 541-551.
- Johnson, B. F., Marciniak, R.A. and Guarente, L. (1998) Telomeres, the nucleolus and aging. *Current Opinion in Cell Biology* 10: 332-338.
- Kallajoki M., Weber K. and Osborn M. (1991) A 210 kDa nuclear matrix protein is a functional part of the mitotic spindle; a microinjection study using SPN monoclonal antibodies. *EMBO Journal* 10: 3351-3362.
- Kanemaki, M., Kurokawa, Y., Matsuura, T., Makino, Y., Masani, A., Okazaki, K., Morishita, T. and Tamura, T. (1999) TIP49b, a new RuvB-like DNA helicase, is included in a complex together with another RuvB-like helicase. *Journal of Biological Chemistry* 274: 22437-22444.
- Kanemaki, M., Makino, Y., Yoshida, T., Kishimoto, T., Koga, A., Yamamoto, K., Yamamoto, M., Moncollin, V., Egly, J.-M., Muramatsu, M. and Tamura, T. (1997) Molecular cloning of a rat 49kDa TBP-interacting protein (TIP49) that is highly homologous to the bacterial RuvB. *Biochemical and Biophysical Research Communications* 235: 64-68.
- Käs, E., Poljak, L., Adachi, Y. and Laemmli, U.K. (1993) A model for chromatin opening: stimulation of topoisomerase II and restriction enzyme cleavage of chromatin by distamycin. *EMBO Journal* 12: 115-126.
- Kay, V. and Bode, J., (1994) Binding specificity of a nuclear scaffold: supercoiled, single-stranded and scaffold-attached DNA. *Biochemistry* 33: 367-374.
- Kiledjian, M. and Dreyfuss, G. (1992) Primary structure and binding activity of the hnRNP U protein: binding RNA through RGG box. *EMBO Journal* 11: 2655-2664.
- Kim M.K., Lesoon-Wood L.A., Weintraub B.D. and Chung J.H. (1996) A soluble transcription factor, Oct-1, is also found in the insoluble nuclear matrix and possesses silencing activity in its alanine-rich domain. *Molecular and Cell Biology* 16: 4366-4377.

- Kohwi-Shigematsu, T., Maass, K. and Bode, J. (1997) A thymocyte factor SATB1 suppresses transcription of stably integrated MAR-linked reporter genes. *Biochemistry* 36: 12005-12010.
- Kramer J. A., Singh, G. B. and Krawetz, S. A. (1996) Computer-assisted search for sites of nuclear matrix attachment. *Genomics* 33: 305-308.
- Krauss, S.W., Chasis, J.A., Rogers, C., Mohandas, N., Krockmalnic, G and Penman S. (1997) Structural protein 4.1 is located in mammalian centrosomes. *Proceedings of the National Academy of Sciences* 94: 7297-7302.
- Krecic A.M., Swanson M.S. (1999) hnRNP complexes: composition, structure, and function. *Current Opinion in Cell Biology* 11: 363-371.
- Lafontaine, D.L.J., Bousquet-Antonelli, C., Henry, Y., Caizergues-Ferrer, M. and Tollervey D. (1998) The box H+ACA snoRNAs carry Cbf5p, the putative rRNA pseudouridine synthase. *Genes and Development* 12: 527-537.
- Lafontaine, D.L. and Tollervey, D. (1999) Nop58p is a common component of the box C+D snoRNPs that is required for snoRNA stability. *RNA* 5:455-467.
- Lafontaine, D.L. and Tollervey, D. (2000) Synthesis and assembly of the box C+D small nucleolar RNPs. *Molecular and Cellular Biology* 20: 2650-2659.
- Lamond, A.I. and Earnshaw, W.C. (1998) Structure and function in the nucleus. *Science* 280:547-553.
- Lawrence, J.B., Singer, R.H., Marselle, L.M. (1989) Highly localized tracks of specific transcripts within interphase nuclei visualized by in situ hybridization. *Cell* 57: 493-502.
- Leader, D.J., Sanders, J.F., Waugh, R., Shaw, P.J. and Brown, J.W.S. (1994) Molecular characterization of plant U14 small nucleolar RNA genes: Closely linked genes are transcribed as a polycistronic U14 transcript. *Nucleic Acids Research* 22: 5196-5200.
- Leader, D.J., Clark, G.P., Watters, J., Beven, A.F., Shaw, P.J. and Brown, J.W.S. (1997) Clusters of multiple different small nucleolar RNA genes in plants are expressed as and processed from polycistronic pre-snoRNAs. *EMBO Journal* 16: 5742-5751.
- Leader, D.J., Clark, G.P., Watters, J., Beven, A.F., Shaw, P.J. and Brown, J.W.S. (1999) Splicing-independent processing of plant box C/D and box H/ACA small nucleolar RNAs. *Plant Molecular Biology* 39: 1091-1100.
- Leuba, S.H., Yang, G., Robert, C., Samori, B., van Holde, K., Zlatanova, J. and Bustamante, C. (1994) Three-dimensional structure of extended chromatin fibers as revealed by tapping-mode scanning force microscopy. *Proceedings of the National Academy of Sciences* 91: 11621-11625.
- Lewis, J.D. and Tollervey, D. (2000) Like attracts like: getting RNA processing together in the nucleus. *Science* 288: 1385-1389.
- Liu, J., D. Bramblett, Q. Zhu, M. Lozano, R. Kobayashi, S. R. Ross and J. P. Dudley. (1997) The matrix attachment region-binding protein SATB1 participates in negative regulation of tissue-specific gene expression. *Molecular and Cellular Biology* 17:5275-5287.

- Lodish, H., Berk, A., Matsudaira, P., Baltimore, D., Zipursky, L. and Darnell, J. (1996) *Molecular Cell Biology*. (New York, New York: W. H. Freeman and Company).
- Lowe, T.M. and Eddy, S.R. (1999) A computational screen for methylation guide snoRNAs in yeast. *Science* 283: 1168-1171.
- Luderus, M.E.E., den Blaauwen, J.L., de Smit, O.J.B., Compton, D.A. and van Driel, R. (1992) Binding of matrix attachment regions to lamin B1. *Cell* 70: 949-959.
- Luderus, M.E.E., de Graaf, A., Mattia, E., den Blaauwen, J.L. Grande, M.A., de Jong, L. and van Driel, R. (1994) Binding of matrix attachment regions to lamin polymers involves single-stranded regions and the minor groove. *Molecular and Cellular Biology* 14: 6297-6305.
- Marshall, W.F. and Sedat, J.W. (1997) Deconstructing the nucleus: global architecture from local interactions. *Current Opinions in Genetics and Development*. 1997 7: 259-263.
- Matakatsu H., Tadokoro R., Gamo S. and Hayashi S. (1999) Repression of the wing vein development in *Drosophila* by the nuclear matrix protein plexus. *Development* 126: 5207-5216.
- Mathews, C.K. and van Holde, K.E. (1990) *Biochemistry*. The Benjamin/Cummings Publishing Company, Inc. Redwood City, CA.
- Mattern K.A., Humbel B.M., Muijsers A.O., de Jong L. and van Driel R. (1996) hnRNP proteins and B23 are the major proteins of the internal nuclear matrix of HeLa S3 cells. *Journal of Cellular Biochemistry* 62: 275-289.
- Mattern, K.A., van der Kraan, I., Schul, W., de Jong, L. and van Driel, R. (1999) Spatial organization of four hnRNP proteins in relation to sites of transcription, to nuclear speckles, and to each other in interphase nuclei and nuclear matrices of HeLa cells. *Experimental Cell Research* 246: 461-470.
- Mattern, K.A., van Goethem, R.E., de Jong, L. and van Driel, R. (1997) Major internal nuclear matrix proteins are common to different human cell types. *Journal of Cellular Biochemistry* 65: 42-52.
- McNulty A.K. and Saunders MJ. (1992) Purification and immunological detection of pea nuclear intermediate filaments: evidence for plant nuclear lamins. *Journal of Cell Science* 103 : 407-414.
- Meier, I., Phelan, T., Gruissem, W., Spiker, S. and Schneider, D. (1996) MFP1, a novel plant filament-like protein with affinity for matrix attachment region DNA. *Plant Cell* 8: 2105-2115.
- Meier, U.T. and Blobel, G. (1994) NAP57, a mammalian nucleolar protein with a putative homolog in yeast and bacteria. *Journal of Cell Biology* 127: 1505-1514.
- Meissner, M., Dechat, T., Gerner, C., Grimm, R., Foisner, R. and Saueremann, G. (2000) Differential nuclear localization and nuclear matrix association of the splicing factors PSF and PTB. *Journal of Cellular Biochemistry* 76: 559-66.
- Merriman, H.L., van Wijnen, A.J., Hiebert, S., Bidwell, J.P., Fey, E., Lian, J., Stein, J. and Stein, G.S. (1995) The tissue-specific nuclear matrix protein, NMP-2, is a member of the AML/CBF/PEBP2/runt domain transcription factor family: interactions with the osteocalcin gene promoter. *Biochemistry* 34: 13125-13132.

- Miau, L.H., Chang, C.J., Shen, B.J., Tsai, W.H. and Lee, S.C. (1998) Identification of heterogeneous nuclear ribonucleoprotein K (hnRNP K) as a repressor of C/EBPbeta-mediated gene activation. *Journal of Biological Chemistry* 273: 10784-10791.
- Michalowski, S. M., Allen, G. C., Hall, Jr., G. E., Thompson, W. F. and Spiker, S. (1999) Characterization of randomly-obtained matrix attachment regions (MARs) from higher plants. *Biochemistry* 38 12795-12804.
- Michelotti, E.F., Michelotti, G.A., Aronsohn, A.I. and Levens, D. (1996) Heterogeneous nuclear ribonucleoprotein K is a transcription factor. *Molecular and Cellular Biology* 16: 2350-2360.
- Mielke, C., Kohwi, Y., Kohwi-Shigematsu, T. and Bode, J. (1990) Hierarchical binding of DNA fragments derived from scaffold-attached regions: Correlation of properties *in vitro* and function *in vivo*. *Biochemistry* 29: 7475-7485.
- Minguez, A. and Moreon Diaz de la Espina, S. (1993) Immunological characterization of lamins in the nuclear matrix of onion cells. *Journal of Cell Science* 106: 431-439.
- Mirkovitch, J., Mirault, M.-E. and Laemmli, U. (1984) Organisation of the higher-order chromatin loop: specific DNA attachment sites on nuclear scaffold. *Cell* 39: 223-232.
- Mirkovitch, J., Spiere, P. and Laemmli, U. K. (1986) Genes and loops in 320,000 base-pairs of the *Drosophila melanogaster* chromosome. *Journal of Molecular Biology* 190: 255-258.
- Miskimins, W.K., Roberts, M.P., McClelland, A. and Ruddle, F.H. (1985) Use of a protein-blotting procedure and a specific DNA probe to identify nuclear proteins that recognize the promoter region of the transferrin receptor gene. *Proceedings of the National Academy of Sciences* 82: 6741-6744.
- Moir, R.D., Spann, T.P. and Goldman, R.D. (1995) The dynamic properties and possible functions of nuclear lamins. *International Review of Cytology* 162B: 141-182.
- Moreno Diaz de la Espina, S. M. (1995) Nuclear matrix isolated from plant cells. *International Review of Cytology* 162B, 75-139.
- Morin, P.J., Downs, J.A., Snodgrass, A.M. and Gilmore, T.D. (1995) Genetic analysis of growth inhibition by GAL4-L kappa B-alpha in *Saccharomyces cerevisiae*. *Cell Growth and Differentiation* 6: 789-798.
- Morisawa, G., Han-yama, A., Moda, I., Tamai, A., Iwabuchi, M. and Meshi, T. (2000) AHM1, a novel type of nuclear matrix-localized, MAR-binding protein with a single AT hook and a J domain-homologous region. *Plant Cell* 12: 1903-1916.
- Mortillaro, M.J., Blencowe, B.J., Wei, X.Y., Nakayasu, H., Du, L., Warren, S.L., Sharp, P.A. and Berezney, R. (1996) A hyperphosphorylated form of the large subunit of RNA polymerase II is associated with splicing complexes and the nuclear matrix. *Proceedings of the National Academy of Sciences* 93: 8253-8257.
- Nan, X.; Tate, P.; Li, E. and Bird, A. (1996) DNA methylation specifies chromosomal localization of MeCP2. *Molecular and Cellular Biology* 16: 414-421.

- Nayler O., Strätling W., Bourquin J.P., Stagljar I., Lindemann L., Jasper H., Hartmann A.M., Fackelmayer F.O., Ullrich A. and Stamm S. (1998) SAF-B protein couples transcription and pre-mRNA splicing to SAR/MAR elements. *Nucleic Acids Research* 26: 3542-3549.
- Neri L.M., Fackelmayer F.O., Zweyer M., Kohwi-Shigematsu T. and Martelli A.M. (1997) Subnuclear localization of S/MAR-binding proteins is differently affected by in vitro stabilization with heat or Cu²⁺. *Chromosoma* 106: 81-93.
- Newberry E.P., Latifi T. and Towler D.A. (1999) The RRM domain of MINT, a novel Msx2 binding protein, recognizes and regulates the rat osteocalcin promoter. *Biochemistry* 38: 10678-10690.
- Newman, D.R., Kuhn, J.F., Shanab, G.M. and Maxwell, E.S. (2000) Box C/D snoRNA-associated proteins: Two pairs of evolutionary ancient proteins and possible links to replication and transcription. *RNA* 6: 861-879.
- Nickerson, J.A., Blencowe, B. and Penman, S. (1995) The architectural organization of nuclear metabolism. *International Review of Cytology* 162A: 67-123.
- Nickerson, J.A., Krockmalnic, G., Wan, K. and Penman, S. (1997) The nuclear matrix revealed by eluting chromatin from a cross-linked nucleus. *Proceedings of the National Academy of Sciences* 94: 4446-4450.
- Oesterreich, S.; Lee, A. V.; Sullivan, T. M.; Samuel, S. K.; Davie, J. R. and Fuqua, S. A. W. (1997) Novel nuclear matrix protein HET binds to and influences activity of the HSP27 promoter in human breast cancer cells. *Journal of Cellular Biochemistry* 67: 275-286.
- Olson, M.O.J., Dundr, M. and Szebeni, A. (2000) The nucleolus: an old factory with unexpected capabilities. *Trends in Cell Biology* 10: 189-196.
- Omer, A., Lowe, T.M., Russell, A.G., Ehardt, H., Eddy, S.R. and Dennis, P.P. (2000) Homologs of small nucleolar RNAs in Archaea. *Science* 288: 517-22.
- Patturajan, M., Wei, X., Berezney, R. and Corden, J.L. (1998) A nuclear matrix protein interacts with the phosphorylated C-terminal domain of RNA polymerase II. *Molecular and Cellular Biology* 18: 2406-2415.
- Paul, A. L. and Ferl, R. J. (1998) Higher order chromatin structures in maize and Arabidopsis. *Plant Cell* 10: 1349-1359.
- Paulson, J. R. and Laemmli, U. K. (1977) The structure of histone-depleted metaphase chromosomes. *Cell* 12: 817-828.
- Pedersen, T.J., Arwood, L.J., Spiker, S., Gultinan, M.J. and Thompson, W.F. (1991) High mobility group chromosomal proteins bind to AT-rich tracts flanking plant genes. *Plant Molecular Biology* 16: 95-104.
- Pederson, T. (1998) Thinking about a nuclear matrix. *Journal of Molecular Biology* 277: 147-159.
- Pederson, T. (2000) Half a century of the "nuclear matrix". *Molecular Biology of the Cell* 11: 799-805.

- Perrin, L., Demakova, O., Fanti, L., Kallenbach, S., Saingery, S., Mal'ceva, N. I., Pimpinelli, S., Zhimulev, I. and Pradel, J. (1998) Dynamics of the sub-nuclear distribution of Modulo and the regulation of position-effect variegation by nucleolus in *Drosophila*. *Journal of Cell Science* 111: 2753-2761.
- Phi-Van, L. and Strätling, W. H. (1996) Dissection of the ability of the chicken lysozyme gene 5' matrix attachment region to stimulate transgene expression and to dampen position effects. *Biochemistry* 35: 10735-10742.
- Phi-Van, L., von Kries, J. P., Ostertag, W. and Strätling, W. H. (1990) The chicken lysozyme 5' matrix attachment region increases transcription from a heterologous promoter in heterologous cells and dampens position effects on the expression of transfected genes. *Molecular and Cellular Biology* 10: 2302-2307.
- Pikaard, C. S. (1998) Chromosome topology - Organizing genes by loops and bounds. *Plant Cell* 10: 1229-1232.
- Poljak, L., Seum, C., Mattioni, T. and Laemmli, U. K. (1994) SARs stimulate but do not confer position independent gene expression. *Nucleic Acids Research* 22: 4386-4394.
- Pruss, D., Bartholomew, B., Persinger, J., Hayes, J., Arents, G., Moudrianakis, E. N. and Wolffe, A. P. (1996) An asymmetric model for the nucleosome: a binding site for linker histones inside the DNA gyres. *Science* 274: 614-617.
- Rando, O.J., Zhao, K. and Crabtree, G.R. (2000) Searching for a function for nuclear actin. *Trends in Cell Biology* 10: 92-97.
- Razin, S.V., Hancock, R., Iarovaia, O., Westergaard, O., Gromova, I. and Georgiev, G.P. (1993) Structural-functional organization of chromosomal DNA domains. *Cold Spring Harbor Symposia on Quantitative Biology* 58: 25-35.
- Renz, A. and Fackelmayer, F. O. (1996) Purification and molecular cloning of the scaffold attachment factor B (SAF-B), a novel human nuclear protein that specifically binds to S/MAR-DNA. *Nucleic Acids Research* 24: 843-849.
- Reyes, J.C., Muchardt, C. and Yaniv, M. (1997) Components of the human SWI/SNF complex are enriched in active chromatin and are associated with the nuclear matrix. *Journal of Cell Biology* 137: 263-74.
- Roder, K., Hung, M.-S., Lee, T.-L., Lin, T.-Y., Xiao, H., Isobe, K.-I., Juang, J.-L. and Shen, C.-K. J. (2000) transcriptional repression by *Drosophila* methyl-CpG-binding proteins. *Molecular and Cellular Biology* 20: 7401-7409.
- Romig, H.; Fackelmayer, F. O.; Renz, A.; Ramsperger, U. and Richter, A. (1992) Characterization of SAF-A, a novel nuclear DNA binding protein from HeLa cells with high affinity for nuclear matrix/scaffold attachment DNA elements. *EMBO Journal* 11: 3431-3440.
- Rzepecki, R., Markiewicz, E., Adamiec, R. and Szopa, J. (1995a) Interaction of the *Pisium sativum* nuclear matrix proteins with SAR DNA. *Acta Biochimica Polonica* 42: 75-81.

- Rzepecki, R., Markiewicz, E., Adamiec, R. and Szopa, J. (1995b) Identification of the proteins responsible for SAR DNA binding in nuclear matrix of *Cucurbita pepa*. *Acta Biochimica Polonica* 42: 171-176.
- Saitoh, N., Gldberg, I. and Earnshaw, W.C. (1995) The SMC proteins and the coming of age of the chromosome scaffold hypothesis. *Bioessays* 17: 759-766.
- Sambrook, J., E. F. Fritsch, and Maniatis, T. (1989) *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Samuel, S.K., Spencer, V.A., Bajno, L., Sun, J.M., Holth, L.T., Oesterreich, S. and Davie, J.R. (1998) In situ cross-linking by cisplatin of nuclear matrix-bound transcription factors to nuclear DNA of human breast cancer cells. *Cancer Research* 58: 3004-3008.
- Sanvito F., Piatti S., Villa A., Bossi M., Lucchini G., Marchisio P.C. and Biffo S. (1999) The beta4 integrin interactor p27(BBP/eIF6) is an essential nuclear matrix protein involved in 60S ribosomal subunit assembly. *Journal of Cell Biology* 144: 823-837.
- Singh, G.B., Kramer, J.A. and Krawetz, S.A. (1997) Mathematical model to predict regions of chromatin attachment to the nuclear matrix. *Nucleic Acids Research*: 25: 1419-1425.
- Sleeman, J., Lyon, C.E., Platani, M., Kreivi, J.-P. and Lamond, A.I. (1998) Dynamic interactions between splicing snRNPs, coiled bodies and nucleoli revealed using snRNP protein fusions to the green fluorescent protein. *Experimental Cell Research* 243: 290-304.
- Smith, C.M. and Steitz, J.A. (1997) Sno storm in the nucleolus: New role for myriad small RNPs. *Cell* 89: 669-672.
- Spector, D. L. (1990) Higher order nuclear organization: three-dimensional distribution of small nuclear ribonucleoprotein particles. *Proceedings of the National Academy of Sciences* 87: 147-151.
- Spector, D. L. (1994) RNA processing: Cycling splicing factors. *Nature* 23: 369: 604.
- Spiker, S. and Thompson, W. F. (1996) Nuclear matrix attachment regions and transgene expression in plants. *Plant Physiology* 110: 15-21.
- Stief, A., Winter, D. M., Strätling, W. H. and Sippel, A. E. (1989) A nuclear DNA attachment element mediates elevated and position-independent gene activity. *Nature* 341: 343-345.
- Stuurman, N., Meijne, A. M. L., van der Pol, A. J., de Jong, L., van Driel, R. and van Renswoude, J. (1990) The nuclear matrix from cells of different origin. *Journal of Biological Chemistry* 265: 5460-5465.
- Szekeres, M., Haizel, T., Adam, E. and Nagy, F. (1995) Molecular characterization and expression of a tobacco histone H1 cDNA. *Plant Molecular Biology* 27: 597-605.
- Tan J.H., Wooley J.C. and LeStourgeon, W.M. (2000) Nuclear matrix-like filaments and fibrogranular complexes form through the rearrangement of specific nuclear ribonucleoproteins. *Molecular Biology of the Cell* 11: 1547-1554.

- Tatusova, T. A. and Madden, T.L. (1999) Blast 2 sequences - a new tool for comparing protein and nucleotide sequences. *FEMS Microbiology Letters* 174: 247-250.
- The Arabidopsis Genome Initiative. (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815.
- Trubiani O., De Fazio P., Pieri C., Mazzanti L. and Di Primio R. (2000) Nuclear matrix provides linkage sites for translocated NF-kappa B: morphological evidence. *Histochemistry and Cell Biology* 113: 369-377.
- Tsutsui, K., Tsutsui, K., Okada, S., Watarai, s., Seki, S., Yasuda, T. and Shohmor, T. (1993) Identification and characterization of a nuclear scaffold protein that binds the matrix attachment region DNA. *Journal of Biological Chemistry* 268: 12886-12894.
- Tuteja, R. and Tuteja N., (1999) Nucleolin: a multifunctional major nucleolar phosphoprotein. *Critical Reviews in Biochemistry and Molecular Biology* 33: 407-436.
- Tycowski K.T., You Z.H., Graham P.J. and Steitz J.A. (1998) Modification of U6 spliceosomal RNA is guided by other small RNAs. *Molecular Cell*. 2: 1-20.
- Ülker, B., Allen, G. C., Thompson, W. F., Spiker, S. and Weissinger, A. K. (1999) A tobacco matrix attachment region reduces the loss of transgene expression in the progeny of transgenic tobacco plants. *Plant Journal* 18: 253-263.
- Van Driel, R. and Otte, A. P., (1997) *Nuclear Organization, Chromatin Structure, and Gene Expression*. Oxford University Press, Inc. New York.
- Van Holde, K. E. (1989) *Chromatin*. Springer-Verlag New York.
- Venema J. and Tollervey D. (1999) Ribosome synthesis in *Saccharomyces cerevisiae*. *Annual Reviews in Genetics*. 33: 261-311.
- Verheijen, R., van Venrooij, W. and Ramaekers, F. (1988) The nuclear matrix: structure and composition. *Journal of Cell Science* 90: 11-36.
- Visintin R. and Amon A. (2000) The nucleolus: the magician's hat for cell cycle tricks. *Current Opinion in Cell Biology* 12: 372-377.
- Vishwanatha, J.K., Jindal, H.K. and Davis, R.G. (1992) The role of primer reinitiation proteins in DNA-replication-association with nuclear matrix in HELA-cells. *Journal of Cell Science* 101: 25-34.
- von Kries, J. P., Buhrmester, H. and Strätling, W. H. (1991) A matrix/scaffold attachment region binding protein: Identification, purification and mode of binding. *Cell* 64: 123-135.
- von Kries, J. P., Buck, F. and Strätling, W. H. (1994) Chicken MAR binding protein p120 is identical to human heterogeneous nuclear ribonucleoprotein (hnRNP) U. *Nucleic Acids Research* 22: 1215-1220.
- Wan, K. M., Nickerson, J. A., Krockmalnic, G. and Penman, S. (1999) The nuclear matrix prepared by amine modification. *Proceedings of the National Academy of Sciences* 96: 933-938.

- Wang, Z., Goldstein, A., Zong, R.-T., Lin, D., Neufeld, E. J., Scheuermann, R. H. and Tucker, P. W. (1999) Cux/CDP homeoprotein is a component of NF- μ NR and represses the immunoglobulin heavy chain intronic enhancer by antagonizing the bright transcription activator. *Molecular and Cellular Biology* 19: 284-295.
- Watkins, N.J., Leverette, R.D., Xia, L., Andrews, M.T. and Maxwell, E.S. (1996) Elements essential for processing intronic U14 snoRNA are located at the termini of the mature snoRNA sequence and include conserved nucleotide boxes C and D. *RNA* 2: 118-133.
- Watkins N.J., Segault V., Charpentier B., Nottrott S., Fabrizio P., Bachi A., Wilm M., Rosbash M., Branlant C. and Luhrmann R. (2000) A common core RNP structure shared between the small nucleolar box C/D RNPs and the spliceosomal U4 snRNP. *Cell* 103: 457-466.
- Watkins, N.J., Newman, D.R., Kuhn, J.F. and Maxwell, E.S. (1998) In vitro assembly of the mouse U14 snoRNP core complex and identification of a 65-kDa box C/D-binding protein. *RNA* 4: 582-593.
- Weighardt, F., Cobianchi, F., Cartegni, L., Chiodi, I., Villa, A., Riva, S. and Biamonti, G. (1999) A novel hnRNP protein (HAP/SAF-B) enters a subset of hnRNP complexes and relocates in nuclear granules in response to heat shock. *Journal of Cell Science* 112: 1465-1476.
- Weipoltshammer, K., C. Schofer, Almeder, M., Philimonenko, V.V., Frei, K., Wachtler, F. and Hozak, P. (1999) Intranuclear anchoring of repetitive DNA sequences: centromeres, telomeres, and ribosomal DNA. *Journal of Cell Biology* 147: 1409-1418.
- Weitzel, J.M., Buhrmester, H. and Strätling, W.H. (1997) Chicken MAR-binding protein ARBP is homologous to rat methyl-CpG-binding protein MeCP2. *Molecular and Cellular Biology* 17: 5656-5666.
- Weitzel, J. and Strätling, W.H. (1997) The case of chicken attachment region binding protein reinforces the importance of pre-screening of several libraries for cDNA cloning. *Biochemical Journal* 321: 261-264.
- Widom, J. (1997) Chromatin: the nucleosome unwrapped. *Current Biology* 7: R653-R655.
- Wu, P., Brockenbrough, S., Metcalfe, A.C., Chen, S. and Aris, J. (1998) Nop5p is a small nucleolar ribonucleoprotein component required for pre-18S rRNA processing in yeast. *Journal of Biological Chemistry* 273: 16453-16463.
- Xing, Y.G. and Lawrence, J.B. (1991) Preservation of specific RNA distribution within the chromatin-depleted nuclear substructure demonstrated by in situ hybridization coupled with biochemical fractionation. *Journal of Cell Biology* 112: 1055-1063.
- Yang, Y., Isaac, C., Wang, C., Dragon, F., Pogacic, V. and Meier, U.T. (2000) Conserved composition of mammalian box H/ACA and box C/D small nucleolar ribonucleoprotein particles and their interaction with the common factor Nopp140. *Molecular Biology of the Cell* 11: 567-577.
- Yu F., Thiesen J., Strätling, W.H. (2000) Histone deacetylase-independent transcriptional repression by methyl-CpG-binding protein 2. *Nucleic Acids Research* 28: 2201-2206.

- Zeng, C., McNeil, S., Pockwinse, S., Nickerson, J., Shopland, L., Lawrence, J.B., Penman, S., Hiebert, S., Lian, J.B., van Wijnen, A.J., Stein, J.L. and Stein G.S. (1998) Intranuclear targeting of AML/CBF regulatory factors to nuclear matrix-associated transcriptional domains. *Proceedings of the National Academy of Sciences* 95: 1585-1589.
- Zhao, J. and Grafi, G. (2000) The high mobility group I/Y protein is hypophosphorylated in endoreduplicating maize endosperm cells and is involved in alleviating histone H1-mediated transcriptional repression. *Journal of Biological Chemistry* 275: 27494-27499.
- Zhao, K., Käs, E., Gonzalez, E. and Laemmli, U.K. (1993) SAR dependent mobilization of histone H1 by HMG I/Y in vitro: HMG I/Y is enriched in H1-depleted chromatin. *EMBO Journal* 12: 3237-3247.
- Zhelev N.Z., Todorov I.T, Philipova R.N., Hadjiolov A.A. (1990) Phosphorylation-related accumulation of the 125K nuclear matrix protein mitotin in human mitotic cells. *Journal of Cell Science* 95: 59-64.
- Zini, N., Santi, S., Ognibene, A., Bavelloni, A., Neri, L.M., Valmori, A., Mariani, E., Negri, C., Astaldi-Ricotti, G.C. and Maraldi N.M. (1994) Discrete localization of different DNA topoisomerases in HeLa and K562 cell nuclei and subnuclear fractions. *Experimental Cell Research* 10: 336-348.