

REFERENCES

REFERENCES

- [1] American Heart Association, *Arrhythmias and Sudden Cardiac Death*. WWW Site, <http://gauss.technion.ac.il/~rl/Gilead/scd>, 1997.
- [2] M.J. Janse, F. J. G. Wilmsschopman, and R. Coronel, "Ventricular-fibrillation is not always due to multiple wavelet reentry," *J. Cardiovasc. Electrophysiol.*, vol. 6, pp. 512-521, July 1995.
- [3] J. M. Rogers, P. V. Bayly, R. E. Ideker, and W. M. Smith, "Quantitative techniques for analyzing high-resolution cardiac mapping data," *IEEE Eng. in Med. and Biol. Magazine*, vol. 17, pp. 62-72, Jan. - Feb. 1998.
- [4] M. S. Spach, W. T. Miller, E. Miller-Jones, R. B. Warren, and R. C. Barr, "Extracellular potentials related to intracellular action potentials during impulse conduction in anisotropic canine cardiac muscle," *Circ. Res.*, vol. 45, pp. 188-204, Aug. 1979.
- [5] T. Lewis and M. A. Rothschild, "The excitatory process in the dog's heart. Part II. The ventricles," *Phil. Trans. Roy. Soc.*, vol. 206, pp. 181-226, 1915.
- [6] F. X. Witkowski, R. Plonsey, P. A. Penkoske, and K. M. Kavanagh, "Significance of inwardly directed transmembrane current in determination of local myocardial electrical activation during ventricular fibrillation," *Circ. Res.*, vol. 74, pp. 507-524, March 1994.
- [7] S. M. Blanchard, A. S. Tohmaz, D. J. Cervino, T. S. Pool, B. P. Clark, E. E. Johnson, and R. E. Ideker, "Detecting activations in unipolar epicardial electrograms recorded during ventricular fibrillation," in *Proc. of Computers in Cardiol.*, pp. 793-796, 1995.
- [8] C. F. Pieper and A. Pacifico, "Improved spatial resolution for cardiac mapping using current source density-based electrode arrays," *PACE* vol. 18, pp. 34-44, Jan. 1995.
- [9] D. B. Geselowitz, P. E. Crombie, S. M. Smith, and E. J. Berbari, "Use of surface Laplacian to extract local activation from epicardial electrograms," in *Proc. of Computers in Cardiol.*, pp. 129-132, 1996.
- [10] D. Itchhaporia, P. B. Snow, R. J. Almassy, and W. J. Oetgen, "Artificial neural networks: current status in cardiovascular medicine," *J. Am. Col. Cardiol.*, vol. 28, pp. 515-521, Aug. 1996.

- [11] F. M. Ham and S. Han, "Classification of cardiac arrhythmias using fuzzy ARTMAP," *IEEE Trans. Biomed. Eng.*, vol. 43, pp. 425-430, April 1996.
- [12] W. G. Baxt, "Use of an artificial neural network for the diagnosis of myocardial infarction," *Ann. Intern. Med.*, vol. 115, pp. 843-848, Dec. 1991.
- [13] B. Heden, H. Ohlin, R. Rittner, and L. Edenbrandt, "Acute myocardial infarction detected in the 12-lead ECG by artificial neural networks," *Circulation*, vol. 96, pp. 1798-1802, Sept. 1997.
- [14] R. L. Kennedy, R. F. Harrison, A. M. Burton, H. S. Fraser, W. G. Hamer, D. MacArthur, R. McAllum, and D. J. Steedman, "An artificial neural network system for diagnosis of acute myocardial infarction (AMI) in the accident and emergency department: evaluation and comparison with serum myoglobin measurements," *Computer Methods and Programs in Biomedicine*, vol. 52, pp. 93-103, Feb. 1997.
- [15] S. Barro, M. Fernandez-Delgado, J. A. Vila-Sobrino, C. V. Regueiro, and E. Sanchez, "Classifying multichannel ECG patterns with an adaptive neural network," *IEEE Eng. Med. and Biol.*, vol. 17, pp. 45-55, Jan. - Feb. 1998.
- [16] R. H. Clayton, A. Murray, and R. W. F. Campbell, "Recognition of ventricular fibrillation using neural networks," *Med. & Biol. Eng. & Comput.*, vol. 32, pp. 217-220, March 1994.
- [17] W. R. Dassen, V. L. Karthaus, J. L. Talmon, R. G. Mulleneers, J. L. Smeets, and H. J. Wellens, "Evaluation of new self-learning techniques for the generation of criteria for differentiation of wide-QRS tachycardia in supraventricular tachycardia and ventricular tachycardia," *Clin. Cardiol.*, vol. 18, pp. 103-108, Feb. 1995.
- [18] M. Jabri and E. Tinker, "Classifier architectures for single chamber arrhythmia recognition," *Applied Intelligence*, vol. 6, pp. 215-224, July 1996.
- [19] P. Werbos, *Beyond Regression: New Tools for Prediction and Analysis in the Behavioral Sciences*. Cambridge, MA: Harvard University Department of Applied Mathematics, 1974.
- [20] D. E. Rumelhart, J. L. McClelland, and the PDP Research Group, *Parallel Distributed Processing*. Cambridge, MA: The MIT Press, 1986.

- [21] P. V. Bayly, E. E. Johnson, S. F. Idriss, R. E. Ideker, and W. M. Smith, "Efficient electrode spacing for examining spatial organization during ventricular fibrillation," *IEEE Trans. Biomed. Eng.*, vol. 40, pp. 1060-1066, Oct. 1993.
- [22] S. M. Blanchard, W. M. Smith, R. J. Damiano, D. W. Molter, R. E. Ideker, and J. E. Lowe, "Four digital algorithms for activation detection from unipolar epicardial electrograms," *IEEE Trans. Biomed. Eng.*, vol. 36, pp. 256-261, Feb. 1989.
- [23] S. M. Blanchard, R. J. Damiano, T. Asano, W. M. Smith, R. E. Ideker, and J. E. Lowe, "The effects of distant cardiac electrical events on local activation in unipolar epicardial electrograms," *IEEE Trans. Biomed. Eng.*, vol. 34, pp. 539-546, July 1987.
- [24] X. Zhou, P. D. Wolf, D. L. Rollins, Y. Afework, W. M. Smith, and R. E. Ideker, "Effects of monophasic and biphasic shocks on action potentials during ventricular fibrillation in dogs," *Circ. Res.*, vol. 73, pp. 325-334, Aug. 1993.
- [25] F. X. Witkowski, K. M. Kavanagh, P. A. Penkoske, and R. Plonsey, "In vivo estimation of cardiac transmembrane current," *Circ. Res.*, vol. 72, pp. 424-439, Feb. 1993.
- [26] K. M. Kavanagh, R. Plonsey, P. A. Penkoske, and F. X. Witkowski, "Cardiac transmembrane current can differentiate propagating electrical activity from electrotonic activity during ventricular fibrillation," *Circulation*, vol. 88, pp. 1-26, 1993.
- [27] D. E. Rumelhart, G. H. Hinton, and R. J. Williams, "Learning representations by back-propagating errors," *Nature*, vol. 323, pp. 533-536, 1986.
- [28] C. M. Bishop, *Neural Networks for Pattern Recognition*. Oxford: Clarendon Press, 1995.
- [29] Association for the Advancement of Medical Instrumentation (AAMI), "Testing and reporting results of ventricular arrhythmia detection algorithms," AAMI Recommended Practice, 1987.