



US006455762B1

(12) **United States Patent**  
**Chiang et al.**(10) **Patent No.: US 6,455,762 B1**  
(45) **Date of Patent: Sep. 24, 2002**(54) **METHODS OF MODIFYING LIGNIN IN PLANTS BY TRANSFORMATION WITH A 4-COUMARATE COENZYME A LIGASE NUCLEIC ACID**(75) Inventors: **Vincent Lee C. Chiang; Chung-Jui Tsai**, both of Hancock, MI (US);  
**Wen-Jing Hu**, Houston, TX (US)(73) Assignee: **Board of Control of Michigan Technological University**, Houghton, MI (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **08/969,046**(22) Filed: **Nov. 12, 1997**(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00; C12N 15/82**(52) **U.S. Cl.** ..... **800/298; 800/278; 800/279; 800/286; 800/294; 800/301**(58) **Field of Search** ..... 536/23.6; 425/468, 425/295, 419, 410; 800/285, 278, 290, 279, 286, 294, 298, 301(56) **References Cited****U.S. PATENT DOCUMENTS**5,107,065 A 4/1992 Shewmaker et al. .... 800/205  
5,190,931 A 3/1993 Inouye ..... 435/91  
5,850,020 A \* 12/1998 Bloksberg et al. .... 800/205  
5,952,486 A \* 9/1999 Bloksberg et al. .... 536/23.6  
6,204,434 B1 \* 3/2001 Bloksberg et al. .... 800/290**FOREIGN PATENT DOCUMENTS**WO 93/05160 3/1993 ..... C12N/15/54  
WO 98/11205 3/1998  
WO 98/11205 3/1998 ..... C12N/9/02**OTHER PUBLICATIONS**GenBank Accession No. M62755, 1993.\*  
GenBank Accession No. X13324, 1993.\*  
GenBank Accession No. U12012, 1996.\*  
GenBank Accession No. U12013, 1996.\*  
GenBank Accession No. X52623, 1993.\*  
Haug et al. In Vitro Cellular and Developmental Biology. 1991. vol. 27P: 201–207.\*  
Kajita et al. 1996. Plant Cell Physiol. vol.: 37: 957–965.\*  
Napoli et al. The Plant Cell. 1989. vol. 2: 278–289.\*  
Smith et al. Nature. 1988. vol. 334: 724–726.\*  
Tsai et al. Plant Cell Reports. 1994. vol. 14: 94–97.\*  
Lee et al. Plant Molecular Biology, 1995. vol. 28: 871–884.\*  
Ehltting et al, Three 4-coumarate: coenzyme A ligases in *Arabidopsis thaliana* represent two evolutionary divergent classes in angiosperms.; *The Plant Journal*, 19 (1) 9–20 (1991).  
Dandekar, A.M., et al., “Transformation and Foreign Gene Expression in Micropropagated Douglas–Fir (*Pseudotsuga menziesii*)”, *Bio/technology*, 5, 587–590 (Jun. 1987).  
Douglas, C.J., et al., “Exonic Sequences Are Required for Elicitor and Light Activation of a Plant Defense Gene, but Promoter Sequences Are Sufficient for Tissue Specific Expression”, *The EMBO Journal*, 10, 1767–1775 (1991).Fillatti, J.J., et al., “Agrobacterium Mediated Transformation and Regeneration of Populus”, *Molecular and General Genetics*, 206, 192–199 (1987).Hu, W., et al., “Compartmentized Expression of Two Structurally and Functionally Distinct 4–Coumarate:CoA Ligase Genes in Aspen (*Populus tremuloides*)”, *Proc. Natl. Acad. Sci. USA*, 95, 5407–5412 (Apr. 1998).Huang, Y., et al., “*Agrobacterium rhizogenes*–Mediated Genetic Transformation and Regeneration of a Conifer: *Larix decidua*”, *In Vitro Cell. Dev. Biol.*, 27P, 201–207 (Oct. 1991).Kajita, S., et al., “Alterations in the Biosynthesis of Lignin in Transgenic Plants with Chimeric Genes for 4–Coumarate:Coenzyme A Ligase”, *Plant Cell Physiology*, 37, 957–965, (1996).McGranahan, G.H., et al., “Agrobacterium–Mediated Transformation of Walnut Somatic Embryos and Regeneration of Transgenic Plants”, *Bio/technology*, 6, 800–804, (Jul. 1988).Minocha, S.C., et al., “Tissue Culture and Genetic Transformation in *Betula Papyrifera* and *Populus Tremuloides*”, Proceedings, 1986 TAPPI Research and Development Conference, Raleigh, NC, 89–92 (Sep. 28–Oct. 1, 1986).Sullivan, J., et al., “Transformation of *Liquidambar styraciflua* using *Agrobacterium tumefaciens*”, *Plant Cell Reports*, 12, 303–306 (1993).Uhlmann, A., et al., “Molecular Cloning and Expression of 4–Coumarate:coenzyme A Ligase, an Enzyme Involved in the Resistance Response of Soybean (*Glycine Max L.*) Against Pathogen Attack”, *Plant Physiology*, 102, 1147–1156 (1993).Wilde, H.D., et al., “Expression of Foreign Genes in Transgenic Yellow–Poplar Plants”, *Plant Physiology*, 98, 114–120 (1992).Zhang, X., et al., “Molecular Cloning of 4–Coumarate:Coenzyme A Ligase in Loblolly Pine and the Roles of the Enzyme in the Biosynthesis of Lignin in Compression Wood”, *Plant Physiology*, 113, 65–74 (1997).Kajita, et al. “Alterations in the Biosynthesis of Lignin in Transgenic Plants with Chimeric Genes for 4–Coumarate:Coenzyme A Ligase”, *Plant Cell Physiol.*, 37(7): pp. 957–965, 1996.Holmberg, et al. “Transgenic Tobacco Expressing Vitreoscilla Hemoglobin Exhibits Enhanced Growth and Altered Metabolite Production”, *Nature Biotechnology*, vol. 15, pp. 244–247, Mar. 1997.Doerner, et al. “Control of Root Growth and Development by Cyclin Expression”, *Nature*, vol. 380, pp. 520–523, Apr. 11, 1996.Szerszen, et al. “*iaglu*, a Gene from *Zea mays* Involved in Conjugation of Growth Hormone Indole–3–Acetic Acid”, *Science*, vol. 265, pp. 1699–1701, Sep. 16, 1994.

\* cited by examiner

Primary Examiner—Amy J. Nelson

(74) Attorney, Agent, or Firm—Joseph A. Gemignani; Teresa J. Welch; Sara Dastgheib-Vinarov

(57) **ABSTRACT**

The invention pertains to methods of altering growth, lignin content, coniferyl and sinapyl alcohol units in the lignin structure, disease resistance and cellulose content in plants by transformation with a lignin pathway p-coumarate Co-enzyme A ligase (4CL) nucleic acid.

**38 Claims, 8 Drawing Sheets**




FIG. 1




FIG. 2

Pt4CL1g--4




FIG. 3





FIG. 4



**FIG. 5**




FIG. 6

Pt4CL1p-GUS




FIG. 7



Pt4CL2p-GUS




FIG. 8

**METHODS OF MODIFYING LIGNIN IN  
PLANTS BY TRANSFORMATION WITH A 4-  
COUMARATE COENZYME A LIGASE  
NUCLEIC ACID**

**STATEMENT OF GOVERNMENT RIGHTS**

The present invention was made at least in part with the support of the United States Government via a grant from the U.S. Department of Agriculture (Grant No. 95-37103-2061). The Government may have certain rights in the invention.

**FIELD OF THE INVENTION**

The invention relates to genetically modifying trees through manipulation of the lignin biosynthesis pathway, and more particularly, to genetically modifying trees through the down regulation of p-coumarate Co-enzyme A ligase (CCL) to achieve faster growth, and/or altered lignin content, and/or altered lignin structure, and/or altered cellulose content and/or disease resistance of the trees and to the use of promoters of the CCL genes to drive gene expression specifically in xylem tissue or specifically in epidermal tissues.

**BACKGROUND OF THE INVENTION**

Genetic engineering of forest tree species to conform to desired traits has shifted the emphasis in forest tree improvement away from the traditional breeding programs during the past decade. Although research on genetic engineering of forest trees has been vigorous, the progress has been slow due.

The ability to make trees grow faster and be disease resistant to produce the highest volume of wood in the shortest period of time has been and continues to be the top objective of many forest products company worldwide. The ability to genetically increase the optimal growth of trees would be a commercially significant improvement. Faster growing trees could be used by all sectors of the forest and wood products industry worldwide.

Lignin, a complex phenolic polymer, is a major component in cell walls of secondary xylem. In general, lignin constitutes 25% of the dry weight of the wood, making it the second most abundant organic compound on earth after cellulose. Although lignin plays an important role in plants, it usually represents an obstacle to utilizing biomass in several applications. For example, in woodpulp production, lignin has to be removed through expensive and polluting processes in order to recover cellulose.

Thus, it is desirable to genetically engineer plants with reduced lignin content and/or altered lignin composition that can be utilized more efficiently. Trees that could be genetically engineered with a reduced amount of lignin would be commercially valuable. These genetically engineered trees would be less expensive to pulp because, in essence, part of the pulping has already been performed due to the reduced amount of lignin.

Trees with increased cellulose content would also be commercially valuable to the pulp and paper industry.

Disease resistance in plants is also a most desirable plant trait. The impact of disease resistance in trees on the economy of forest products industry worldwide is significant.

Promoters that target specific plant tissue could be useful in manipulating gene expression to enable the engineering of traits of interest in specific tissue of plants, such as, xylem and epidermal tissues.

Although studies have revealed several general properties of plant p-coumarate Co-enzyme A ligase (CCL), the role of CCL in regulating the synthesis of monolignols in response to different states of development and environmental stress in tree species remains largely unknown. Furthermore, multiple CCL isoforms are normally present in plants, channeling phenolic compounds to the biosynthesis of not only lignin but also other phenylpropanoids, such as flavonoids. Since CCL isoforms have not been previously cloned from tree species for the identification of their biochemical functions, the presence of CCL isoforms remains so far as a challenge to a specific control of metabolic flux to the lignin biosynthesis in tree species.

**SUMMARY OF THE INVENTION**

The invention provides a method to genetically alter trees through the down regulation of p-coumarate Co-enzyme A ligase (CCL). Such down regulation of CCL results in faster growth, and/or reduced lignin content, and/or altered lignin structure, and/or altered cellulose content and/or disease resistance. The invention also provides for genetically engineered trees which have been altered to down regulate p-coumarate Co-enzyme A ligase (CCL) to achieve faster growth, and/or reduced lignin content, and/or altered lignin structure, and/or increased cellulose content and/or increased disease resistance. The invention also provides tissue specific promoters of the CCL genes that can be used to manipulate gene expression in target tissue such as xylem and epidermal tissues.

It is one object of the present invention to down regulate p-coumarate Co-enzyme A ligase (CCL) in trees.

It is another object of the present invention to provide a method to genetically alter trees to grow faster.

It is another object of the present invention to provide a method to genetically alter the growth of trees through manipulation the lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to provide genetically altered trees with an accelerated growth characteristic.

It is another object of the present invention to provide transgenic trees with an accelerated growth characteristic which have been genetically altered by down regulating lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to provide a method to genetically alter trees to reduce their lignin content.

It is another object of the present invention to provide a method to genetically alter the lignin content of trees through manipulation of a lignin pathway enzyme.

It is another object of the present invention to genetically engineer trees which have reduced lignin content through manipulation of lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to provide genetically altered trees with a reduced lignin content.

It is another object of the present invention to provide transgenic trees with reduced lignin content which have been genetically altered by down regulating the p-coumarate Co-enzyme A ligase (CCL).

It is another object of the present invention to provide a method to genetically alter trees to change their lignin structure through manipulation of lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to provide trees with altered lignin structure.

## 3

It is another object of the present invention to provide a method to increase the cellulose content in trees.

It is another object of the present invention to provide a method to increase the cellulose content of trees through the manipulation of a lignin pathway enzyme.

It is another object of the present invention to provide trees with increased cellulose content.

It is another object of the present invention to provide transgenic trees having increased cellulose content from the down regulation of CCL.

It is another object of the present invention to provide a method to genetically alter trees to increase their disease resistance.

It is another object of the present invention to provide a method to genetically alter trees to be more disease resistant through manipulation of the lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to genetically alter trees to increase their disease resistance to fungal pathogens.

It is another object of the present invention to provide trees with increased disease resistance.

It is another object of the present invention to provide transgenic trees with increased disease resistance through down regulation of the lignin pathway p-coumarate Co-enzyme A ligase.

It is another object of the present invention to provide a method using a promoter of a CCL gene to target gene expression in specific plant tissue.

It is another object of the present invention to provide a method using a promoter of a CCL gene to target gene expression specifically in plant xylem.

It is another object of the present invention to provide a method using a promoter of the CCL gene to target gene expression specifically in the epidermal tissues of plants.

It is another object of the present invention to provide a CCL gene promoter that targets gene expression specifically in the xylem of plants.

It is another object of the present invention to provide a CCL gene promoter that targets gene expression specifically in the epidermal tissues of plants.

Other features and advantages of the invention will become apparent to those of ordinary skill in the art upon review of the following drawing, detailed description and claims.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic of a phenylpropanoid pathway;

FIG. 2 is a diagram of Agrobacterium T-DNA construct pACCL1;

FIG. 3 is a restriction map of genomic clone PtCCL1g-4;

FIG. 4 is a restriction map of genomic clone PtCCL2g-11;

FIG. 5 is a restriction map of subcloned PtCCL1 gene promoter p7Z-4XS;

FIG. 6 is a restriction map of subcloned PtCCL2 gene promoter pSK-11HE

FIG. 7 is an Agrobacterium T-DNA construct of PtCCL1 promoter and GUS fusion gene, PtCCL1p-GUS; and

FIG. 8 is an Agrobacterium T-DNA construct of PtCCL2 promoter and GUS fusion gene, PtCCL2p-GUS.

Before one embodiment of the invention is explained in detail, it is to be understood that the invention is not limited in its application to the details set forth in the following

## 4

description of the preferred embodiment. The invention is capable of other embodiments and of being practiced or being carried out in various ways. Also, it is to be understood that the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The invention pertains to genetically down regulating a lignin pathway p-coumarate Co-enzyme A ligase (CCL). Trees which have been genetically transformed to down regulate CCL will hereafter be called transgenic trees. Such down regulation can result in faster growing trees. Such down regulation can result in a reduction in the lignin content of the trees and/or altered lignin structure. Such down regulation can result in increased cellulose content. Such down regulation can result in increased tree disease resistance. Further, by using a specific promoter of CCL, targeted tissue gene expression can be achieved in either the xylem or the epidermal tissues of the plant.

## A. CCL

Lignin is synthesized by the oxidative coupling of three monolignols (coumaryl, coniferyl and sinapyl alcohols) formed via the phenylpropanoid pathway as shown in FIG. 1. Reactions in the phenylpropanoid pathway include the deamination of phenylalanine to cinnamic acid followed by hydroxylations, methylations and activation of substituted cinnamic acids to coenzyme A (CoA) esters. The CoA esters are then reduced to form monolignols which are secreted from cells to form lignin.

The products of the phenylpropanoid pathway are not only required for the synthesis of lignin but also required for the synthesis of a wide range of aromatic compounds including flavonoids, phytoalexins, stilbenes and suberin.

In angiosperms (hardwoods), lignin is composed of both coniferyl and sinapyl alcohol and is classified as guaiacyl-syringyl lignin. Grasses synthesize a third precursor (p-coumaryl alcohol) which is polymerized along with coniferyl and sinapyl alcohol. In gymnosperms (softwoods), lignin is composed of mainly coniferyl alcohol and is classified as guaiacyl lignin.

In the phenylpropanoid pathway, CCL activates a number of cinnamic acid derivatives, including p-coumaric acid, caffeic acid, ferulic acid, 5-hydroxyferulic acid and sinapic acid. The resulting products, CoA esters, serve as substrates for entry into various branch pathways, such as lignin, flavonoids, phytoalexins, stilbenes and suberin. The esterification reactions catalyzed by CCL require high energy and the reactions are not likely to occur without CCL. CCL is important in making a continuous flow of the lignin biosynthesis pathway. CCL is also important because it is located at the branching points of the phenylpropanoid metabolism. CCL is suggested to play a pivotal role in regulating carbon flow into specific branch pathways of the phenylpropanoid metabolism in response to stages of development and environmental stress.

The basic properties of CCL are quite uniform. CCL depends on ATP as a cosubstrate and requires  $Mg^{2+}$  as a cofactor. The optimal pH for CCL ranges from pH 7.0 to 8.5 and the molecular weight of CCL isoforms from various plant species ranges from 40 kd to 75 kd. Most CCLs have high affinity with substituted cinnamic acids. CCL has the highest activity with p-coumaric acid.

CCL cDNA sequences have been reported for parsley, potato, soybean, loblolly pine, Arabidopsis, Lithospermum

and tobacco. CCL genes have been isolated and sequenced for parsley, rice, potato and loblolly pine. The analysis of CCL cDNAs and genes indicates that CCL is encoded by multiple/divergent genes in rice, soybean, and *Lithospermum*, very similar genes in parsley, potato, tobacco and loblolly pine, and a single gene in *Arabidopsis*. CCL promoters have been isolated and sequenced for parsley, rice and potato.

Alignment of deduced amino acid sequences of cloned plant CCL sequences reveals two highly conserved regions. The first conserved region (SSGTTGLPKG)(SEQ ID NO:7), proposed to designate a putative AMP-binding region, is very rich in Gly, Ser and Thr and is followed by a conserved Lys. The second conserved region (GEICIRG)(SEQ ID NO:8) contains one common Cys residue. The amino acid sequences of CCL from plants contain a total of five conserved Cys residues.

The CCL genes of parsley, potato and rice contain five exons and four introns. The CCL genes also share the same exon/intron splice junction sites but have different lengths of introns. The genomic sequences of loblolly pine CCL are composed of four exons and three introns. It has been found that two similar CCL genes of the same species may differ slightly in length of intron as shown in two parsley genes (PC4CL1 and PC4CL2) and in two loblolly pine genes (LP4CL1 and LP4CL2).

By Northern blot analysis, it has been shown that CCL is expressed in leaf, shoot tip, stem, root, flower and cell culture. Two similar CCL cDNAs in parsley, potato and tobacco have been shown to be expressed at similar level in response to the environmental stress and during different developmental stages. Two distinct CCL cDNAs in soybean and *Lithospermum* have shown different expression levels when pathogens or chemicals were applied to the cell cultures. It appears that the expression of the CCL genes is developmentally regulated and inducible by many environmental stresses at the transcription level.

Genetic transformation with a CCL sequence can result in several significant affects. The description of the invention hereafter refers to aspen, and in particular quaking aspen (*Populus tremuloides* Michx) when necessary for the sake of example. However, it should be noted that the invention is not limited to genetic transformation of aspen. The method of the present invention is capable of being practiced for other trees, including for example, other angiosperms, other gymnosperm forest tree species, etc.

Preferably, the CCL down regulation is accomplished through transformation with a homologous CCL sequence in an antisense orientation. However, it should be noted that a heterologous antisense CCL sequence could be utilized and incorporated into a tree species to down regulate CCL if the heterologous CCL gene sequence has a high nucleotide sequence homology, approximately higher than 70%, to the endogenous CCL gene sequence of that tree species.

In addition, trees transformed with a sense CCL sequence could also cause a sequence homology-based cosuppression of the expression of the transgene and endogenous CCL gene, thereby achieving down regulation of CCL in these trees.

#### B. Isolation of CCL cDNAs

The present invention utilizes a homologous CCL sequence to genetically alter trees. The preferred embodiment of the invention as further described below utilizes a cDNA clone of the quaking aspen CCL gene.

Two aspen (*Populus tremuloides* Michx) cDNAs encoding two distinct CCL isoforms, PtCCL1 and PtCCL2 have

been cloned. PtCCL1 cDNA is lignin pathway-specific and is different from PtCCL2 cDNA, which is involved in flavonoid synthesis. The cloning of PtCCL1 and PtCCL2 cDNAs and the identification of their biochemical functions will be discussed in more length below. PtCCL1 and PtCCL2 genomic clones including their 5'-end regulatory promoter sequences were also isolated. The promoter of PtCCL1 (PtCCL1p) directs xylem tissue-specific gene expression in a plant, whereas the promoter of PtCCL2 (PtCCL2p) drives the expression of genes specifically in epidermal tissues of stem and leaf of a plant. These tissue specific promoters will be discussed in more length in Section I below.

Two CCL cDNAs, PtCCL1 and PtCCL2, have been isolated from quaking aspen using either a conventional cDNA library screening method or a PCR-based cDNA cloning method. It should be noted that the methods described below are set forth as an example and should not be considered limiting. These CCL cDNA clones are available from Michigan Technological University, Institute of Wood Research, Houghton, Mich.

Young leaves and shoot tips are collected from greenhouse-grown quaking aspen (*Populus tremuloides* Michx). Differentiating xylem and sclerenchyma are collected from three to four year old quaking aspen. The bark is peeled from the tree exposing the developing secondary xylem on the woody stem and the sclerenchyma on the inner side of the bark. Developing secondary xylem and sclerenchyma are scraped from the stem and bark with a razor blade and immediately frozen in liquid nitrogen until use.

Total RNA is isolated from the young leaves, shoot tips, xylem and sclerenchyma following the method of Bugos RC et al. (1995), RNA Isolation from Plant Tissue Recalcitrant to Extraction in Guanidine, *Biotechniques* 19(5):734-737. Poly(A)<sup>+</sup>RNA is purified from total RNA using Poly(A)<sup>+</sup>mRNA Isolation Kit from Tel-test B. Inc. A unidirectional Lambda gt22 expression cDNA library was constructed from the xylem mRNA using Superscript  $\lambda$  System from Life Technologies, Inc. and Gigapack Packaging Extracts from Stratagene. The PtCCL1 cDNA was obtained by screening the cDNA library with a <sup>32</sup>P-labeled parsley 4CL cDNA probe. The parsley 4CL cDNA (pc4CL2) has Genbank accession number X13325(SEQ ID NO:15).

The PtCCL2 cDNA was obtained by RT-PCR. The reverse transcription of total RNA isolated from shoot tips was carried out using the Superscript II reverse transcriptase from Life Technologies. Two sense primers (R1S, 5'-TTGGATCCGGIACIACIGGIYTIICIAARGG-3'(SEQ ID NO:9) and H1S, 5'-TTGGATCCGTIGCICARCARGTIGAYGG-3')(SEQ ID NO:10) are designed around the first consensus AMP-binding region of CCL that was previously discussed. One antisense primer (R2A, 5'-ATGTCGACCICGDATRCADATYTCICC-3')(SEQ ID NO:11) is designed based on the sequence of the putative catalytic motif GEICIRG(SEQ ID NO:8). One fifth of the reverse transcription reaction (4  $\mu$ l) is used as the template in a 50  $\mu$ l PCR reaction containing 1 $\times$  reaction buffer, 200  $\mu$ M each deoxyribonucleotide triphosphate, 2  $\mu$ M each R1S and oligo-dT (20 mer) primers, and 2.5 units of Taq DNA polymerase. The PCR reaction mixture was denatured at 94 $^{\circ}$  C. for 5 minutes followed by 30 cycles of 94 $^{\circ}$  C./45 seconds, 50 $^{\circ}$  C./1 minute, 72 $^{\circ}$  C./90 seconds and is ended with a 5 minute extension at 72 $^{\circ}$  C. 2  $\mu$ l of the PCR amplification products are used for a second run PCR re-amplification using primers H1S and R2A. A 0.6 kb PCR fragment is cloned using the TA Cloning Kit from Invitrogen and used

as a probe to screen an aspen genomic library to obtain the PtCCL2 genomic clone. Two primers (2A, 5'-TCTGTCTAGATGATGTCGTGGCCACGG-3'(SEQ ID NO:12) and 2B, 5'-TTAGATCTCTAGGACATGGTGGTGGC-3')(SEQ ID NO:13) are designed based on the genomic sequence of PtCCL2 at around the deduced transcription start site and stop codon for the cloning of PtCCL2 cDNA by RT-PCR as described above using total RNA isolated from shoot tips.

The DNA sequences of PtCCL1 and PtCCL2 cDNA were determined using ΔTaq Cycle Sequencing Kit from Amersham.

The PtCCL1 cDNA has an open reading frame of 1620 bp which encodes a polypeptide of 540 amino acid residues with a predicted molecular weight of 59 kd and pI of 5.8. The nucleotide sequence of the aspen CCL cDNA clone PtCCL1 is set forth as SEQ ID NO:1. The deduced amino acid sequence for the aspen CCL1 protein is set forth as SEQ ID NO:2.

The PtCCL2 cDNA has an open reading frame of 1713 bp which encodes a polypeptide of 571 amino acid residues with a predicted molecular weight of 61.8 kd and pI of 5.1. The nucleotide sequence of the aspen CCL cDNA clone PtCCL2 is set forth as SEQ ID NO:3. The deduced amino acid sequence for the aspen CCL2 protein is set forth as SEQ ID NO:4.

The aspen PtCCL1 cDNA shares a 59–74% identity at the nucleotide level and 59–81% identity at the amino acid level with other prior reported CCL cDNAs and genes, whereas the PtCCL2 cDNA shares a 60–73% identity at the nucleotide level and 57–74% at the amino acid level with other CCL cDNAs and genes as set forth in the following table.

TABLE 1

cDNA*	Comparison of PtCCL1 and PtCCL2 Nucleotide and Predicted Amino Acid Sequence to Each Other and Other CCL Sequences			
	DNA IDENTITY % PtCCL1	DNA IDENTITY % PtCCL2	AMINO ACID IDENTITY % PtCCL1	AMINO ACID IDENTITY % PtCCL2
PtCCL1		62		63
LE4CL1	69	62	71	64
LE4CL2	60	71	59	73
GM14	74	67	81	69
GM16	62	73	65	73
NT4CL1	67	62	75	74
NT4CL2	66	63	75	66
PC4CL1	66	64	71	64
PC4CL2	66	63	72	64
ST4CL1	67	63	75	64
AT4CL	66	63	70	61
LP4CL	61	64	63	67
OS4CL1	59	60	59	57

\*PtCCL1: aspen CCL

PtCCL2: aspen CCL

LE4CL1 and LE4CL2: *Lithospermum erythrorhizon* CCL

GM14 and GM16: soybean CCL

NT4CL1 and NT4CL2: tobacco CCL

PC4CL1 and PC4CL2: parsley CCL

ST4CL1: potato CCL

AT4CL: Arabidopsis CCL

LP4CL: loblolly pine CCL

OS4CL1: rice CCL

The results of sequence analysis, phylogenetic tree and genomic Southern blot analysis indicate that PtCCL1 and PtCCL2 cDNAs encode two distinct CCLs that belong to two divergent gene families in aspen. The deduced amino acid sequence for the PtCCL2 protein contains a longer

N-terminal sequence than the PtCCL1 protein but shows profound similarity in the central and C-terminal portions of protein to the PtCCL1 protein.

PtCCL1 and PtCCL2 cDNAs display distinct tissue-specific expression patterns. The PtCCL1 sequence is expressed highly in the secondary developing xylem and in the 6th to 10th internodes whereas the PtCCL2 sequence is expressed in the shoot tip and leaves. These tissue-specific expression patterns were investigated by fusing promoters of PtCCL1 and PtCCL2 genes to a GUS reporter gene. The tissue specific promoters for PtCCL1 and PtCCL2 will be discussed in more length in Section I below.

The substrate specificity of PtCCL1 and PtCCL2 is also different from each other as determined using recombinant proteins produced in *E. coli*. PtCCL1 utilized p-coumaric acid, caffeic acid, ferulic acid and 5-hydroxyferulic acid as substrates. PtCCL2 showed activity to p-coumaric acid, caffeic acid and ferulic acid but not to 5-hydroxyferulic acid.

Specifically, PtCCL1 and PtCCL2 were used to construct expression vectors for *E. coli* expression. The substrate specificity of PtCCL1 and PtCCL2 were tested using fusion proteins produced in *E. coli*. Two plasmids, pQE/CCL1 and pQE/CCL2, were constructed in which the coding regions of PtCCL1 and PtCCL2, respectively were fused to N-terminal His tags in expression plasmids pQE-31 and pQE-32 (QIAGEN, Chatsworth, Calif). The recombinant proteins of PtCCL1 and PtCCL2 produced by *E. coli* are approximately 59 kd and 63 kd, respectively.

The two recombinant proteins were tested for their activity in utilizing cinnamic acid derivatives. PtCCL1 recombinant protein showed 100, 58, 71, 18 and 0% relative activity to p-coumaric acid, caffeic acid, ferulic acid, 5-hydroxyferulic acid and sinapic acid, respectively. PtCCL2 recombinant protein exhibited 100, 14, 27, 0 and 0% relative activity to p-coumaric acid, caffeic acid, ferulic acid, 5-hydroxyferulic acid and sinapic acid, respectively. Neither recombinant protein showed detectable activity to sinapic acid.

The results of the tissue-specific expression pattern and substrate specificity suggests that in addition to the general function of CCL, PtCCL1 apparently is more related to lignin synthesis in the xylem tissue and PtCCL2 apparently is more likely involved in flavonoid synthesis and UV protection.

It should be noted that the isolation and characterization of the PtCCL1 and PtCCL2 cDNA clones is described in Kawaoka A, Chiang VL (1995), The Molecular Cloning and Expression of Syringyl- and Guaiacyl-Specific Hydroxycinnamate:CoA Ligases from Aspen (*Populus tremuloides*), Proceedings of the 6th International Conference on Biotechnology in the Pulp and Paper Industry, Vienna, Austria; and in Hu, Wen-Jing, Isolation and Characterization of p-coumarate Co-enzyme A ligase cDNAs and Genes from Quaking Aspen (*Populus tremuloides* Michx), Ph.D Dissertation, Michigan Technological University, Houghton, Mich. (1997); which are both herein incorporated by reference.

### C. Transformation and Regeneration

Several methods for gene transformation of plant species with the CCL sequence are available such as the use of a transformation vector, agroinfection, electroinjection, particle bombardment with a gene gun or microinjection.

Preferably, a CCL cDNA clone is positioned in a binary expression vector in an antisense orientation under the control of double cauliflower mosaic virus 35S promoter.

The vector is then preferably mobilized into a strain of *Agrobacterium* species such as *tumefaciens* strain C58/pMP90 and is used as the DNA delivery system due to its efficiency and low cost.

For example, with reference to FIG. 2, the binary expression pACCL1 used for plant transformations is shown. Specifically, the PtCCL1 CDNA is inserted in an antisense orientation into Pac I and BamH I sites between the double CaMV 35S/AMV RNA4 and the 3' terminator sequence of the nopaline synthase gene in a binary cloning vector pACCL1 (FIG. 2). The binary vector containing hygromycin phosphotransferase (HPT) gene is modified from pBin 19.

The gene construct pACCL1 is available from Michigan Technological University, Institute of Wood Research, Houghton, Mich.

The binary vector construct is mobilized in *Agrobacterium tumefaciens* using the freeze-thaw method of Holsters et al., *Mol. Gen. Genet.* 163:181-187 (1978). For the freeze-thaw method, 1.5 ml of overnight cultures *Agrobacterium tumefaciens* strain C58/pMP90 is pelleted at 5000×g for 3 minutes at 4° C. and suspended in 1 ml of ice cold 20 mM CaCl<sub>2</sub>. To the suspension is added 10 μl binary vector DNA (from an alkaline lysis miniprep) and mixed by pipetting. The microcentrifuge tube is then frozen in liquid nitrogen for 5 minutes and thawed at 37° C. for 5 minutes. After being cooled on ice, 1 ml of LB is added and the mixture is incubated at 28° C. for 2 hours with gentle shaking. 200 μl of the cells is spread onto LB plates containing gentamycin and kanamycin and incubated at 28° C. for 2 days. Colonies grown on the selection plates are randomly picked or miniprep and restriction enzyme digestion analysis is used to verify the integration.

The resulting binary vector containing *Agrobacterium* strain is used to transform quaking aspen according to Tsai et al., *Agrobacterium-Mediated Transformation of Quaking Aspen (Populus tremuloides) and Regeneration of Transgenic Plants*, *Plant Cell Rep.* 14:94-97 as set forth below.

Explants of young leaves from cuttings of aspen are obtained by cutting leaf disks of approximately 7 mm square from the young leaves along the midrib of the leaves. The explants are surface sterilized in 20% commercial bleach for 10 minutes followed by rinsing 3 times with sterile double-distilled water.

All of the culture media used includes the basal medium of woody plant medium (WPM) as described in Lloyd et al., *Proc. Int. Plant Prop. Soc.* 30:421-437 (1980) and supplemented with 2% sucrose, 650 mg/L calcium gluconate and 500 mg/L MES are added as pH buffers as described in Tsai, *Plant Cell Reports*, 1994. All culture media is adjusted to pH 5.5 prior to the addition of 0.075% Difco Bacto Agar and then autoclaved at 121° C. and 15 psi for 20 minutes. Filter sterilized antibiotics are added to all culture media after autoclaving. All culture media are maintained at 23±1° C. in a growth chamber with 16 hour photoperiods (160 μE×m<sup>-2</sup>×S<sup>-1</sup>) except for callus induction (as will be described later) which is maintained in the dark.

The sterilized explants are then inoculated with the mobilized vector with an overnight-grown agrobacterial suspension containing 20 μM acetosyringone. After cocultivation for 2 days, the explants are washed in 1 mg/ml claforan and ticarcillin for 2 hours with shaking to kill *Agrobacterium*. The explants are blotted dry with sterile Whatman No. 1 filter paper and transferred onto callus induction medium containing 50 mg/L kanamycin and 300 mg/L claforan to induce and select transformed callus. The callus induction medium is the basal medium with the addition of

6-benzyladenine (BA) and 2,4-dichlorophenoxyacetic acid (2,4-D) at concentrations of 0.5 mg/L and 1 mg/L, respectively, to induce callus.

The kanamycin-resistant explants are then subcultured on fresh callus induction media every two weeks. Callus formation occurs after approximately four weeks. Formed callus are separated from the explant and subcultured periodically for further proliferation.

When the callus clumps reach approximately 3 mm in diameter, the callus clumps are transferred to shoot regeneration medium. The shoot regeneration medium is the basal medium containing 50 mg/L kanamycin, 0.5 mg/L thidiazuron (TDZ) as a plant growth regulator and cefotaxime at 300 mg/L to kill *Agrobacterium*. Shoots were regenerated about 4 weeks after callus is transferred to regeneration medium.

As soon as the shoots are regenerated, they are immediately transferred to hormone-free elongation medium containing 50 mg/L kanamycin and, whenever necessary, cefotaxime (300 mg/L), to promote elongation. Green and healthy shoots elongated to 2-3 cm in length are excised and planted separately in a hormone-free rooting medium containing 50 mg/L kanamycin. The efficient uptake of kanamycin by shoots during their rooting stage provides the most effective selection for positive transformants. Transgenic plants are then transplanted into soil medium of vermiculite:peatmoss:perlite at 1:1:1 and grown in the greenhouse.

The above described transformation and regeneration protocol is readily adaptable to other tree species. Other published transformation and regeneration protocols for tree species include Danekar et al., *Bio/Technology* 5:587-590 (1987); McGranahan et al., *Bio/Technology* 6:800-804 (1988); McGranahan et al., *Plant Cell Reports* 8:512-616 (1990); Chen, PhD Thesis, North Carolina State University, Raleigh, N.C. (1991); Sullivan et al., *Plant Cell Reports* 12:303-306 (1993); Huang et al., *In Vitro Cell Dev. Bio.* 4:201-207 (1991); Wilde et al., *Plant Physiol.* 98:114-120 (1992); Minocha et al., 1986 *Proc. TAPPI Research and Development Conference*, TAPPI Press, Atlanta, pp. 89-91 (1986); Parsons et al., *Bio/Technology* 4:533-536 (1986); Fillatti et al., *Mol. Gen. Genet* 206:192-199 (1987); Pythoud et al., *Bio/Technology* 5:1323-1327 (1987); De Block, *Plant Physiol.* 93:1110-1116 (1990); Brasileiro et al., *Plant Mol. Bio* 17:441-452 (1991); Brasileiro et al., *Transgenic Res.* 1:133-141 (1992); Howe et al., *Woody Plant Biotech.*, Plenum Press, New York, pp.283-294 (1991); Klopfenstein et al., *Can. J. For. Res.* 21:1321-1328 (1991); Leple et al., *Plant Cell Reports* 11:137-141 (1992); and Nilsson et al. *Transgenic Res.* 1:209-220 (1992).

#### D. Phenotype Changes

The results of the transformation can be confirmed with conventional PCR and Southern analysis. For example, transferring CCL CDNA in an antisense orientation down regulates CCL in the tree. Expression of the CCL has been found to be blocked up to 96 percent in some transgenic trees.

After acclimation, the transgenic aspen display an unusual phenotype, including big curly leaves, thick diameters, longer internodes, more young leaves in the shoot tip and a red pigmentation in the petioles extending into midvein leaves. Red coloration of the developing secondary xylem tissues is observed after peeling of the bark in the transgenic plants.

#### E. Accelerated Growth

Down regulation of CCL alters growth of the transgenic trees. For example, transformation with an antisense CCL

sequence accelerates the growth of the tree. Enhanced growth is markedly noticeable at all ages. In particular, the transgenic trees show enhanced growth in the form of thicker stems and enlarged leaves as compared to control trees. These characteristics are retained in the vegetative propagules of these transgenic trees. Table 2 sets forth exemplary data with respect to several lines of transgenic quaking aspen grown in the greenhouse after eight months. Volume represents the overall quantitative growth of the tree.

TABLE 2

Growth Measurement for Control and Transgenic Plants				
PLANT #	HEIGHT (cm)	DIAMETER (cm)*	VOLUME (cm <sup>3</sup> )*	AVERAGE LENGTH OF INTERNODE (cm)
Control 1	247.7	1.08	75.6	2.6
Control 2	250.2	1.01	66.8	2.8
11-1	304.8	1.15	105.5	3.1
11-2	248.9	1.01	66.4	3.4
11-3	241.3	0.84	44.6	3.2
11-4	288.3	0.94	66.7	3.4
11-5	246.4	0.92	54.6	3.3
11-7	226.7	1.13	75.7	3.4
11-8	289.6	1.16	102.0	3.3
11-9	287.0	1.76	232.6	4.3
11-10	252.7	0.83	45.6	3.1
11-11	247.7	0.86	48.0	3.5
12-1	247.7	1.1	78.4	2.7
12-2	199.4	0.96	48.1	2.5
12-6	294.6	0.92	65.2	3.2
16-1	227.3	0.95	53.7	2.8
16-2	278.1	0.97	68.5	3.4
16-3	265.4	1.09	82.5	3.5
17-2	243.8	0.89	50.5	2.6

\*at 10 cm above ground

The averages for height, diameter, volume and average length between internodes for the control plants are as follows:

Height (cm)	248.95
Diameter (cm)	1.045
Volume (cm <sup>3</sup> )	71.2
Ave. Length of Internodes (cm)	2.7

With respect to height alone, for those transgenic plants (11-1, 11-4, 11-8, 11-9, 12-6, 16-2, 16-3) having a statistically larger height than the control plants, the average height was 286.83 cm as compared to the control plant average height of 248.95 cm.

With respect to diameter alone, for those transgenic plants (11-1, 11-7, 11-8, 11-9) having a statistically larger diameter than the control plants, the average diameter was 1.30 cm as compared to the control plant average diameter of 1.045 cm.

With respect to volume alone, for those transgenic plants (11-1, 11-8, 11-9, 12-1, 16-3) having a statistically larger volume than the control plants, the average volume was 120.2 cm<sup>3</sup> as compared to the control plant average volume of 71.2 cm<sup>3</sup>.

With respect to average length of internodes alone, for those transgenic plants (11-1, 11-2, 11-3, 11-4, 11-5, 11-7, 11-8, 11-9, 11-10, 12-6, 16-2, 16-3) having a statistically larger average length of internodes than the control plants, the average average length of internodes was 3.39 cm as

compared to the control plant average average length of internodes of 2.7° cm.

As demonstrated in Table 2, while there are variations in growth among the transgenic trees, the average length of the internodes for the transgenic trees is consistently and significantly higher than that of the control plants. Variations in the growth of the transgenic trees is normal and to be expected. Preferably, a transgenic tree with a particular growth rate is selected and this tree is vegetatively propagated to produce an unlimited number of clones that all exhibit the identical growth rate.

F. Lignin

Down regulation of lignin pathway CCL results in production of trees with reduced lignin content.

The following table shows the reduction of lignin content and CCL enzyme activity in several transgenic aspen which have been transformed with an homologous antisense CCL sequence.

TABLE 3

Characterization of Transgenic Aspen Plants Harboring Antisense CCL Sequence				
Transgenic Plant #	Lignin Content % Based On Wood Weight	% Lignin Reduction	CCL Enzyme Activity*	% CCL Enzyme Activity Reduction
control	21.4	0.0	868	0
11-1	20.5	4.2	1171	-25
11-2	19.2	10.3	515	45
11-3	20.9	2.3	922	6
11-4	19.7	7.9	1032	-19
11-5	19.7	7.9	691	20
11-7	19.9	7.0	578	38
11-8	20.2	5.6	694	20
11-9	20.4	4.7	806	14
11-10	19.4	9.3	455	51
11-11	20.4	4.7	726	22
12-1	12.8	40.2	49	95
12-2	12.6	41.1	62	93
12-3	11.9	44.4	61	94
12-6	19.8	7.5	786	16
16-1	12.8	40.2	35	96
16-2	20.6	3.7	780	17
16-3	21.0	1.9	795	15
17-1	20.5	4.2	855	9
17-2	21.4	0.0	925	1

\*activity is expressed as pkat/(mg protein) using p-coumaric acid as the substrate

Lignin content was determined according to Chiang and Funaoka (1990) *Holzforchung* 44:147-155. CCL enzyme activity was determined according to Ranjeva et al. (1976), *Biochimie* 58:1255-1262.

The data in Table 3 demonstrates a correlation between down regulation of CCL and reduction in lignin content.

Transgenic trees with reduced lignin content have an altered phenotype in that the stem is more elastic to the touch and the leaves are typically curlier.

It should also be noted that for those transgenic trees (12-1, 12-2, 12-3 and 16-1) with the approximately 40% reduction in lignin content and the corresponding approximately 95% reduction in CCL enzyme levels, all of those transgenic trees had a consistent deep red coloration in the wood of the plant. Accordingly, the deep red color can be used as an identifier of reduced lignin content.

Down regulation of lignin pathway CCL also results in production of trees with an altered lignin structure. Based

upon thioacidolysis (Rolando et al. (1992) Thioacidolysis, Methods in Lignin Chemistry, Springer-Verlag, Berlin, pp 334-349) of plants 12-3 and 16-1, coniferyl alcohol and sinapyl alcohol lignin units are significantly reduced in these two trees as compared to the control tree, as shown in the following table.

TABLE 4

Plant #	Altered Lignin Structure	
	Coniferyl Alcohol Units*	Sinapyl Alcohol Units*
control	733	1700
12-3	283	592
16-1	247	445

\*micro-mole/g of lignin

The alteration of the frequency of the structural units in lignin of these transgenic trees is evidence that the overall structure of lignin in these plants has been genetically altered.

#### G. Cellulose Content

Down regulation of lignin pathway CCL results in increased cellulose content of the transgenic plants. Analysis of control and transgenic aspen for carbohydrate content demonstrate a higher cellulose content in the transgenic trees than the control trees. Particularly, the transgenic trees that have over 40% lignin reduction have about 10-15% higher cellulose content than the control. Data is set forth in the following tables for trees that were transformed with homologous CCL in an antisense orientation:

TABLE 5

Plant #	Analysis of Carbohydrate Components in Transgenic and Control Aspen					
	Glucan	Arabinan	Galactan	Rhamnan	Xylan	Mannan
Control	44.23%	0.47%	0.79%	0.37%	17.19%	1.91%
11-2	49.05%	0.36%	1.05%	0.38%	15.34%	2.04%
11-9	45.95%	0.40%	0.80%	0.37%	17.12%	1.83%
11-10	47.49%	0.43%	0.99%	0.40%	16.24%	2.35%
12-3	50.83%	0.55%	1.24%	0.48%	17.25%	1.77%
16-1	48.14%	0.56%	1.07%	0.48%	19.14%	1.58%
16-2	46.55%	0.34%	0.82%	0.37%	16.75%	2.31%

TABLE 6

Plant #	Comparison of Lignin and Cellulose (glucan) Contents in Transgenic and Control Aspen			
	Lignin		Cellulose	
	Content % on wood	% reduction	Content % on wood	% increase
Control	21.4	0	44.23	0
11-2	19.2	10.3	49.05	11.0
11-9	20.4	4.7	45.95	3.9
11-10	19.4	9.3	47.49	7.4
12-3	11.9	44.5	50.83	15.0
16-1	12.8	40.2	48.14	6.8
16-2	20.6	3.7	46.55	5.2

The procedure for carbohydrate analysis utilized is as follows. About 100 mg of powdery woody tissue with sizes that pass a 80-mesh screen was hydrolyzed with 1 mL of 72% (W/W) H<sub>2</sub>SO<sub>4</sub> for 1 hr at 30° C. Samples were then

diluted to 4% (W/W) H<sub>2</sub>SO<sub>4</sub> with distilled water, fucose was added as an internal standard, and a secondary hydrolysis was performed for 1 hr at 121° C. After secondary hydrolysis, the sugar contents of the hydrolysates are determined by anion exchange high performance liquid chromatography using pulsed amperometric detection. Sugar contents are expressed as % of the weight of the woody tissue used. The above procedures are similar to those in a publication by RC Pettersen and VH Schwandt, 1991, J. Wood Chem & Technol. 11:495-501.

#### H. Increased Disease Resistance

Down regulation of lignin pathway CCL results in production of trees with increased disease resistance, and in particular, with increased fungal pathogen resistance.

In particular, greenhouse transgenic aspen plants showed a disease resistance to fungi such as those which induce leaf-blight disease.

#### I. Promoters

Two distinct genes encoding CCL and their promoters were cloned. The promoter of PtCCL1 can drive gene expression specifically in xylem tissue and the promoter for PtCCL2 confers gene expression exclusively in the epidermal tissues. These promoters can be used to manipulate gene expression to engineer traits of interest in specific tissues of target plants. The significance of the promoters is the application of the xylem-specific promoter to direct the expression of any relevant genes specifically in the xylem for engineering lignin content, lignin structure, enhanced tree growth, cellulose content and other value-added wood qualities, etc. The importance of the epidermis-specific promoter is its ability to drive the expression of any relevant genes specifically in epidermal tissues for engineering disease-, UV light-, cold-, heat-, drought-, and other stress resistance traits in trees.

Specifically, the promoters of the PtCCL1 and PtCCL2 were conventionally isolated as follows. An aspen genomic library was screened with PtCCL1cDNA and PtCCL2 partial cDNA fragment to isolate genomic clones of PtCCL1 and PtCCL2. Eleven and seven positive genomic clones were identified for PtCCL1 and PtCCL2 gene, respectively. Among 11 positive clones for PtCCL1, PtCCL1g-4 contained a full length coding sequence and at least 2 kb 5' flanking regions. The restriction map of PtCCL1g-4 is set forth at FIG. 3.

With respect to PtCCL2, restriction map analysis was performed on  $\lambda$ DNA of positive genomic clone PtCCL2g-11. The restriction map of PtCCL2g-11 is set forth at FIG. 4.

Approximately a 2.3 kb 5' flanking region of PtCCL1 was digested from PtCCL1g-4 using Xba I and Sac I sites and cloned into pGEM7Z Xba I and Sac I sites. The subcloned PtCCL1 promoter was named p7Z-4XS and the restriction map of p7Z-4XS is set forth at FIG. 5. The 5' unilateral deletion of p7Z-4XS was generated for DNA sequencing by exonuclease III/S1 nuclease digestion using Erase-a-Base System (Promega, Madison, Wis.). The deletion series was sequenced using a primer on pGEM7Z vector.

A 1.6 kb Hind III and EcoR I fragment containing a 1.2 kb 5' flanking region of PtCCL2 and 0.4 kb coding region of PtCCL2g-11 were subcloned in pBluescript II SK+ Hind III and EcoR I sites. The restriction map of the resulting clone, pSK-11HE, was determined by digesting the plasmid with several restriction enzymes, as in set forth at FIG. 6. In order



to determine the sequence of the PtCCL2 promoter, pSK-11HE was further digested into small fragments according to the restriction map and subcloned into vectors with suitable cloning sites. The DNA sequence was determined using M13 universal primer and reverse primer on the vector.

The DNA sequences of the two promoters was determined and analyzed using  $\Delta$ Taq cycle sequencing Kit (USB, Cleveland, Ohio), and GENETYX-MAC 7.3 sequence analysis software from Software Development Co., Ltd. The nucleotide sequence of promoter region of PtCCL1 is set forth as SEQ ID NO:5 and the nucleotide sequence of the promoter region of PtCCL2 is set forth as SEQ ID NO:6. The promoter gene constructs PtCCL1p and PtCCL2p are available from Michigan Technological University, Institute of Wood Research, Houghton, Mich.

Tissue-specific expression can be achieved by conventionally fusing the promoters of PtCCL1 or PtCCL2 to a gene of interest and transferred to a plant species via *Agrobacterium*. For the sake of example, the promoters of PtCCL1 and PtCCL2 were fused to a GUS reporter gene as detailed below. However, it should be noted that genes other than the GUS reporter gene can be fused to these promoters for tissue specific expression.

In order to construct PtCCL1 promoter-GUS binary vector, a 1 Kb fragment covering 5'-flanking region and 117 bp coding region of PtCCL1 was subcloned into pGEM7Z Sph I and EcoR I sites for constructing promoter-GUS binary vector. In this 1 kb DNA fragment, it is found that one Xho I site locates at 486 bases proximal to the translation start site and the EcoR I site at 117 bases downstream the translation site. This 0.6 Kb fragment was subcloned into pGEM7Z Xho I and EcoR I sites and used as a template in PCR amplification.

In order to construct a promoter-GUS transcriptional fusion, a BamH I site was introduced in front of the translation start site of PtCCL1 by PCR. PCR amplification was performed using p7Z-4XE as the template, M13 universal primer on pGEM7Z vector as 5' end primer and PtCCL1p-1 primer containing a BamH I site at the end is complementary to a sequence upstream of the translation start site. The reaction was carried out in 100  $\mu$ l reaction mix containing 1 $\times$ pfu reaction buffer, 200  $\mu$ l each dNTPs, 100  $\mu$ M each primer and 5 units of pfu. The PCR reaction mixture was denatured at 94 $^{\circ}$  C. for 5 minutes followed by 30 cycles of 94 $^{\circ}$  C. (1 minute), 55 $^{\circ}$  C. (1 minute), 72 $^{\circ}$  C. (1 minute, 30 seconds) and was ended with a 5 minute extension at 72 $^{\circ}$  C.

The amplified 0.6 Kb fragment was cloned and sequenced to confirm the sequence. The engineered 0.6 Kb fragment was ligated to p7Z-4SE which was digested with Xho I and BamH I. In order to incorporate a Hind III site in the 5' end of PtCCL1 promoter, the 1 kb Sph I-BamH I PtCCL1promoter region was the cloned into pNoTA (5 prime $\rightarrow$ 3 prime Inc., Boulder, Colo.) Sph I and BamH I site. The 1 Kb PtCCL1promoter was then released from pNoTA vector with Hind III and BamH digestion and subsequently transcriptionally fused to pBI101 Hind III and BamH I sites in front of GUS. The resulting binary vector was named PtCCL1p-GUS and is set forth at FIG. 7.

In order to construct PtCCL2 promoter-GUS binary vector, pSK-11HE was digested with Sph I and EcoR I to release 0.2 Kb Sph I and EcoR I fragment. The 0.2 Kb fragment was cloned into pGEM7Z Sph I and EcoR I sites. A primer, PtCCL2p-3' (5'-CATCGGATCCTGAGATGGAAGGGAGTTTCT-3')(SEQ ID NO:14) was designed to be complementary to a sequence upstream of the translation start site of PtCCL2 and to incorporate BamH I site at the end. Amplification was performed using p7Z11SE as a template, M13 universal primer as the 5' end primer and PtCCL2p-3 as the 3' end primer. A PCR reaction was carried out and the amplified PCR product was cloned and sequenced to check the fidelity of the PCR amplification. The 0.2 Kb Sph I-BamH I DNA fragment with correct sequence was fused to pSK-11HE linearized with Sph I and BamH I. The resulting plasmid was named pSK-11HB. The promoter of PtCCL2 was then excised from pSK-11HB with Hind III and BamH I and ligated to pBI101 Hind III and BamH I site to make PtCCL2p-GUS transcriptional fusion binary vector as shown in FIG. 8.

The PtCCL1p-GUS and PtCCL2p-GUS constructs are then mobilized into *Agrobacterium tumefaciens* strain C58/pMP90 by freeze and thaw method as explained previously.

Leaf disk transformation of tobacco with these two *Agrobacterium* constructs is conducted according to the method of Horsch R. B. (1988) Leaf Disk Transformation, Plant Molecular Biology Manual, A5:1-9. Histochemical GUS staining of promoter-GUS transgenic tobacco plants demonstrated that the PtCCL1 promoter restricted GUS expression in xylem tissue whereas PtCCL2 promoter regulated GUS expression in epidermal cells.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1  
<211> LENGTH: 1915  
<212> TYPE: DNA  
<213> ORGANISM: *Populus tremuloides* Michx.  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (83)...(1687)

<400> SEQUENCE: 1

```
ccctcgcgaa actccgaaaa cagagagcac ctaaaactca coactctctcc ctctgcatct      60
ttagcccgca atggagcgca ca atg aat cca caa gaa ttc atc ttt cgc tca      112
Met Asn Pro Gln Glu Phe Ile Phe Arg Ser
1 5 10
```

-continued

aaa tta cca gac atc tac atc ccg aaa aac ctt ccc ctg cat tca tac	160
Lys Leu Pro Asp Ile Tyr Ile Pro Lys Asn Leu Pro Leu His Ser Tyr	
15 20 25	
gtt ctt gag aac ttg tct aaa cat tca tca aaa cct tgc ctg ata aat	208
Val Leu Glu Asn Leu Ser Lys His Ser Ser Lys Pro Cys Leu Ile Asn	
30 35 40	
ggc gcg aat gga gat gtc tac acc tat gct gat gtt gag ctc aca gca	256
Gly Ala Asn Gly Asp Val Tyr Thr Tyr Ala Asp Val Glu Leu Thr Ala	
45 50 55	
aga aga gtt gct tct ggt ctg aac aag att ggt att caa caa ggt gac	304
Arg Arg Val Ala Ser Gly Leu Asn Lys Ile Gly Ile Gln Gln Gly Asp	
60 65 70	
gtg atc atg ctc ttc cta cca agt tca cct gaa ttc gtg ctt gct ttc	352
Val Ile Met Leu Phe Leu Pro Ser Ser Pro Glu Phe Val Leu Ala Phe	
75 80 85 90	
cta ggc gct tca cac aga ggt gcc atg atc act gct gcc aat cct ttc	400
Leu Gly Ala Ser His Arg Gly Ala Met Ile Thr Ala Ala Asn Pro Phe	
95 100 105	
tcc acc cct gca gag cta gca aaa cat gcc aag gcc tcg aga gca aag	448
Ser Thr Pro Ala Glu Leu Ala Lys His Ala Lys Ala Ser Arg Ala Lys	
110 115 120	
ctt ctg ata aca cag gct tgt tac tac gag aag gtt aaa gat ttt gcc	496
Leu Leu Ile Thr Gln Ala Cys Tyr Tyr Glu Lys Val Lys Asp Phe Ala	
125 130 135	
cga gaa agt gat gtt aag gtc atg tgc gtg gac tct gcc ccg gac ggt	544
Arg Glu Ser Asp Val Lys Val Met Cys Val Asp Ser Ala Pro Asp Gly	
140 145 150	
gct tca ctt ttc aga gct cac aca cag gca gac gaa aat gaa gtg cct	592
Ala Ser Leu Phe Arg Ala His Thr Gln Ala Asp Glu Asn Glu Val Pro	
155 160 165 170	
cag gtc gac att agt cct gat gat gtc gta gca ttg cct tat tca tca	640
Gln Val Asp Ile Ser Pro Asp Asp Val Val Ala Leu Pro Tyr Ser Ser	
175 180 185	
ggg act aca ggg ttg cca aaa ggg gtc atg tta acg cac aaa ggg cta	688
Gly Thr Thr Gly Leu Pro Lys Gly Val Met Leu Thr His Lys Gly Leu	
190 195 200	
ata acc agt gtg gct caa cag gta gat gga gac aat cct aac ctg tat	736
Ile Thr Ser Val Ala Gln Gln Val Asp Gly Asp Asn Pro Asn Leu Tyr	
205 210 215	
ttt cac agt gaa gat gtg att ctg tgt gtg ctt cct atg ttc cat atc	784
Phe His Ser Glu Asp Val Ile Leu Cys Val Leu Pro Met Phe His Ile	
220 225 230	
tat gct ctg aat tca atg atg ctc tgt ggt ctg aga gtt ggt gcc tcg	832
Tyr Ala Leu Asn Ser Met Met Leu Cys Gly Leu Arg Val Gly Ala Ser	
235 240 245 250	
att ttg ata atg cca aag ttt gag att ggt tct ttg ctg gga ttg att	880
Ile Leu Ile Met Pro Lys Phe Glu Ile Gly Ser Leu Leu Gly Leu Ile	
255 260 265	
gag aag tac aag gta tct ata gca cca gtt gtt cca cct gtg atg atg	928
Glu Lys Tyr Lys Val Ser Ile Ala Pro Val Val Pro Pro Val Met Met	
270 275 280	
gca att gct aag tca cct gat ctt gac aag cat gac ctg tct tct ttg	976
Ala Ile Ala Lys Ser Pro Asp Leu Asp Lys His Asp Leu Ser Ser Leu	
285 290 295	
agg atg ata aaa tct gga ggg gct cca ttg ggc aag gaa ctt gaa gat	1024
Arg Met Ile Lys Ser Gly Gly Ala Pro Leu Gly Lys Glu Leu Glu Asp	
300 305 310	
act gtc aga gct aag ttt cct cag gct aga ctt ggt cag gga tat gga	1072
Thr Val Arg Ala Lys Phe Pro Gln Ala Arg Leu Gly Gln Gly Tyr Gly	
315 320 325 330	

-continued

---

```

atg acc gag gca gga cct gtt cta gca atg tgc ttg gca ttt gcc aag      1120
Met Thr Glu Ala Gly Pro Val Leu Ala Met Cys Leu Ala Phe Ala Lys
          335                      340                      345

gaa cca ttc gac ata aaa cca ggt gca tgt gga act gta gtc agg aat      1168
Glu Pro Phe Asp Ile Lys Pro Gly Ala Cys Gly Thr Val Val Arg Asn
          350                      355                      360

gca gag atg aag att gtt gac cca gaa aca ggg gtc tct cta ccg agg      1216
Ala Glu Met Lys Ile Val Asp Pro Glu Thr Gly Val Ser Leu Pro Arg
          365                      370                      375

aac cag cct ggt gag atc tgc atc cgg ggt gat cag atc atg aaa gga      1264
Asn Gln Pro Gly Glu Ile Cys Ile Arg Gly Asp Gln Ile Met Lys Gly
          380                      385                      390

tat ctt aat gac ccc gag gca acc tca aga aca ata gac aaa gaa gga      1312
Tyr Leu Asn Asp Pro Glu Ala Thr Ser Arg Thr Ile Asp Lys Glu Gly
          395                      400                      405                      410

tgg ctg cac aca ggc gat atc ggc tac att gat gat gat gat gag ctt      1360
Trp Leu His Thr Gly Asp Ile Gly Tyr Ile Asp Asp Asp Asp Glu Leu
          415                      420                      425

ttc atc gtt gac aga ttg aag gaa ttg atc aag tat aaa ggg ttt cag      1408
Phe Ile Val Asp Arg Leu Lys Glu Leu Ile Lys Tyr Lys Gly Phe Gln
          430                      435                      440

gtt gct cct act gaa ctc gaa gct ttg tta ata gcc cat cca gag ata      1456
Val Ala Pro Thr Glu Leu Glu Ala Leu Leu Ile Ala His Pro Glu Ile
          445                      450                      455

tcc gat gct gct gta gta gga ttg aaa gat gag gat gcg gga gaa gtt      1504
Ser Asp Ala Ala Val Val Gly Leu Lys Asp Glu Asp Ala Gly Glu Val
          460                      465                      470

cct gtt gca ttt gta gtg aaa tca gaa aag tct cag gcc acc gaa gat      1552
Pro Val Ala Phe Val Val Lys Ser Glu Lys Ser Gln Ala Thr Glu Asp
          475                      480                      485                      490

gaa att aag cag tat att tca aaa cag gtg atc ttc tac aag aga ata      1600
Glu Ile Lys Gln Tyr Ile Ser Lys Gln Val Ile Phe Tyr Lys Arg Ile
          495                      500                      505

aaa cga gtt ttc ttc att gaa gca att ccc aag gca cca tca ggc aag      1648
Lys Arg Val Phe Phe Ile Glu Ala Ile Pro Lys Ala Pro Ser Gly Lys
          510                      515                      520

atc ctg agg aag aat ctg aaa gag aag ttg cca ggc ata taactgaaga      1697
Ile Leu Arg Lys Asn Leu Lys Glu Lys Leu Pro Gly Ile
          525                      530                      535

tggtactgaa catttaacc tctgtcttat ttctttaata cttgcgaatc attgtagtgt      1757

tgaaccaagc atgcttgaa aagacacgta cccaacgtaa gacagttact gttcctagta      1817

tacaagctct ttaatgttcg ttttgaactt gggaaaacat aagttctcct gtogccatat      1877

ggagtaattc aattgaatat tttggtttct ttaatgat      1915

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 535

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Populus tremuloides Michx.

&lt;400&gt; SEQUENCE: 2

```

Met Asn Pro Gln Glu Phe Ile Phe Arg Ser Lys Leu Pro Asp Ile Tyr
 1          5          10          15

Ile Pro Lys Asn Leu Pro Leu His Ser Tyr Val Leu Glu Asn Leu Ser
 20          25          30

Lys His Ser Ser Lys Pro Cys Leu Ile Asn Gly Ala Asn Gly Asp Val
 35          40          45

Tyr Thr Tyr Ala Asp Val Glu Leu Thr Ala Arg Arg Val Ala Ser Gly

```

-continued

50					55					60						
Leu	Asn	Lys	Ile	Gly	Ile	Gln	Gln	Gly	Asp	Val	Ile	Met	Leu	Phe	Leu	80
65					70					75						
Pro	Ser	Ser	Pro	Glu	Phe	Val	Leu	Ala	Phe	Leu	Gly	Ala	Ser	His	Arg	95
				85					90							
Gly	Ala	Met	Ile	Thr	Ala	Ala	Asn	Pro	Phe	Ser	Thr	Pro	Ala	Glu	Leu	110
			100					105					110			
Ala	Lys	His	Ala	Lys	Ala	Ser	Arg	Ala	Lys	Leu	Leu	Ile	Thr	Gln	Ala	125
		115					120						125			
Cys	Tyr	Tyr	Glu	Lys	Val	Lys	Asp	Phe	Ala	Arg	Glu	Ser	Asp	Val	Lys	140
	130					135					140					
Val	Met	Cys	Val	Asp	Ser	Ala	Pro	Asp	Gly	Ala	Ser	Leu	Phe	Arg	Ala	160
145					150					155					160	
His	Thr	Gln	Ala	Asp	Glu	Asn	Glu	Val	Pro	Gln	Val	Asp	Ile	Ser	Pro	175
				165					170							
Asp	Asp	Val	Val	Ala	Leu	Pro	Tyr	Ser	Ser	Gly	Thr	Thr	Gly	Leu	Pro	190
			180					185						190		
Lys	Gly	Val	Met	Leu	Thr	His	Lys	Gly	Leu	Ile	Thr	Ser	Val	Ala	Gln	205
		195					200						205			
Gln	Val	Asp	Gly	Asp	Asn	Pro	Asn	Leu	Tyr	Phe	His	Ser	Glu	Asp	Val	220
		210					215						220			
Ile	Leu	Cys	Val	Leu	Pro	Met	Phe	His	Ile	Tyr	Ala	Leu	Asn	Ser	Met	240
225					230					235					240	
Met	Leu	Cys	Gly	Leu	Arg	Val	Gly	Ala	Ser	Ile	Leu	Ile	Met	Pro	Lys	255
			245					250						255		
Phe	Glu	Ile	Gly	Ser	Leu	Leu	Gly	Leu	Ile	Glu	Lys	Tyr	Lys	Val	Ser	270
			260					265						270		
Ile	Ala	Pro	Val	Val	Pro	Pro	Val	Met	Met	Ala	Ile	Ala	Lys	Ser	Pro	285
		275					280						285			
Asp	Leu	Asp	Lys	His	Asp	Leu	Ser	Ser	Leu	Arg	Met	Ile	Lys	Ser	Gly	300
	290					295					300					
Gly	Ala	Pro	Leu	Gly	Lys	Glu	Leu	Glu	Asp	Thr	Val	Arg	Ala	Lys	Phe	320
305					310					315					320	
Pro	Gln	Ala	Arg	Leu	Gly	Gln	Gly	Tyr	Gly	Met	Thr	Glu	Ala	Gly	Pro	335
				325					330					335		
Val	Leu	Ala	Met	Cys	Leu	Ala	Phe	Ala	Lys	Glu	Pro	Phe	Asp	Ile	Lys	350
			340					345						350		
Pro	Gly	Ala	Cys	Gly	Thr	Val	Val	Arg	Asn	Ala	Glu	Met	Lys	Ile	Val	365
		355					360						365			
Asp	Pro	Glu	Thr	Gly	Val	Ser	Leu	Pro	Arg	Asn	Gln	Pro	Gly	Glu	Ile	380
	370					375					380					
Cys	Ile	Arg	Gly	Asp	Gln	Ile	Met	Lys	Gly	Tyr	Leu	Asn	Asp	Pro	Glu	400
385					390					395					400	
Ala	Thr	Ser	Arg	Thr	Ile	Asp	Lys	Glu	Gly	Trp	Leu	His	Thr	Gly	Asp	415
				405					410					415		
Ile	Gly	Tyr	Ile	Asp	Asp	Asp	Asp	Glu	Leu	Phe	Ile	Val	Asp	Arg	Leu	430
			420					425					430			
Lys	Glu	Leu	Ile	Lys	Tyr	Lys	Gly	Phe	Gln	Val	Ala	Pro	Thr	Glu	Leu	445
		435					440					445				
Glu	Ala	Leu	Leu	Ile	Ala	His	Pro	Glu	Ile	Ser	Asp	Ala	Ala	Val	Val	460
	450					455					460					
Gly	Leu	Lys	Asp	Glu	Asp	Ala	Gly	Glu	Val	Pro	Val	Ala	Phe	Val	Val	480
465					470					475					480	

-continued

Lys Ser Glu Lys Ser Gln Ala Thr Glu Asp Glu Ile Lys Gln Tyr Ile  
 485 490 495

Ser Lys Gln Val Ile Phe Tyr Lys Arg Ile Lys Arg Val Phe Phe Ile  
 500 505 510

Glu Ala Ile Pro Lys Ala Pro Ser Gly Lys Ile Leu Arg Lys Asn Leu  
 515 520 525

Lys Glu Lys Leu Pro Gly Ile  
 530 535

<210> SEQ ID NO 3  
 <211> LENGTH: 1710  
 <212> TYPE: DNA  
 <213> ORGANISM: Populus tremuloides Michx.  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)...(1710)

<400> SEQUENCE: 3

atg atg tcc gtg gcc acg gtt gag ccc ccg aaa ccg gaa ctc tcc cct 48  
 Met Met Ser Val Ala Thr Val Glu Pro Pro Lys Pro Glu Leu Ser Pro  
 1 5 10 15

cca caa aac caa aac gca cca tcc tct cat gaa act gat cac att ttc 96  
 Pro Gln Asn Gln Asn Ala Pro Ser Ser His Glu Thr Asp His Ile Phe  
 20 25 30

aga tca aaa cta cca gac ata acc atc tcg aac gac ctc cct ctg cac 144  
 Arg Ser Lys Leu Pro Asp Ile Thr Ile Ser Asn Asp Leu Pro Leu His  
 35 40 45

gca tac tgc ttt gaa aac ctc tct gat ttc tca gat agg cca tgc ttg 192  
 Ala Tyr Cys Phe Glu Asn Leu Ser Asp Phe Ser Asp Arg Pro Cys Leu  
 50 55 60

att tca ggt tcc acg gga aaa acc tat tct ttt gcc gaa act cac ctc 240  
 Ile Ser Gly Ser Thr Gly Lys Thr Tyr Ser Phe Ala Glu Thr His Leu  
 65 70 75 80

ata tct cgg aag gtc gct gct ggg tta tcc aat ttg ggc atc aag aaa 288  
 Ile Ser Arg Lys Val Ala Ala Gly Leu Ser Asn Leu Gly Ile Lys Lys  
 85 90 95

ggc gat gta atc atg acc ctg ctc caa aac tgc cca gaa ttc gtc ttc 336  
 Gly Asp Val Ile Met Thr Leu Leu Gln Asn Cys Pro Glu Phe Val Phe  
 100 105 110

tcc ttc atc ggt gct tcc atg att ggt gca gtc atc acc act gcg aac 384  
 Ser Phe Ile Gly Ala Ser Met Ile Gly Ala Val Ile Thr Thr Ala Asn  
 115 120 125

cct ttc tac act caa agt gaa ata ttc aag caa ttc tct gct tct cgt 432  
 Pro Phe Tyr Thr Gln Ser Glu Ile Phe Lys Gln Phe Ser Ala Ser Arg  
 130 135 140

gcg aaa ctg att atc acc cag tct caa tat gtg aac aag cta gga gat 480  
 Ala Lys Leu Ile Ile Thr Gln Ser Gln Tyr Val Asn Lys Leu Gly Asp  
 145 150 155 160

agt gat tgc cat gaa aac aac caa aaa ccg ggg gaa gat ttc ata gta 528  
 Ser Asp Cys His Glu Asn Asn Gln Lys Pro Gly Glu Asp Phe Ile Val  
 165 170 175

atc acc att gat gac ccg cca gag aac tgt cta cat ttc aat gtg ctt 576  
 Ile Thr Ile Asp Asp Pro Pro Glu Asn Cys Leu His Phe Asn Val Leu  
 180 185 190

gtc gag gct agc gag agt gaa atg cca aca gtt tca atc ctt ccg gat 624  
 Val Glu Ala Ser Glu Ser Glu Met Pro Thr Val Ser Ile Leu Pro Asp  
 195 200 205

gat cct gtg gca tta cca ttc tct tca ggg aca aca ggg ctc cca aaa 672  
 Asp Pro Val Ala Leu Pro Phe Ser Ser Gly Thr Thr Gly Leu Pro Lys  
 210 215 220

-continued

gga gtg ata ctg acc cac aag agc ttg ata aca agt gtg gct caa caa	720
Gly Val Ile Leu Thr His Lys Ser Leu Ile Thr Ser Val Ala Gln Gln	
225 230 235 240	
gtt gat gga gag atc cca aat tta tac ttg aaa caa gat gac gtt gtt	768
Val Asp Gly Glu Ile Pro Asn Leu Tyr Leu Lys Gln Asp Asp Val Val	
245 250 255	
tta tgc gtt tta cct ttg ttt cac atc ttt tca ttg aac agc gtg ttg	816
Leu Cys Val Leu Pro Leu Phe His Ile Phe Ser Leu Asn Ser Val Leu	
260 265 270	
tta tgc tcg ttg aga gcc ggt tct gct gtt ctt tta atg caa aag ttt	864
Leu Cys Ser Leu Arg Ala Gly Ser Ala Val Leu Leu Met Gln Lys Phe	
275 280 285	
gag ata gga tca ctg cta gag ctc att cag aaa cac aat gtt tcg gtt	912
Glu Ile Gly Ser Leu Leu Glu Leu Ile Gln Lys His Asn Val Ser Val	
290 295 300	
gcg gct gtg gtg cca cca ctg gtg ctg gcg ttg gcc aag aac cca ttg	960
Ala Ala Val Val Pro Pro Leu Val Leu Ala Leu Ala Lys Asn Pro Leu	
305 310 315 320	
gag gcg aac ttc gac ttg agt tcg atc agg gta gtc ctg tca ggg gct	1008
Glu Ala Asn Phe Asp Leu Ser Ser Ile Arg Val Val Leu Ser Gly Ala	
325 330 335	
gcg cca ctg ggg aag gag ctc gag gac gcc ctc agg agc agg gtt cct	1056
Ala Pro Leu Gly Lys Glu Leu Glu Asp Ala Leu Arg Ser Arg Val Pro	
340 345 350	
cag gcc atc ctg gga cag ggt tat ggg atg aca gag gcc ggg cct gtg	1104
Gln Ala Ile Leu Gly Gln Gly Tyr Gly Met Thr Glu Ala Gly Pro Val	
355 360 365	
cta tca atg tgc tta gcc ttt tca aag caa cct ttc cca acc aag tct	1152
Leu Ser Met Cys Leu Ala Phe Ser Lys Gln Pro Phe Pro Thr Lys Ser	
370 375 380	
ggg tcg tgt gga acg gtg gtt aga aac gca gag ctc aag gtc att gac	1200
Gly Ser Cys Gly Thr Val Val Arg Asn Ala Glu Leu Lys Val Ile Asp	
385 390 395 400	
cct gag acc ggt cgc tct ctt ggt tac aac caa cct ggt gaa atc tgc	1248
Pro Glu Thr Gly Arg Ser Leu Gly Tyr Asn Gln Pro Gly Glu Ile Cys	
405 410 415	
atc cgt gga tcc caa atc atg aaa gga tat ttg aat gac gcg gaa gcc	1296
Ile Arg Gly Ser Gln Ile Met Lys Gly Tyr Leu Asn Asp Ala Glu Ala	
420 425 430	
acg gca aac acc ata gac gtt gag ggt tgg ctc cac act gga gat ata	1344
Thr Ala Asn Thr Ile Asp Val Glu Gly Trp Leu His Thr Gly Asp Ile	
435 440 445	
ggt tat gtc gac gac gac gag att ttc att gtt gat aga gtg aag	1392
Gly Tyr Val Asp Asp Asp Asp Glu Ile Phe Ile Val Asp Arg Val Lys	
450 455 460	
gaa atc ata aaa ttc aaa ggc ttc cag gtg ccg cca gcg gag ctt gag	1440
Glu Ile Ile Lys Phe Lys Gly Phe Gln Val Pro Pro Ala Glu Leu Glu	
465 470 475 480	
gct ctc ctt gta aac cac cct tca att gcg gat gcg gct gtt gtt ccg	1488
Ala Leu Leu Val Asn His Pro Ser Ile Ala Asp Ala Ala Val Val Pro	
485 490 495	
caa aaa gac gag gtt gct ggt gaa gtt cct gtc gcg ttt gtg gtc cgc	1536
Gln Lys Asp Glu Val Ala Gly Glu Val Pro Val Ala Phe Val Val Arg	
500 505 510	
tca gat gat ctt gac ctt agt gaa gag gct gta aaa gaa tac att gca	1584
Ser Asp Asp Leu Asp Leu Ser Glu Glu Ala Val Lys Glu Tyr Ile Ala	
515 520 525	
aag cag gtg gtg ttc tac aag aaa ctg cac aag gtg ttc ttc gtt cat	1632
Lys Gln Val Val Phe Tyr Lys Lys Leu His Lys Val Phe Phe Val His	

-continued

530	535	540	
tct att ccc aaa tcg gct	tct gga aag att	cta aga aaa gac ctc aga	1680
Ser Ile Pro Lys Ser Ala	Ser Gly Lys Ile Leu	Arg Lys Asp Leu Arg	
545	550	555 560	
gcc aag ctt gcc aca gcc	acc acc atg tcc		1710
Ala Lys Leu Ala Thr Ala	Thr Thr Met Ser		
	565	570	
<210> SEQ ID NO 4			
<211> LENGTH: 570			
<212> TYPE: PRT			
<213> ORGANISM: Populus tremuloides Michx.			
<400> SEQUENCE: 4			
Met Met Ser Val Ala Thr	Val Glu Pro Pro Lys Pro	Glu Leu Ser Pro	
1	5	10 15	
Pro Gln Asn Gln Asn Ala	Pro Ser Ser His Glu Thr	Asp His Ile Phe	
	20	25 30	
Arg Ser Lys Leu Pro Asp	Ile Thr Ile Ser Asn Asp	Leu Pro Leu His	
	35	40 45	
Ala Tyr Cys Phe Glu Asn	Leu Ser Asp Phe Ser Asp	Arg Pro Cys Leu	
	50	55 60	
Ile Ser Gly Ser Thr Gly	Lys Thr Tyr Ser Phe Ala	Glu Thr His Leu	
65	70	75 80	
Ile Ser Arg Lys Val Ala	Ala Ala Gly Leu Ser Asn	Leu Gly Ile Lys Lys	
	85	90 95	
Gly Asp Val Ile Met Thr	Leu Leu Gln Asn Cys Pro	Glu Phe Val Phe	
	100	105 110	
Ser Phe Ile Gly Ala Ser	Met Ile Gly Ala Val Ile	Thr Thr Ala Asn	
	115	120 125	
Pro Phe Tyr Thr Gln Ser	Glu Ile Phe Lys Gln Phe	Ser Ala Ser Arg	
	130	135 140	
Ala Lys Leu Ile Ile Thr	Gln Ser Gln Tyr Val Asn	Lys Leu Gly Asp	
145	150	155 160	
Ser Asp Cys His Glu Asn	Asn Gln Lys Pro Gly Glu	Asp Phe Ile Val	
	165	170 175	
Ile Thr Ile Asp Asp Pro	Pro Glu Asn Cys Leu His	Phe Asn Val Leu	
	180	185 190	
Val Glu Ala Ser Glu Ser	Glu Met Pro Thr Val Ser	Ile Leu Pro Asp	
	195	200 205	
Asp Pro Val Ala Leu Pro	Phe Ser Ser Gly Thr Thr	Gly Leu Pro Lys	
	210	215 220	
Gly Val Ile Leu Thr His	Lys Ser Leu Ile Thr Ser	Val Ala Gln Gln	
225	230	235 240	
Val Asp Gly Glu Ile Pro	Asn Leu Tyr Leu Lys	Gln Asp Asp Val Val	
	245	250 255	
Leu Cys Val Leu Pro Leu	Phe His Ile Phe Ser Leu	Asn Ser Val Leu	
	260	265 270	
Leu Cys Ser Leu Arg Ala	Gly Ser Ala Val Leu Leu	Met Gln Lys Phe	
	275	280 285	
Glu Ile Gly Ser Leu Leu	Glu Leu Ile Gln Lys His	Asn Val Ser Val	
	290	295 300	
Ala Ala Val Val Pro Pro	Leu Val Leu Ala Leu Ala	Lys Asn Pro Leu	
305	310	315 320	
Glu Ala Asn Phe Asp Leu	Ser Ser Ile Arg Val Val	Leu Ser Gly Ala	

-continued

	325		330		335										
Ala	Pro	Leu	Gly	Lys	Glu	Leu	Glu	Asp	Ala	Leu	Arg	Ser	Arg	Val	Pro
			340					345					350		
Gln	Ala	Ile	Leu	Gly	Gln	Gly	Tyr	Gly	Met	Thr	Glu	Ala	Gly	Pro	Val
		355					360						365		
Leu	Ser	Met	Cys	Leu	Ala	Phe	Ser	Lys	Gln	Pro	Phe	Pro	Thr	Lys	Ser
	370					375					380				
Gly	Ser	Cys	Gly	Thr	Val	Val	Arg	Asn	Ala	Glu	Leu	Lys	Val	Ile	Asp
385					390					395					400
Pro	Glu	Thr	Gly	Arg	Ser	Leu	Gly	Tyr	Asn	Gln	Pro	Gly	Glu	Ile	Cys
				405					410					415	
Ile	Arg	Gly	Ser	Gln	Ile	Met	Lys	Gly	Tyr	Leu	Asn	Asp	Ala	Glu	Ala
			420					425					430		
Thr	Ala	Asn	Thr	Ile	Asp	Val	Glu	Gly	Trp	Leu	His	Thr	Gly	Asp	Ile
		435					440					445			
Gly	Tyr	Val	Asp	Asp	Asp	Asp	Glu	Ile	Phe	Ile	Val	Asp	Arg	Val	Lys
	450					455					460				
Glu	Ile	Ile	Lys	Phe	Lys	Gly	Phe	Gln	Val	Pro	Pro	Ala	Glu	Leu	Glu
465					470					475					480
Ala	Leu	Leu	Val	Asn	His	Pro	Ser	Ile	Ala	Asp	Ala	Ala	Val	Val	Pro
				485					490					495	
Gln	Lys	Asp	Glu	Val	Ala	Gly	Glu	Val	Pro	Val	Ala	Phe	Val	Val	Arg
			500					505					510		
Ser	Asp	Asp	Leu	Asp	Leu	Ser	Glu	Glu	Ala	Val	Lys	Glu	Tyr	Ile	Ala
	515						520					525			
Lys	Gln	Val	Val	Phe	Tyr	Lys	Lys	Leu	His	Lys	Val	Phe	Phe	Val	His
	530					535					540				
Ser	Ile	Pro	Lys	Ser	Ala	Ser	Gly	Lys	Ile	Leu	Arg	Lys	Asp	Leu	Arg
545					550					555					560
Ala	Lys	Leu	Ala	Thr	Ala	Thr	Thr	Met	Ser						
			565					570							

<210> SEQ ID NO 5  
 <211> LENGTH: 1172  
 <212> TYPE: DNA  
 <213> ORGANISM: Populus tremuloides Michx.

<400> SEQUENCE: 5

tgtaggattg	gtggaatggg	atcattccta	atcccttaat	gacggtggca	tgaacacaaa	60
gcaaagagaa	gtaggtgcac	tcctccttta	tatatatata	tatatgcatg	catgaggacc	120
atggctatga	tgaaggttaa	tagaggtagt	tgtgattgag	atatgtccag	cactagtttt	180
ttgttggtgt	gattttctcat	gatgacgcga	aaatittata	tatatatata	atgaataata	240
tgattgatta	ttctctgtaa	ttttgtgaaa	tagattaata	cagctcaatg	tgagggtgacc	300
agttgtcaaa	tgaccactcg	acttggggca	tggtgatttt	tcaaatcaca	actcaatttg	360
aaaactaaaa	ttaaaaaaga	tttagattat	taaattatta	ggtaattca	cgggttggtc	420
aatcaattat	tattaattaa	aacgatagta	tttttgataa	tttaattaa	atttttattg	480
atttgaatga	actcaattac	atcacaaaaa	acctaataca	attaatatct	tatgtgatat	540
aatttagaaa	tataaatgat	taacctttaa	atctcgagtt	tctcttataa	aaaacacgta	600
taattgggct	agatttaaca	gctattattc	aaactggcca	ggacaattat	taaaattaat	660
aattattatt	ttttctaata	aagcacttcc	taattgttaa	aatatatgtc	taaacactaa	720



-continued

---

taataaaatt tatttgtgta tctttggcag taggtgagag gtgctgacaa ataaattagt	780
gcataaaata taatggattg gtggctctgtg aaaagacagg tggaggacaa gccacctctc	840
tcaagtcaaa aggccatttc acaaccaacc caaatgggaa cccaccaccg ttccccgcca	900
ttaaaatccc taatctcacc aaccaactc cacagattct tcaccaaacg caactgattt	960
ttcaatcaat gttttcccta tactaccccc ccaacaactc cataataccc aatttgcct	1020
ttcaccaacc cccgtcctcc gtgccagcca attctatatac agcaggaatg ctctgcaactc	1080
tgctttctca ggtctctac cataagaaaa cagagagcac ctaaaactcg ccatctctcc	1140
ctctgcatct ttagcccgca atggacgca ca	1172

<210> SEQ ID NO 6  
 <211> LENGTH: 1180  
 <212> TYPE: DNA  
 <213> ORGANISM: Populus tremuloides Michx.

<400> SEQUENCE: 6

aagctttgag tattcatatg ggtattcatc cgaccattat tttcaattt gtgttggtt	60
gatccaattt tcaacttatt tttttttcac ttatttttta ttagttattt ttatttttat	120
tattttttta aaaatttaaa aattaaatta taacattttt attttatccc tcattaacta	180
aaatagggat ggtaatagat attcatgaag ggagttatat atcaaatgat attagttaag	240
ctattttgat atttataccc tactcattac ttatggaata aaaaatttag atatttataa	300
aatattttatc ggatttcagg tattcatatg aatatttatt tgattattat ttattcaaca	360
aaaaataaaa caattaatat gcatgtttga agtttatata tatattaagt taggtttaga	420
tagatttttg gtggggttaa ttaatatcca tacctatct actatctatc aaataatcca	480
aataaaatcca cctaaattag gttgggtttg tattcatcaa gttaacatta aattgtaatt	540
ccgtaagtaa ctaaacaagt acaaagact ctattttatc ttatatatta ccataaagcc	600
aactatattt cctattcttt ttcacccctt ctatcgtaat tttctgtgac ttttttattt	660
atatattaac ggtaacgaaa cacagcaata aaagtattg tgaagatat ggataattat	720
tatggtgact atgaaagagt aaatttgcca tgcactaagt tcctagtgtc atctcataaa	780
agacttgtct gccacgtaag ctggttgtag tgtcgtttat ttacgctgt caaccaatcg	840
ctgccaattg actcctgagg gtaggtgaga gcttcggctt tgatgggaac tgcattgaggc	900
ataggggttg gtttcttgaa tgtgagatgg gcatgctttg gctcccttgc tactcacctc	960
atcttcaatt tgccagctca gctaccagtc totcaccact agtttcacca aactttctct	1020
gctcctgtat ttattacacc ttgctcgatt ggctccgtcc tcgtacacgc atccacaccg	1080
atcgatcgat tagaaccata cagaattggg attggttggg tttacattct gcgtagata	1140
catctatcac agaaagaaac tccttccat ctgagaaac	1180

<210> SEQ ID NO 7  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Highly conserved region of amino acids as  
 determined following alignment of cloned plant CCL  
 sequences.

<400> SEQUENCE: 7

Ser Ser Gly Thr Thr Gly Leu Pro Lys Gly Val  
 1 5 10

-continued

---

```

<210> SEQ ID NO 8
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Highly conserved region of amino acids as
determined following alignment of cloned plant CCL
sequences.

```

```

<400> SEQUENCE: 8

```

```

Gly Glu Ile Cys Ile Arg Gly
  1             5

```

```

<210> SEQ ID NO 9
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sense primer designed around the first
consensus AMP-binding region of CCL.

```

```

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (11)...(11)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (14)...(14)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (17)...(17)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)...(20)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (23)...(23)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (26)...(26)
<223> OTHER INFORMATION: I

```

```

<400> SEQUENCE: 9

```

```

ttggatccgg aacaacagga ytaccaaarg g

```

31

```

<210> SEQ ID NO 10
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sense primer designed around the first
consensus AMP-binding region of CCL.

```

```

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (11)...(11)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (14)...(14)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (23)...(23)
<223> OTHER INFORMATION: I

```

```

<400> SEQUENCE: 10

```

```

ttggatccgt agcacarcar gtagaygg

```

28

```

<210> SEQ ID NO 11

```

-continued

---

```

<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Antisense primer designed based on the
sequence of the putative catalytic motif GEICIRG (SEQ ID NO:12).
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: I
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: I

<400> SEQUENCE: 11

atgtcgacca cgdatrcada tytcacc 27

<210> SEQ ID NO 12
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer designed based on the genomic
sequence of PtCCL2.

<400> SEQUENCE: 12

tctgtctaga tgatgtcgtg gccacgg 27

<210> SEQ ID NO 13
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer designed based on the genomic
sequence of PtCCL2.

<400> SEQUENCE: 13

ttagatctct aggacatggt ggtggc 26

<210> SEQ ID NO 14
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A primer designed to be complementary to a
sequence upstream of the translation start site of
PtCCL2 and to incorporate BamH I site at the end.

<400> SEQUENCE: 14

catcggatcc tgagatggaa gggagtttct 30

<210> SEQ ID NO 15
<211> LENGTH: 1927
<212> TYPE: DNA
<213> ORGANISM: Petroselinum crispum
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (64)..(1698)

<400> SEQUENCE: 15

gaattcccat catcgtttca acacaaaaa cacacacaca actcatattt tcatattttc 60

ata atg gga gac tgt gta gca ccc aaa gaa gac ctt att ttc cga tcg 108
Met Gly Asp Cys Val Ala Pro Lys Glu Asp Leu Ile Phe Arg Ser
1 5 10 15

aaa ctc cct gat att tac atc ccg aaa cac ctt ccg tta cat act tat 156
Lys Leu Pro Asp Ile Tyr Ile Pro Lys His Leu Pro Leu His Thr Tyr
20 25 30

```

-continued

tgt ttc gaa aac atc tcg aaa gtt ggc gac aag tcc tgt tta ata aat Cys Phe Glu Asn Ile Ser Lys Val Gly Asp Lys Ser Cys Leu Ile Asn 35 40 45	204
ggc gct aca ggc gaa acg ttc act tat tct caa gtt gag ctc ctt tcc Gly Ala Thr Gly Glu Thr Phe Thr Tyr Ser Gln Val Glu Leu Leu Ser 50 55 60	252
agg aaa gtt gca tca ggg tta aac aaa ctc ggc att caa cag ggc gat Arg Lys Val Ala Ser Gly Leu Asn Lys Leu Gly Ile Gln Gln Gly Asp 65 70 75	300
acc atc atg ctt ttg ctc ccc aac tcc cct gag tat ttt ttc gct ttc Thr Ile Met Leu Leu Leu Pro Asn Ser Pro Glu Tyr Phe Phe Ala Phe 80 85 90 95	348
tta ggc gca tcg tat cgt ggt gca att tct act atg gcc aat ccg ttt Leu Gly Ala Ser Tyr Arg Gly Ala Ile Ser Thr Met Ala Asn Pro Phe 100 105 110	396
ttc act tct gct gag gtg atc aaa cag ctc aaa gca tcc cta gct aag Phe Thr Ser Ala Glu Val Ile Lys Gln Leu Lys Ala Ser Leu Ala Lys 115 120 125	444
ctc ata att acg caa gct tgt tac gta gac aaa gtg aaa gac tac gca Leu Ile Ile Thr Gln Ala Cys Tyr Val Asp Lys Val Lys Asp Tyr Ala 130 135 140	492
gca gag aaa aat ata cag atc att tgc atc gat gat gct cct cag gat Ala Glu Lys Asn Ile Gln Ile Ile Cys Ile Asp Asp Ala Pro Gln Asp 145 150 155	540
tgt tta cat ttc tcc aaa ctt atg gaa gct gat gaa tca gaa atg ccc Cys Leu His Phe Ser Lys Leu Met Glu Ala Asp Glu Ser Glu Met Pro 160 165 170 175	588
gag gta gtg atc gat tca gac gat gtc gtc ggc tta cct tac tca tcg Glu Val Val Ile Asp Ser Asp Asp Val Val Ala Leu Pro Tyr Ser Ser 180 185 190	636
ggc act aca gga cta ccg aaa ggt gtt atg ttg acc cac aaa gga ctt Gly Thr Thr Gly Leu Pro Lys Gly Val Met Leu Thr His Lys Gly Leu 195 200 205	684
gtt act agc gtg gca caa caa gtt gat gga gac aat ccg aat tta tat Val Thr Ser Val Ala Gln Gln Val Asp Gly Asp Asn Pro Asn Leu Tyr 210 215 220	732
atg cat agc gag gat gtg atg atc tgc ata ttg cct ttg ttt cat att Met His Ser Glu Asp Val Met Ile Cys Ile Leu Pro Leu Phe His Ile 225 230 235	780
tat tcg ctt aac gcg gtg ttg tgc tgt gga ctc aga gca ggg gtg acg Tyr Ser Leu Asn Ala Val Leu Cys Cys Gly Leu Arg Ala Gly Val Thr 240 245 250 255	828
atc ttg att atg cag aaa ttt gat att gtg cca ttt ttg gaa ctg ata Ile Leu Ile Met Gln Lys Phe Asp Ile Val Pro Phe Leu Glu Leu Ile 260 265 270	876
cag aaa tat aaa gtt aca att gga ccg ttt gtg cca cca att gtg ttg Gln Lys Tyr Lys Val Thr Ile Gly Pro Phe Val Pro Pro Ile Val Leu 275 280 285	924
gca att gcg aaa agt cca gtg gtg gat aaa tat gac ttg tcg tcg gtg Ala Ile Ala Lys Ser Pro Val Val Asp Lys Tyr Asp Leu Ser Ser Val 290 295 300	972
agg acg gtt atg tct gga gct gct ccg tta ggg aag gag ctt gaa gat Arg Thr Val Met Ser Gly Ala Ala Pro Leu Gly Lys Glu Leu Glu Asp 305 310 315	1020
gct gtt aga gct aag ttt cct aat gcc aaa ctt ggt cag gga tat gga Ala Val Arg Ala Lys Phe Pro Asn Ala Lys Leu Gly Gln Gly Tyr Gly 320 325 330 335	1068
atg aca gag gca ggg cca gtt tta gca atg tgc ctg gcg ttt gca aag Met Thr Glu Ala Gly Pro Val Leu Ala Met Cys Leu Ala Phe Ala Lys 1116	1116

-continued

340	345	350	
gaa cca tac gag atc aaa tcg ggt gcc tgt gga act gtt gtg agg aat Glu Pro Tyr Glu Ile Lys Ser Gly Ala Cys Gly Thr Val Val Arg Asn 355 360 365			1164
gct gaa atg aaa att gtg gat cct gag acc aac gcc tct ctt cca cga Ala Glu Met Lys Ile Val Asp Pro Glu Thr Asn Ala Ser Leu Pro Arg 370 375 380			1212
aac caa cgc gga gag att tgc att cga ggt gac caa att atg aaa ggc Asn Gln Arg Gly Glu Ile Cys Ile Arg Gly Asp Gln Ile Met Lys Gly 385 390 395			1260
tac ctc aat gat cct gaa tca aca agg aca aca ata gac gaa gaa ggc Tyr Leu Asn Asp Pro Glu Ser Thr Arg Thr Thr Ile Asp Glu Glu Gly 400 405 410 415			1308
tgg ttg cac aca gga gat ata ggc ttc att gac gac gat gat gag cta Trp Leu His Thr Gly Asp Ile Gly Phe Ile Asp Asp Asp Asp Glu Leu 420 425 430			1356
ttt att gtt gat aga ctt aag gaa ata atc aaa tac aaa ggc ttc cag Phe Ile Val Asp Arg Leu Lys Glu Ile Ile Lys Tyr Lys Gly Phe Gln 435 440 445			1404
gtt gcc cct gct gaa ctt gaa gct ctg cta ctt act cat cct acc att Val Ala Pro Ala Glu Leu Glu Ala Leu Leu Leu Thr His Pro Thr Ile 450 455 460			1452
tcc gat gct gca gtt gtt ccc atg ata gat gag aaa gca gga gag gtg Ser Asp Ala Ala Val Val Pro Met Ile Asp Glu Lys Ala Gly Glu Val 465 470 475			1500
cct gtg gct ttt gtt gtg aga aca aac ggt ttc acc acc act gag gaa Pro Val Ala Phe Val Val Arg Thr Asn Gly Phe Thr Thr Thr Glu Glu 480 485 490 495			1548
gaa atc aag caa ttc gtc tcg aaa cag gtg gtg ttc tac aag aga ata Glu Ile Lys Gln Phe Val Ser Lys Gln Val Val Phe Tyr Lys Arg Ile 500 505 510			1596
ttt cgt gta ttt ttt gtt gat gca att ccg aaa tca cca tct gga aag Phe Arg Val Phe Phe Val Asp Ala Ile Pro Lys Ser Pro Ser Gly Lys 515 520 525			1644
att ctt cga aag gac ttg aga gca aaa ata gca tcc ggt gat ctt ccc Ile Leu Arg Lys Asp Leu Arg Ala Lys Ile Ala Ser Gly Asp Leu Pro 530 535 540			1692
aaa taa gtaatctcta caaacagaaa tggcataaag ctgaagctgt atgtgtatct Lys			1748
ttacaaagta aattctacct aaaagagctc cgagttgtaa cttgtttgta tattttattt			1808
tttgaatgaa ggaagattta taagatcatg taatcactca tcaaagtta aatatcatca			1868
tttgtatcac tacattcgggt ttttcogatc ataaacattg attttttcat gttaaaagt			1927

We claim:

1. A method of altering a characteristic of a plant comprising the step of incorporating into the genome of the plant a nucleotide sequence encoding p-coumarate Co-enzyme A ligase (4CL), such that when the nucleotide sequence is expressed in the plant, the characteristic of the plant is altered, wherein the characteristic is selected from the group consisting of altered growth, altered lignin content, increased or decreased coniferyl and sinapyl alcohol units in the lignin structure, increased or decreased disease resistance, altered cellulose content and combinations thereof compared to a control plant that is not transformed with the nucleotide sequence.

2. A plant having a characteristic genetically altered through incorporation into the genome of the plant a nucleotide sequence encoding p-coumarate Co-enzyme A ligase

(4CL), such that when the nucleotide sequence is expressed in the plant, the characteristic of the plant is altered, wherein the characteristic is selected from the group consisting of altered growth, altered lignin content, increased or decreased coniferyl and sinapyl alcohol units in the lignin structure, increased or decreased disease resistance, altered cellulose content and combinations thereof compared to a control plant that is not transformed with the nucleotide sequence.

3. The method as set forth in claim 1 wherein the nucleotide sequence is in the anti-sense orientation.

4. The method as set forth in claim 1 wherein the nucleotide sequence is in the sense orientation.

5. The method as set forth in claim 1 wherein the 4CL comprises an AMP-binding region conserved in all 4CL enzymes in the lignin biosynthetic pathway.

6. The method as set forth in claim 1 wherein the nucleotide sequence comprises:

## 41

- (a) the nucleotide sequence of SEQ ID NO:1;
- (b) a nucleotide sequence having at least about 60% sequence identity to SEQ ID NO:1; or
- (c) a nucleotide sequence that is complementary to (a) or (b).
7. The method as set forth in claim 1 wherein the nucleotide sequence has at least 60% identity to the endogenous 4CL gene.
8. The method as set forth in claim 1 wherein the incorporating is by plant transformation.
9. The method as set forth in claim 8 wherein the transformation is Agrobacterium-mediated transformation.
10. The method as set forth in claim 1 wherein the nucleotide sequence is a cDNA.
11. The method as set forth in claim 1 wherein the nucleotide sequence is operably linked to the CaMV 35S promoter.
12. The method as set forth in claim 1 wherein the plant is a tree.
13. The method as set forth in claim 12 wherein the tree is an angiosperm.
14. The method as set forth in claim 12 wherein the tree is a gymnosperm.
15. The method as set forth in claim 1 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in the average internode length.
16. The method as set forth in claim 1 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in plant height.
17. The method as set forth in claim 1 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in plant diameter.
18. The method as set forth in claim 1 wherein the altered characteristic is increased disease resistance and wherein the increased disease resistance is increased fungal pathogen resistance.
19. The method as set forth in claim 1 wherein the nucleotide sequence is contained in a binary vector.
20. The plant as set forth in claim 2 wherein the nucleotide sequence is in the anti-sense orientation.
21. The plant as set forth in claim 2 wherein the nucleotide sequence is in the sense orientation.
22. The plant as set forth in claim 2 wherein the 4CL comprises an AMP-binding region conserved in all 4CL enzymes in the lignin biosynthetic pathway.
23. The plant as set forth in claim 2 wherein the nucleotide sequence comprises:
- (a) the nucleotide sequence of SEQ ID NO:1;

## 42

- (b) a nucleotide sequence having at least about 60% sequence identity to SEQ ID NO:1; or
- (c) a nucleotide sequence that is complementary to (a) or (b).
24. The plant as set forth in claim 2 wherein the nucleotide sequence has at least 60% identity to the endogenous 4CL gene.
25. The plant as set forth in claim 2 wherein the nucleotide sequence is incorporated into the genome of the plant by transformation.
26. The plant as set forth in claim 25 wherein the transformation is Agrobacterium-mediated transformation.
27. The plant as set forth in claim 2 wherein the nucleotide sequence is a cDNA.
28. The plant as set forth in claim 2 wherein the nucleotide sequence is operably linked to the CaMV 35S promoter.
29. The plant as set forth in claim 2 wherein the plant is a tree.
30. The plant as set forth in claim 29 wherein the tree is an angiosperm.
31. The plant as set forth in claim 29 wherein the tree is a gymnosperm.
32. The plant as set forth in claim 2 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in the average internode length.
33. The plant as set forth in claim 2 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in plant height.
34. The plant as set forth in claim 2 wherein the altered characteristic is accelerated growth and wherein the accelerated growth is manifested as an increase in plant diameter.
35. The plant as set forth in claim 2 wherein the altered characteristic is increased disease resistance and wherein the increased disease resistance is increased fungal pathogen resistance.
36. The plant as set forth in claim 2 wherein the nucleotide sequence is contained in a binary vector.
37. The method as set forth in claim 1 wherein the nucleotide sequence encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or a polypeptide comprising an AMP-binding region conserved in all 4CL enzymes in the lignin biosynthetic pathway.
38. The plant as set forth in claim 2 wherein the nucleotide sequence encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or a polypeptide comprising an AMP-binding region conserved in all 4CL enzymes in the lignin biosynthetic pathway.

\* \* \* \* \*